Mitochondrial proteins and complexes in *Leishmania* and *Trypanosoma* involved in U-insertion/deletion RNA editing

LARRY SIMPSON,^{1,2} RUSLAN APHASIZHEV,¹ GUANGHAN GAO,² and XUEDONG KANG²

¹Department of Microbiology, Immunology and Molecular Genetics, University of California, Los Angeles, California 90095, USA

ABSTRACT

A number of mitochondrial proteins have been identified in *Leishmania sp.* and *Trypanosoma brucei* that may be involved in U-insertion/deletion RNA editing. Only a few of these have yet been characterized sufficiently to be able to assign functional names for the proteins in both species, and most have been denoted by a variety of species-specific and laboratory-specific operational names, leading to a terminology confusion both within and outside of this field. In this review, we summarize the present status of our knowledge of the orthologous and unique putative editing proteins in both species and the functional motifs identified by sequence analysis and by experimentation. An online Supplemental sequence database (<http://164.67.60.200/proteins/protsmi1.asp>) is also provided as a research resource.

Keywords: editing; trypanosomes; RNA; L-complex

INTRODUCTION

The identification of proteins putatively involved in U-insertion/deletion RNA editing in trypanosome mitochondria has progressed very rapidly recently, owing to the application of mass spectrometry gene identification methods to components of macromolecular editing complexes isolated by a variety of methods, combined with the availability of partial genome databases for both *Leishmania major* and *Trypanosoma brucei* (<http://www.genedb.org/>). Our 2003 review (Simpson et al. 2003) is already considerably out of date.

Putative editing proteins have been identified both by direct biochemical isolations targeted for known enzymatic or RNA binding activities, and by sequencing of protein components of macromolecular complexes active in in vitro editing. Possibly functional motifs have been identified by sequence analysis but only confirmed by analysis of recombinant proteins in a few cases. Evidence for an involvement in editing has been obtained for several of these proteins by down-regulation of expression using conditional gene disruptions and RNA interference, but even this evidence is frequently difficult to interpret.

THE 20S “EDITOSOME”

A full cycle, although fairly inefficient, of gRNA-dependent in vitro editing activity that sedimented at approximately 20S in glycerol gradients was detected in mitochondrial extracts from procyclic *T. brucei* (Corell et al. 1996; Kable et al. 1996; Seiwert et al. 1996). Enrichment of this editing activity in the Sollner-Webb laboratory using sequential Q Sepharose and DNA cellulose column fractionations of mitochondrial extract led to a complex reported to contain seven major polypeptides; this sedimented at approximately 20S in glycerol gradients and migrated as a single band in a native gel (Rusché et al. 1997). This preparation was capable of both full round U-insertion and U-deletion editing at one or two sites using synthetic pre-edited A6 mRNA and the cognate gRNA, and the efficiency of in vitro editing could be substantially increased by modifications of the mRNA and gRNA sequences (Cruz-Reyes and Sollner-Webb 1996; Cruz-Reyes et al. 1998, 2001, 2002). The polypeptides were operationally labeled Bands I–VII. Bands IV and V proved to be the REL1 and REL2 RNA ligases (Rusché et al. 2001), which had been functionally identified previously as adenylatable proteins (Sabatini and Hajduk 1995; Peris et al. 1997) and Band III, a zinc finger-containing protein (Huang et al. 2002). Down-regulation of Band III by RNAi was lethal and affected in vivo endonucleolytic cleavage and RNA ligation but not addition or removal of uridines. Recognizable nuclease motifs are absent in the Band III protein, indicating these are indirect effects, con-

Reprint requests to: Larry Simpson, Department of Microbiology, Immunology and Molecular Genetics, University of California, Los Angeles, CA 90095, USA; e-mail: simpson@kdna.ucla.edu; fax: (310) 206-8967.

Article and publication are at <http://www.rnajournal.org/cgi/doi/10.1261/rna.5170704>.

sistent with the decrease of the sedimentation value of the entire complex (Huang et al. 2002). The notion that these seven proteins represented the complete set of proteins required for editing (Rusché et al. 1997) has evolved into the realization that a variety of additional components, which are most likely present in substoichiometric amounts or interact transiently, are also required (Aphasizhev et al. 2003a; Panigrahi et al. 2003a). For example, the RET2 TUTase required for U-insertion editing (see below) is not one of these seven polypeptides (B. Sollner-Webb, pers. comm.).

A similar fractionation procedure of in vitro editing activity in the Stuart laboratory involving sequential SP Sepharose, Q Sepharose, and Superose 6 chromatography of mitochondrial extract led to a product that contained more than 20 major polypeptides (Corell et al. 1996; Panigrahi et al. 2001a,b, 2003b). A product of a similar complexity was isolated by immunoprecipitation of the 20S region from a glycerol gradient of mitochondrial extract using an antibody against TbMP63, one of the protein components of the core complex (Sollner-Webb lab Band III). Several non-mitochondrial and mitochondrial metabolic or structural proteins were, however, also identified by mass spectrometric analysis of tryptic peptides derived from the biochemically purified material and the monoclonal antibody affin-

ity-purified material (Panigrahi et al. 2003a), some of which represented major stained bands in SDS-PAGE, most likely caused by high abundance of these proteins in the mitochondrial cell fractions used as starting material.

Approximately 16 polypeptides with mitochondrial targeting signals were identified that appeared by one or more criteria to represent possible components of the editing machinery (see Table 1). These proteins have been given the operational designation of TbMPx in the Stuart laboratory, where x represents the calculated molecular weight of the mitochondrial pre-protein. The criteria included either cosedimentation in the 20S region in glycerol gradients, coimmunoprecipitation by antibodies against specific proteins demonstrated to be essential for editing, double affinity purification using TAP-tagged proteins, or demonstration of an inhibition of RNA editing by conditional down-regulation of expression in vivo. The genetic method is the most definitive especially because major nonediting protein contaminants were present in the biochemically purified and in the antibody-purified fractions. TbMP123, MP103, MP100, MP99, MP90, MP67, MP61, and MP24 represent proteins with mitochondrial signal sequences that were identified by mass spectrometry analysis of purified editing complexes but for which no definite evidence for inclusion in the editing complex was reported (Table 1).

FIGURE 1. L-complex proteins with protein- and nucleic acid-binding motifs. The locations of the tryptic peptide sequences used for the *L. major* gene identifications are indicated by triangles in this and the following figures. All motif identifications in this and the following figures were performed using SMART (<http://smart.embl-heidelberg.de/>) and PFAM (<http://www.sanger.ac.uk/Software/Pfam/index.shtml>). The *e*-values are indicated in parentheses. Mitochondrial target signals were identified by MitoProt (<http://www.mips.biochem.mpg.de/cgi-bin/proj/medgen/mitofilter>), SMART, or SignalP (<http://www.cbs.dtu.dk/services/SignalP/>). The probabilities from MitoProt are given above the figure. Note that the motifs in LC-10 and LC-7b were identified by similarity with the SSB motif in LC-11. All sequences were obtained from the GeneDB *Leishmania major* and *Trypanosoma brucei* databases.

TABLE 1. Putative editing-related proteins in *Leishmania* and *T. brucei*

Operational names		Leishmania			T. brucei									
Simpson lab	Stuart lab	Sollner-Webb lab	Functional names	Motifs	Possible function		L-complex	Gp-complex	MRP-complex	Effect of down-regulation on editing activity		% ID of Lm and Tb	Ref	
					Putative	Exptl. verified				In vivo	In vitro			
LC-1	MP81	Band II		ZnF_C2H2	Structural scaffold	Interacts with RET2 and REL2	TAP, IP, CC			Lethal, A6, ND7, Cyb	U-insertion	27	Panigrahi et al. 2001b; Aphasizhev et al. 2003a	
LC-2	MP100			Exo_endo_phos	3'-5' exo-nucl. of mRNA		TAP, IP, CC					39	Aphasizhev et al. 2003a; Panigrahi et al. 2003b	
LC-3	MP99			Exo_endo_phos	3'-5' exo-nucl. of mRNA		TAP, IP, CC					25	Aphasizhev et al. 2003a; Panigrahi et al. 2003b	
LC-4	MP63	Band III		ZnF_C2H2	Structural scaffold		TAP, IP, CC			Lethal	mRNA cleavage and ligation	36	Panigrahi et al. 2001b; Aphasizhev et al. 2003a	
LC-5	MP46			ZnF_NFX	RNA binding		TAP, IP, CC					19	Aphasizhev et al. 2003a; Panigrahi et al. 2003b	
LC-6a	MP61			RNase III	Cleavage of mRNA		TAP, IP, CC					38	Aphasizhev et al. 2003a; Panigrahi et al. 2003b	
LC-6b	MP57		RET2	3'-TUTase	Addition of Us at editing site	Recomb. protein TUTase activity	TAP, IP, CC			Lethal, ND8, COIII Cyb, COII	U-insertion	67	Aphasizhev et al. 2003a; Ernst et al. 2003	
LC-7a	MP52	Band IV	REL1	Ligase	Ligation of mRNA fragments	Recomb. protein ligase activity	TAP, IP, CC			Lethal, A6, ND7, RPS12		65	McManus et al. 2001; Panigrahi et al. 2001a; Rusché et al. 2001; Schnauffer et al. 2001; Aphasizhev et al. 2003a	

(Continued)

TABLE 1. (Continued)											
Operational names			Possible function				Effect of down-regulation on editing activity				
<i>Leishmania</i>	Stuart lab	Sollner-Webb lab	Functional names	Motifs	Putative	Exptl. verified	L-complex	GP-complex	MRP-complex	% ID of Lm and Tb	Ref
<i>T. brucei</i>											
LC-7b	MP42	Band VI	ZnF_C2H2	Structural scaffold	TAP, IP, CC					42	Panigrahi et al. 2001b; Aphasizhev et al. 2003a
LC-7c	—				TAP, CS					11	Aphasizhev et al. 2003a
LC-8	MP44		RNase III	Cleavage of mRNA	TAP, IP, CC			Lethal, ND7, A6, ND4, RPS12	Disrupts L-complex	41	Aphasizhev et al. 2003a; Panigrahi et al. 2003b
LC-9	MP48	Band V	REL2	Ligase	Ligation of mRNA fragments	Recomb. protein ligase activity	TAP, IP, CC		No pheno-type	64	McManus et al. 2001; Panigrahi et al. 2001a; Aphasizhev et al. 2003a
LC-10	—				TAP						Aphasizhev et al. 2003a
LC-11	MP18	Band VII	SSB	Binds RNA	TAP, IP, CC					44	Panigrahi et al. 2001b; Aphasizhev et al. 2003a
	MP121				IP, CC						Panigrahi et al. 2003b
Lm MP90	MP90		RNase III	Cleavage of mRNA	IP, CC					38	Panigrahi et al. 2003b
	MP103				IP, CC						Panigrahi et al. 2003b
Lm MP67	MP67		RNase III	Cleavage of mRNA	IP, CC					24	Panigrahi et al. 2003b
	MP24				IP, CC						Panigrahi et al. 2003b

(Continued)

(Continued)

TABLE 1. (Continued)

Operational names			Possible function			Effect of down-regulation on editing activity			% ID
<i>Leishmania</i>	Stuart lab	Sollner-Webb lab	Functional names	Motifs	Putative	Exptl. verified	L-complex	GP-complex	
									In vitro
									In vivo
									Ref
Lm RCGm	TbRCGm-			RRM	RNA binding	IP			57
									Vanhamme et al. 1998; Panigrahi et al. 2003a
									Madison-Antenucci et al. 1998
REAP-1					mRNA binding				72
RBP16				Cold shock domain	RNA binding				Pelletier and Read 2003
Lm mHel61 mHE62					Helicase?				Missel et al. 1997
Lm mHE61 mHE62				DEAD box					Aphasizhev et al. 2002
Lm mHE61 mHE62				3'-TUTase	3'-U addition to gRNAs	Recomb. protein activity	CC	IA, TAP	38
Lm mHE61 mHE62									U-insertion, length of 3'-oligo(U) tail
Lp26	gBP21				gRNA-mRNA binding	Recomb. protein annealing		TAP, CC	45
									Koller et al. 1997; Aphasizhev et al. 2003b
Lp28	gBP25				gRNA-mRNA binding	Recomb. protein annealing		TAP, CC	43
									Blom et al. 2001; Aphasizhev et al. 2003b
AP1								TAP	48
									Aphasizhev et al. 2003b
AP2								TAP	59
									Aphasizhev et al. 2003b
AP3								TAP	42
									Aphasizhev et al. 2003b

Abbreviations: IP, protein detected in immunoprecipitation; TAP, protein detected in tandem affinity purification; CC, protein detected in material purified by column chromatography and/or sedimentation; ID, identical amino acids.

FIGURE 2. L-complex proteins with RNase III motifs. Note that the motif in LC-8 was identified by similarity with the RNase III motif in LC-6a. The *L. major* homologs to the *T. brucei* MP90 and MP67 were identified by database mining.

The TbMP52 and TbMP48 proteins (the Sollner-Webb laboratory Bands IV and V) were identified as the two mitochondrial RNA ligases (Sabatini and Hajduk 1995; McManus et al. 2001) and were designated REL1 (RNA Editing Ligase 1) and REL2 (McManus et al. 2001; Panigrahi et al. 2001a; Rusché et al. 2001; Schnauffer et al. 2001). Down-regulation of REL1 was lethal and inhibited editing, but down-regulation of REL2 (Drozdz et al. 2002; Stuart et al. 2002; Gao and Simpson 2003) had no effect on cell growth. Down-regulation of TbMP81 and Band III (= TbMP63) expression by conditional RNAi inhibited *in vivo* editing but not transcription or mRNA stability.

The TbMP57 protein showed a nucleotidyltransferase motif and poly(A) polymerase-associated motifs, and recombinant protein immunoprecipitated from a reticulocyte transcription-translation system (Ernst et al. 2003) or expressed in *Leishmania tarentolae* as a TAP-tagged cytosolic protein by removal of the mitochondrial signal peptide (Aphasizhev et al. 2003c) both exhibited a 3'-TUTase activity that added mainly a single U residue to the 3'-OH of a single-stranded RNA substrate. This protein has been designated RET2 (RNA Editing TUTase 2; Aphasizhev et al. 2003a; Ernst et al. 2003), as suggested by K. Stuart (pers. comm.). Down-regulation of RET2 in procyclic *T. brucei* was lethal and led to a complete inhibition of U-insertion precleaved *in vitro* editing and no effect on U-deletion editing or on the length

of the 3'-oligo[U] tails of the gRNAs, whereas down-regulation of RET1 expression produced a small effect on *in vitro* U-insertion editing and a large effect on the length distribution of the 3'-oligo[U] tails of gRNAs (Aphasizhev et al. 2003c). It was concluded that RET2 is an integral part of a core editing complex and is responsible for the addition of Us to the editing site and RET1 is responsible for the 3' addition of Us to the gRNAs (Aphasizhev et al. 2003c).

One protein was identified that was homologous to the mitochondrial RNA-binding protein, TcRGGm, and was therefore designated TbRGG1, but there is no evidence that this is a component of an isolated editing complex (Vanhamme et al. 1998; Panigrahi et al. 2003a).

The difference in the number of polypeptides obtained in these two laboratories is most likely due to technical differences in isolation and detection methodologies and to differential stability of the complex components. In both cases, the final preparations showed both U-insertion and U-deletion *in vitro* editing activity. There is clearly a fairly stable macromolecular complex composed of ~16 proteins sedimenting at approximately 20S in glycerol gradients, but it may be premature to designate this as the "editosome" (Panigrahi et al. 2003a) because work in the *L. tarentolae* system discussed below has indicated a higher complexity of interacting components of the editing machinery required for functional editing (Aphasizhev et al. 2003a,b).

THE L-COMPLEX

The ligase-containing complex (L-complex) was first identified as a high-molecular-weight band in native gels of *L.*

FIGURE 3. L-complex proteins with nuclease motifs. The LC-3 protein identified by mass spectrometry lacks most of the Exo_endo_phos motif present in MP99, but otherwise is very similar.

FIGURE 4. The *L. major* L-complex RET2 3'-TUTase. The homolog in *T. brucei* was identified by database searching (Aphasizhev et al. 2002) and has been characterized experimentally (Ernst et al. 2003).

tarentolae and *T. brucei* mitochondrial extracts that could be labeled with [α - 32 P]ATP (Sabatini and Hajduk 1995; Peris et al. 1997; Rusché et al. 1997) because of the presence of the two adenylatable RNA ligases, REL1 and REL2 (Sabatini and Hajduk 1995). This complex sedimented in glycerol gradients in a broad region from 20S to 25S but as a single band in a native gel and is almost certainly identical to the "editosome" complexes isolated from *T. brucei* discussed above. The L-complex was isolated from *L. tarentolae* by the TAP procedure using a REL1 fusion protein (Aphasizhev et al. 2003a). The purified L-complex sedimented as a single homogeneous adenylatable 20S particle both in gradients and in native gels. The isolated material showed precleaved U-insertion and U-deletion editing activity. Substoichiometric amounts of RET1 and MRP1 and MRP2 were also present in the final preparation, indicating a low-affinity interaction of the L-complex with the MRP-complex, the GP-complex, and gRNA (see below).

The gradient-isolated TAP-tagged 20S L-complex was composed of 13–16 bands in an SDS gel in approximate stoichiometric amounts (Aphasizhev et al. 2003a). The absence of contamination with abundant mitochondrial metabolic enzymes was shown by probing with antiserum for glutamate dehydrogenase. Substoichiometric amounts of RET1 and the MRP1,2 proteins could be identified by Western analysis. So far, 15 polypeptides have been identified by MS/MS analysis of tryptic peptides from gel-isolated stained bands. All except LC-10 have orthologs in the *T. brucei* database, and orthologs to MP90 and MP67 were identified in the *L. major* database. LC-6b (MP57) was shown to be a 3'-TUTase and was labeled RET2, as mentioned above.

Functional motifs in the L-complex proteins

All identified *Leishmania* editing proteins and the orthologs in *T. brucei* are shown in Table 1, together with the operational designations and the functional gene

names if assigned. The apparent absences of LC-10 in *T. brucei* and MP90, MP103, and MP121 in *Leishmania* are probably due to database gaps rather than species-specific differences. All sequences in FastA format can be obtained from the online spreadsheet at <http://164.67.60.200/proteins/protsmini1>.

asp. Alignments of orthologs from *Leishmania* and *Trypanosoma* with identified motifs indicated are also presented in the online spreadsheet as pdf files.

The presence of the *Leishmania* LC-proteins in the TAP-affinity isolated complex in approximately stoichiometric yields (Aphasizhev et al. 2003a) indicates that these are, indeed, integral components of the L-complex, and that the orthologs in *T. brucei* are also L-complex components in that species.

RNA and/or protein-binding motifs

LC-1 (MP81), LC-4 (MP63), LC-7b (MP42), and LC-5 (MP46) have zinc finger motifs, which could be nucleic acid or protein-binding. LC-11 (MP18) also has an SSB-like single-strand-binding motif (Fig. 1). SSB-like motifs can be recognized in LC-7b (MP42) and LC-10 by alignment of the LC-11 (MP18) SSB motif sequence, as indicated in Figure 1. Only the zinc finger motifs in LC-4 (MP63) have so far been experimentally shown to have a biochemical function in protein–protein interactions (X. Kang and L. Simpson, unpubl.). Possible RNA-binding functions of these motifs remain to be experimentally demonstrated.

RNase III motifs

Three proteins with RNase III motifs were identified from *Leishmania major* and *T. brucei* (Fig. 2). RNase III enzymes degrade double-stranded RNA and are involved in rRNA

FIGURE 5. The L-complex REL1 and REL2 RNA ligases. Five conserved ligase motifs (McManus et al. 2001) are indicated as is the location of the lysine residues that covalently bind AMP. These motifs can be recognized by multiple alignments with known ligase enzymes.

			I * \$ @	III #	IIIa &	IV \$	V * *
T3	DNA	ATP	29	49	55	58	14
T7	DNA	ATP	28	49	49	58	14
Vac	DNA	ATP	226	42	27	50	13
At	DNA	ATP	439	50	25	50	17
Mj	DNA	ATP	245	39	34	66	15
Cj	DNA	ATP	37	31	26	52	14
Sc	DNA	ATP	414	43	32	50	17
Hu	DNA	ATP	201	48	29	50	15
Ec	DNA	NAD	110	48	45	50	23
Ae	DNA	NAD	137	45	45	52	22
Mg	DNA	NAD	108	45	41	49	22
T4	RNA	ATP	94	42	34	25	6
Sc	tRNA	ATP	109	28	36	59	11
ChlV	CE	GTP	77	39	08	57	14
At	CE	GTP	360	43	12	66	14
Tb	CE	GTP	283	56	06	53	13
Sc	CE	GTP	65	52	12	64	15
Ce	CE	GTP	294	38	14	66	14
Hu	CE	GTP	289	41	10	66	13
Lt	REL1	ATP	100	62	43	64	17
Lm	REL1	ATP	103	62	43	64	17
Tb	REL1	ATP	82	64	43	64	17
Lt	REL2	ATP	53	61	40	63	16
Lm	REL2	ATP	53	61	40	63	16
Tb	REL2	ATP	52	61	40	63	16

FIGURE 6. Multiple alignments of portions of selected genes to illustrate the five conserved motifs in REL1 and REL2 characteristic of DNA and RNA ligases (Shuman and Schwer 1995; Odell et al. 2000; McManus et al. 2001). (T3) Bacteriophage T3; (T7) bacteriophage T7; (Vac) Vaccinia virus; (At) *Arabidopsis thaliana*; (Mj) *Methanocaldococcus jannaschii*; (Cj) *Campylobacter jejuni*; (Sc) *Saccharomyces cerevisiae*; (Hu) human; (Ec) *E. coli*; (Ae) *Aquifex aeolicus*; (Mg) *Mycoplasma genitalium*; (T4) bacteriophage T4; (ChlV) Chlorella virus; (Tb) *T. brucei*; (Ce) *Caenorhabditis elegans*; (Lt) *L. tarentolae*; (Lm) *L. major*; (*) α PO₄-binding residues; (\$) metal-binding residues; (@) ribose O-binding residue; (#) residue binding to Arg; (&) adenine-binding residue.

processing and degradation of some mRNAs. The presence of an RNase III-like motif in LC-8 (MP44) was only detected by alignment of the LC-6a (MP61) RNase III motif sequence (Fig. 2). There is as yet no biochemical evidence for enzymatic activity of these proteins. A possible function is the gRNA-mediated cleavage of mRNA at editing sites.

Exonuclease–endonuclease–phosphatase motifs

There are two L-complex proteins, LC-2 (MP100) and Tb MP99, with exonuclease–endonuclease–phosphatase motifs (Fig. 3). The *L. major* LC-3 protein is clearly an ortholog to MP99 but, interestingly, lacks most of the motif sequence.

3'-TUTase motif

The L-complex LC-6b (MP57; Aphasizhev et al. 2003a; Ernst et al. 2003) has a nucleotidyltransferase motif separated between the second and third catalytic aspartates by a stretch of amino acids that is not found in other nucleotidyltransferase proteins, as exemplified by the RET1 TUTase (Fig. 4; Aphasizhev et al. 2002). There are also poly(A) polymerase-associated motifs. This is the ortholog of RET2 in *T. brucei*, which has been expressed and shown to add mainly a single U residue to the 3'-end of RNA substrates (Aphasizhev et al. 2003c; Ernst et al. 2003).

Ligase motifs

The five motifs in the REL1 and REL2 proteins indicated in Figures 5 and 6 are conserved among both DNA and RNA ligases (McManus et al. 2001; Ho and Shuman 2002). The localizations of the lysine residues that covalently bind AMP were also determined experimentally (Huang et al. 2001). Ligase activities of recombinant TbREL1 and REL2 expressed in an *Escherichia coli* system (McManus et al. 2001; Rusché et al. 2001), TbREL1 and TbREL2 expressed in a reticulocyte cell-free coupled transcription–translation system (Palazzo et al. 2003), and LtREL1 and LtREL2 expressed in the *Baculovirus* system (G. Gao and L. Simpson, unpubl.) have been demonstrated.

THE MRP-COMPLEX

Two mitochondrial RNA-binding proteins (MRPs) from *L. tarentolae*, MRP-1, were shown to form a stable 100-kD complex (Aphasizhev et al. 2003b). These mitochondrial proteins, which were previously called Ltp26 and Lt28 (Aphasizhev et al. 2003b), are orthologs of TbgBP21 (Koller et al. 1997) and TbgBP25. We suggest the MRP designation instead of the original gBP (guide RNA binding protein) designation (Koller et al. 1997) because it is now known that these proteins bind RNA fairly nonspecifically (Aphasizhev et al. 2003b). TAP-isolation us-

FIGURE 7. MRP-complex proteins. The *L. major* MRP RNA-binding proteins and the three associated proteins (AP1, AP2, AP3) identified by double affinity isolation (Aphasizhev et al. 2003b) are indicated.

FIGURE 8. RET1 proteins. The *L. major* and *T. brucei* RET1 TUTases are shown (Aphasizhev et al. 2002).

ing an MRP-1 fusion protein was used to identify three additional interacting proteins, AP1, AP2, and AP3 (Fig. 7; Aphasizhev et al. 2003b). The TAP-isolated ~30S complex contained bound gRNA, and the native and recombinant MRP proteins from both species and the reconstituted 100-kD complex exhibited binding to both double- and single-stranded RNA and catalyzed RNA annealing (Muller et al. 2001; Aphasizhev et al. 2003b). We have designated this as the MRP-complex (Aphasizhev et al. 2003c). There is evidence that down-regulation of both MRP1 and MRP2 together affects viability of procyclic *T. brucei* (A. Simpson, R. Aphasizhev, I. Aphasizheva, and L. Simpson, unpubl.), but it remains to be demonstrated that the complex is involved in vivo with catalyzing the annealing of the gRNA and mRNA.

THE RET1-COMPLEX

A biochemical isolation of the major processive 3'-TUTase activity from *L. tarentolae* mitochondria yielded a 121-kD protein belonging to the nucleotidyltransferase superfamily (Fig. 8; Aphasizhev et al. 2002). The recombinant protein expressed in *E. coli* exhibited a U-specific 3' transferase activity that added multiple Us to the 3'-OH of any single-stranded RNA. An involvement with RNA editing was evidenced by the correlation of down-regulation of expression with a decrease in editing in vivo. The homologous protein was identified in *T. brucei*, and it was shown that down-regulation of expression by RNAi led to a decrease in the average length of the 3'-oligo[U] tail of the gRNAs, indicating that the observed inhibition of editing in vivo was indirect. This protein was designated as RET1 (RNA Editing

TUTase 1; Aphasizhev et al. 2003c), as suggested by K. Stuart (pers. comm.).

The majority of the protein was present as an ~500-kD tetramer, but a small portion was in the form of an ~700-kD complex that could be separated by ion exchange chromatography followed by sedimentation in a glycerol gradient or electrophoresis in a native gel. We have designated this as the RET1-complex (Aphasizhev et al. 2003c). It remains to be investigated whether a previously identified U-specific 3'-5' exonuclease activity (Aphasizhev and Simpson 2001; Igo et al. 2002) is also associated with this complex. It is possible that the 700-kD complex is a breakdown product of a larger complex after ion exchange chromatography, but this must be further examined.

PROTEINS WITH POSSIBLE ROLES IN RNA EDITING

It should be noted that there are several additional mitochondrial RNA-binding proteins that interact with guide RNA and mRNA, such as Tb REAP-1 (Madison-Antenucci

FIGURE 9. Several proteins that may be involved in RNA editing. The *L. major* orthologs of the RNA-binding proteins, RBP16 and RGGm, and the putative helicase, mHel61, are shown. The *T. brucei* REAP-1 is shown because there is no ortholog of this protein in the *L. major* database.

bers of the Simpson laboratory for advice and discussion. We acknowledge the contribution of the parasite genome database sequencing centers, the Seattle Biomedical Research Institute, the Sanger Institute, and the Institute for Genomic Research for supplying the genomic sequences and for maintaining the relational databases (<http://www.genedb.org/>) that allowed the identification of most of these proteins.

REFERENCES

- Allen, T.E., Heidmann, S., Reed, R., Myler, P.J., Goring, H.U., and Stuart, K.D. 1998. Association of guide RNA binding protein gBP21 with active RNA editing complexes in *Trypanosoma brucei*. *Mol. Cell. Biol.* **18**: 6014–6022.
- Aphasizhev, R. and Simpson, L. 2001. Isolation and characterization of a U-specific 3′–5′ exonuclease from mitochondria of *Leishmania tarentolae*. *J. Biol. Chem.* **276**: 21280–21284.
- Aphasizhev, R., Sbicego, S., Peris, M., Jang, S.H., Aphasizheva, I., Simpson, A.M., Rivlin, A., and Simpson, L. 2002. Trypanosome mitochondrial 3′ terminal uridylyl transferase (TUTase): The key enzyme in U-insertion/deletion RNA editing. *Cell* **108**: 637–648.
- Aphasizhev, R., Aphasizheva, I., Nelson, R.E., Gao, G., Simpson, A.M., Kang, X., Falick, A.M., Sbicego, S., and Simpson, L. 2003a. Isolation of a U-insertion/deletion editing complex from *Leishmania tarentolae* mitochondria. *EMBO J.* **22**: 913–924.
- Aphasizhev, R., Aphasizheva, I., Nelson, R.E., and Simpson, L. 2003b. A 100-kD complex of two RNA-binding proteins from mitochondria of *Leishmania tarentolae* catalyzes RNA annealing and interacts with several RNA editing components. *RNA* **9**: 62–76.
- Aphasizhev, R., Aphasizheva, I., and Simpson, L. 2003c. A tale of two TUTases. *Proc. Natl. Acad. Sci.* **100**: 10617–10622.
- Blom, D., Burg, J., Breek, C.K., Speijer, D., Muijsers, A.O., and Benne, R. 2001. Cloning and characterization of two guide RNA-binding proteins from mitochondria of *Crithidia fasciculata*: gBP27, a novel protein, and gBP29, the orthologue of *Trypanosoma brucei* gBP21. *Nucleic Acids Res.* **29**: 2950–2962.
- Corell, R.A., Read, L.K., Riley, G.R., Nellissery, J.K., Allen, T.E., Kable, M.L., Wachal, M.D., Seiwert, S.D., Myler, P.J., and Stuart, K.D. 1996. Complexes from *Trypanosoma brucei* that exhibit deletion editing and other editing-associated properties. *Mol. Cell. Biol.* **16**: 1410–1418.
- Cruz-Reyes, J. and Sollner-Webb, B. 1996. Trypanosome U-deletional RNA editing involves guide RNA-directed endonuclease cleavage, terminal U exonuclease, and RNA ligase activities. *Proc. Natl. Acad. Sci.* **93**: 8901–8906.
- Cruz-Reyes, J., Rusché, L.N., and Sollner-Webb, B. 1998. *Trypanosoma brucei* U insertion and U deletion activities co-purify with an enzymatic editing complex but are differentially optimized. *Nucleic Acids Res.* **26**: 3634–3639.
- Cruz-Reyes, J., Zhelonkina, A., Rusche, L., and Sollner-Webb, B. 2001. Trypanosome RNA editing: Simple guide RNA features enhance U deletion 100-fold. *Mol. Cell. Biol.* **21**: 884–892.
- Cruz-Reyes, J., Zhelonkina, A.G., Huang, C.E., and Sollner-Webb, B. 2002. Distinct functions of two RNA ligases in active *Trypanosoma brucei* RNA editing complexes. *Mol. Cell. Biol.* **22**: 4652–4660.
- Drozdz, M., Palazzo, S.S., Salavati, R., O’Rear, J., Clayton, C., and Stuart, K. 2002. TbMP81 is required for RNA editing in *Trypanosoma brucei*. *EMBO J.* **21**: 1791–1799.
- Ernst, N.L., Panicucci, B., Igo Jr., R.P., Panigrahi, A.K., Salavati, R., and Stuart, K. 2003. TbMP57 is a 3′ terminal uridylyl transferase (TUTase) of the *Trypanosoma brucei* editosome. *Mol. Cell* **11**: 1525–1536.
- Gao, G. and Simpson, L. 2003. Is the *trypanosoma brucei* REL1 RNA ligase specific for U-deletion RNA editing and the REL2 RNA ligase specific for U-insertion editing? *J. Biol. Chem.* **278**: 27570–27574.
- Hayman, M.L. and Read, L.K. 1999. *Trypanosoma brucei* RBP16 is a mitochondrial Y-box family protein with guide RNA binding activity. *J. Biol. Chem.* **274**: 12067–12074.
- Hermann, T., Schmid, B., Heumann, H., and Goring, H.U. 1997. A three-dimensional working model for a guide RNA from *Trypanosoma brucei*. *Nucleic Acids Res.* **25**: 2311–2318.
- Ho, C.K. and Shuman, S. 2002. Bacteriophage T4 RNA ligase 2 (gp24.1) exemplifies a family of RNA ligases found in all phylogenetic domains. *Proc. Natl. Acad. Sci.* **99**: 12709–12714.
- Huang, C.E., Cruz-Reyes, J., Zhelonkina, A.G., O’Hearn, S., Wirtz, E., and Sollner-Webb, B. 2001. Roles for ligases in the RNA editing complex of *Trypanosoma brucei*: Band IV is needed for U-deletion and RNA repair. *EMBO J.* **20**: 4694–4703.
- Huang, C.E., O’Hearn, S.F., and Sollner-Webb, B. 2002. Assembly and function of the RNA editing complex in *Trypanosoma brucei* requires band III protein. *Mol. Cell. Biol.* **22**: 3194–3203.
- Igo, Weston, D.S., Ernst, N.L., Panigrahi, A.K., Salavati, R., and Stuart, K. 2002. Role of uridylylate-specific exoribonuclease activity in *Trypanosoma brucei* RNA editing. *Eukar. Cell* **1**: 112–118.
- Kable, M.L., Seiwert, S.D., Heidmann, S., and Stuart, K. 1996. RNA editing: A mechanism for gRNA-specified uridylylate insertion into precursor mRNA. *Science* **273**: 1189–1195.
- Koller, J., Muller, U.F., Schmid, B., Missel, A., Kruff, V., Stuart, K., and Goring, H.U. 1997. *Trypanosoma brucei* gBP21. An arginine-rich mitochondrial protein that binds to guide RNA with high affinity. *J. Biol. Chem.* **272**: 3749–3757.
- Madison-Antenucci, S. and Hajduk, S.L. 2001. RNA editing-associated protein 1 is an RNA binding protein with specificity for preedited mRNA. *Mol. Cell* **7**: 879–886.
- Madison-Antenucci, S., Sabatini, R.S., Pollard, V.W., and Hajduk, S.L. 1998. Kinetoplastid RNA-editing-associated protein 1 (REAP-1): A novel editing complex protein with repetitive domains. *EMBO J.* **17**: 6368–6376.
- McManus, M.T., Shimamura, M., Grams, J., and Hajduk, S.L. 2001. Identification of candidate mitochondrial RNA editing ligases from *Trypanosoma brucei*. *RNA* **7**: 167–175.
- Missel, A. and Goring, H.U. 1994. *Trypanosoma brucei* mitochondria contain RNA helicase activity. *Nucleic Acids Res.* **22**: 4050–4056.
- Missel, A., Nörskau, G., Shu, H.H., and Goring, H.U. 1995. A putative RNA helicase of the DEAD box family from *Trypanosoma brucei*. *Mol. Biochem. Parasitol.* **75**: 123–126.
- Missel, A., Souza, A.E., Nörskau, G., and Goring, H.U. 1997. Disruption of a gene encoding a novel mitochondrial DEAD-box protein in *Trypanosoma brucei* affects edited mRNAs. *Mol. Cell. Biol.* **17**: 4895–4903.
- Muller, U.F., Lambert, L., and Goring, H.U. 2001. Annealing of RNA editing substrates facilitated by guide RNA-binding protein gBP21. *EMBO J.* **20**: 1394–1404.
- Odell, M., Srisakanda, V., Shuman, S., and Nikolov, D.B. 2000. Crystal structure of eukaryotic DNA ligase-adenylate illuminates the mechanism of nick sensing and strand joining. *Mol. Cell* **6**: 1183–1193.
- Palazzo, S.S., Panigrahi, A.K., Igo, R.P., Salavati, R., and Stuart, K. 2003. Kinetoplastid RNA editing ligases: Complex association, characterization, and substrate requirements. *Mol. Biochem. Parasitol.* **127**: 161–167.
- Panigrahi, A.K., Gygi, S.P., Ernst, N.L., Igo, R.P., Palazzo, S.S., Schnauffer, A., Weston, D.S., Carmean, N., Salavati, R., Aebersold, R., et al. 2001a. Association of two novel proteins, TbMP52 and TbMP48, with the *Trypanosoma brucei* RNA editing complex. *Mol. Cell. Biol.* **21**: 380–389.
- Panigrahi, A.K., Schnauffer, A., Carmean, N., Igo Jr., R.P., Gygi, S.P., Ernst, N.L., Palazzo, S.S., Weston, D.S., Aebersold, R., Salavati, R., et al. 2001b. Four related proteins of the *Trypanosoma brucei* RNA editing complex. *Mol. Cell. Biol.* **21**: 6833–6840.
- Panigrahi, A.K., Allen, T.E., Stuart, K., Haynes, P.A., and Gygi, S.P. 2003a. Mass spectrometric analysis of the editosome and other multiprotein complexes in *Trypanosoma brucei*. *J. Am. Soc. Mass Spectrom.* **14**: 728–735.
- Panigrahi, A.K., Schnauffer, A., Ernst, N.L., Wang, B., Carmean, N., Salavati, R., and Stuart, K. 2003b. Identification of novel components of *Trypanosoma brucei* editosomes. *RNA* **9**: 484–492.
- Pelletier, M. and Read, L.K. 2003. RBP16 is a multifunctional gene

- regulatory protein involved in editing and stabilization of specific mitochondrial mRNAs in *Trypanosoma brucei*. *RNA* **9**: 457–468.
- Pelletier, M., Miller, M.M., and Read, L.K. 2000. RNA-binding properties of the mitochondrial Y-box protein RBP16. *Nucleic Acids Res.* **28**: 1266–1275.
- Peris, M., Simpson, A.M., Grunstein, J., Liliental, J.E., Frech, G.C., and Simpson, L. 1997. Native gel analysis of ribonucleoprotein complexes from a *Leishmania tarentolae* mitochondrial extract. *Mol. Biochem. Parasitol.* **85**: 9–24.
- Rappsilber, J., Ryder, U., Lamond, A.I., and Mann, M. 2002. Large-scale proteomic analysis of the human spliceosome. *Genome Res.* **12**: 1231–1245.
- Rusché, L.N., Cruz-Reyes, J., Piller, K.J., and Sollner-Webb, B. 1997. Purification of a functional enzymatic editing complex from *Trypanosoma brucei* mitochondria. *EMBO J.* **16**: 4069–4081.
- Rusché, L.N., Huang, C.E., Piller, K.J., Hemann, M., Wirtz, E., and Sollner-Webb, B. 2001. The two RNA ligases of the *Trypanosoma brucei* RNA editing complex: Cloning the essential band IV gene and identifying the band V gene. *Mol. Cell. Biol.* **21**: 979–989.
- Sabatini, R. and Hajduk, S.L. 1995. RNA ligase and its involvement in guide RNA/mRNA chimera formation. *J. Biol. Chem.* **270**: 7233–7240.
- Schnauffer, A., Panigrahi, A.K., Panicucci, B., Igo, R.P., Salavati, R., and Stuart, K. 2001. An RNA ligase essential for RNA editing and survival of the bloodstream form of *Trypanosoma brucei*. *Science* **291**: 2159–2161.
- Seiwert, S.D., Heidmann, S., and Stuart, K. 1996. Direct visualization of uridylyte deletion in vitro suggests a mechanism for kinetoplast RNA editing. *Cell* **84**: 831–841.
- Shuman, S. and Schwer, B. 1995. RNA capping enzyme and DNA ligase: A superfamily of covalent nucleotidyl transferases. *Mol. Microbiol.* **17**: 405–410.
- Simpson, L., Sbicego, S., and Aphasizhev, R. 2003. Uridine insertion/deletion RNA editing in trypanosome mitochondria: A complex business. *RNA* **9**: 265–276.
- Stuart, K., Panigrahi, A.K., Schnauffer, A., Drozd, M., Clayton, C., and Salavati, R. 2002. Composition of the editing complex of *Trypanosoma brucei*. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **357**: 71–79.
- Vanhamme, L., Perez-Morga, D., Marchal, C., Speijer, D., Lambert, L., Geuskens, M., Alexandre, S., Ismaili, N., Göringer, U., Benne, R., et al. 1998. *Trypanosoma brucei* TBRGG1, a mitochondrial oligo(U)-binding protein that co-localizes with an in vitro RNA editing activity. *J. Biol. Chem.* **273**: 21825–21833.

RNA

A PUBLICATION OF THE RNA SOCIETY

Mitochondrial proteins and complexes in *Leishmania* and *Trypanosoma* involved in U-insertion/deletion RNA editing

LARRY SIMPSON, RUSLAN APHASIZHEV, GUANGHAN GAO, et al.

RNA 2004 10: 159-170

References

This article cites 48 articles, 32 of which can be accessed free at:
<http://rnajournal.cshlp.org/content/10/2/159.full.html#ref-list-1>

License

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).

To subscribe to *RNA* go to:
<http://rnajournal.cshlp.org/subscriptions>