Post-transcriptional regulation of microRNA expression

GREGOR OBERNOSTERER,^{1,3} PHILIPP J.F. LEUSCHNER,^{1,3} MATTIAS ALENIUS,^{2,3} and JAVIER MARTINEZ¹

¹Institute of Molecular Biotechnology of the Austrian Academy of Sciences (IMBA), A-1030 Vienna, Austria

²Institute of Molecular Pathology (IMP), A-1030 Vienna, Austria

ABSTRACT

microRNAs (miRNAs) are endogenous, noncoding ~22-nucleotide RNA molecules that have recently emerged as fundamental, post-transcriptional regulators of cognate target gene expression. Many mammalian miRNAs are expressed in a tissue-specific manner, a phenomenon that has so far been attributed to transcriptional regulation. We here show by Northern blots and in situ hybridization experiments that the expression of mammalian miRNAs can be regulated at the post-transcriptional level. In particular, miR-138 is spatially restricted to distinct cell types, while its precursor, pre-miR-138-2, is ubiquitously expressed throughout all tissues analyzed. Furthermore, pre-miR-138-2 is exported from the nucleus to the cytoplasm, suggesting that cleavage of this pre-miRNA by Dicer is restricted to certain tissues and cell types. Thus, differential processing of pre-miRNAs might be an alternative mechanism to control miRNA function.

Keywords: microRNAs; post-transcriptional regulation; Dicer; tissue-specific expression

INTRODUCTION

miRNAs are a family of small (~22 nucleotides [nt]), endogenous, noncoding RNAs that, by binding complementary sequences in the 3' untranslated region (3' UTR) of messenger RNAs (mRNAs), either mediate translational repression or direct mRNA cleavage (Pillai 2005). miRNAs are transcribed as mono- or polycistronic, long, primary precursor transcripts (pri-miRNAs) that are cleaved into ~70-nt precursor hairpins, known as pre-miRNAs, by the nuclear RNase III-like enzyme Droscha (Lee et al. 2003). Subsequently, pre-miRNA hairpins are exported to the cytoplasm by Exportin-5 (Yi et al. 2003; Bohnsack et al. 2004; Lund et al. 2004), where they are processed by a second RNase III-like enzyme, termed Dicer, into ~22-nt duplexes (Bernstein et al. 2001; Hutvagner et al. 2001; Ketting et al. 2001; Knight & Bass 2001), followed by the asymmetric assembly of one of the two strands into a functional miRNP or miRISC (Khvorova et al. 2003; Schwarz et al. 2003).

More than 800 miRNAs have so far been experimentally discovered in mammals (Lagos-Quintana et al. 2001, 2002;

Bentwich et al. 2005) and some of them are highly conserved between invertebrates and vertebrates (Bartel 2004). However, through the development of increasingly sophisticated algorithms based on machine learning techniques, the number of predicted miRNAs currently amounts to a few thousand (Lim et al. 2003; Bentwich et al. 2005; Berezikov et al. 2005). Many mammalian miRNAs are tissue- and/or developmental stage-specifically expressed. Several microarray profiling studies have shown that the expression pattern of a large number of miRNAs can be attributed to regulatory sequences present in their promoters (Babak et al. 2004; Barad et al. 2004; Calin et al. 2004; Liu et al. 2004; Miska et al. 2004; Sempere et al. 2004). Furthermore, host genes harboring miRNA sequences in their intronic sites impose their pattern of expression to the respective miRNAs (Bartel 2004; Rodriguez et al. 2004).

The current view suggests that miRNA expression is mainly controlled at the transcriptional level. For example, the transcription factors MyoD, Mef2, and SRF determine the heart-specific expression of miR-1 (Zhao et al. 2005). A recent study by O'Donnell and colleagues (O'Donnell et al. 2005) shows that the proto-oncogene c-Myc activates the expression of the *miR-17-92* cluster. However, like other RNAs, miRNA expression could potentially be controlled at the post-transcriptional level (Pillai 2005). It has been shown in *Caenorhabditis elegans* that *miR-38* is regulated in a temporal manner by differential maturation of pre-miR-38 during development (Ambros et al. 2003). We here

³These authors contributed equally to this work.

Reprint requests to: Javier Martinez, Institute of Molecular Biotechnology of the Austrian Academy of Sciences, Dr.-Bohr-Gasse 3-5, A-1030 Vienna, Austria; e-mail: javier.martinez@imba.oeaw.ac.at; fax: +43 (1) 79044-110.

Article published online ahead of print. Article and publication date are at <http://www.rnajournal.org/cgi/doi/10.1261/rna.2322506>.

provide conclusive evidence that also mammalian miRNA expression can be regulated at the level of miRNA processing.

RESULTS AND DISCUSSION

As a first step in validating targets of a set of brain-specific miRNAs, we isolated total RNA from brain and other mouse tissues, as well as from HeLa cells, and performed Northern blots with miRNA-specific probes. As expected for *miR-138*, the mature ~23-nt miRNA was detectable only in brain tissue (Fig. 1A). In particular, we found that *miR-138* was expressed in the cerebrum, the cerebellum, and the midbrain of adult mice as well as in the murine neuroblastoma cell line N2A (Fig. 1B). Strikingly, its putative ~69-nt precursor was present in all tissues and cells analyzed (Fig. 1A, left panel; 1B), suggesting that the ubiquitously expressed precursor is processed into the mature miRNA in a tissue-specific manner.

Two putative precursors were recently predicted for *miR-138*, termed “pre-miR-138-1” and “pre-miR-138-2,”

which are encoded on different chromosomal loci and are 62 nt and 69 nt in size, respectively (Lagos-Quintana et al. 2002; Griffiths-Jones 2004; Weber 2005). Multiple sequence alignments (Schwartz et al. 2000) of the two chromosomal regions showed high conservation of the mature miRNA among different vertebrate species, whereas only the locus of *miR-138-2* showed an overall conservation pattern in the flanking sequences (Supplemental Material).

To check that (1) the ~69-nt band is indeed a precursor miRNA and, if so, (2) which of the two precursors is ubiquitously expressed, we performed Northern blots with total RNA isolated from HeLa cells and probed against the mature sequence as well as sequences specific for the loop regions of the two different precursors. Interestingly, the probe against the mature *miR-138* showed a single band at 69 nt, which was also revealed by a probe against the loop region of pre-miR-138-2 (Fig. 1C). Mismatches in the probes against the sequences of the mature *miR-138* and of pre-miR-138-2 abolished detection of the 69-nt band (Supplemental Material), indicating that the ubiquitously

expressed ~69-nt band shown in Figure 1, A and B, is indeed a precursor of *miR-138*, and that the mature *miR-138* derives from pre-miR-138-2. In agreement with this, we were not able to detect a 62-nt band that would correspond to pre-miR-138-1 (Fig. 1C) either with a mature probe or with a probe against the loop region of pre-miR-138-1.

To further investigate the overall distribution of *miR-138* and its precursor, we performed in situ hybridizations with 3' DIG-labeled LNA oligonucleotide probes on cryo-sections of E17 mouse embryos (Fig. 2A) and adult brain (Fig. 2B). We observed a strong staining in the central nervous system (CNS) for mature *miR-138* (Fig. 2A, left panel). In particular, *miR-138* was primarily localized to most neurons in the hippocampus and to specific regions of the neocortex, but also to the cerebellum (Fig. 2A, left panel; 2B, upper panel). This clearly demonstrates that expression of *miR-138* is not uniform throughout the brain but restricted to distinct neuronal populations. Surprisingly, *miR-138* was also expressed in fetal liver (Fig. 2A, left panel; 2C) but not in adult liver (Fig. 2C), indicating that the expression of *miR-138* is also developmentally regulated. To confirm the ubiquitous expression observed for pre-miR-138-2 in Northern blots performed with total

FIGURE 1. (A) Northern blot analysis of different brain-specific miRNAs. Thirty micrograms of total RNA from HeLa cells and from different mouse tissues were blotted and probed with a 5'-radiolabeled, LNA-modified oligodeoxynucleotide complementary to the indicated miRNAs. Equal loading of total RNA on the gel was verified by hybridizing with a probe complementary to the U6 snRNA. Note that only the precursor of *miR-138* displays a ubiquitous pattern. The asterisk highlights the faint band of pre-miR-124a seen in the lane of murine brain at ~57 nt. (B) Northern blot analysis of total RNA isolated from HeLa cells, from the murine neuroblastoma cell line N2A, and from different neuronal tissues of murine brain showing the tissue-specific processing of the ubiquitous 69-nt pre-miRNA. The band at 69 nt corresponds to the precursor, which is present in all cells and tissues that were analyzed. The 23- to 24-nt bands represent the mature *miR-138*, which is specifically expressed in the cerebrum as well as in the cerebellum, midbrain, and N2A cells, albeit to a lower extent. (C) Northern blot analysis of pre-miR-138-2 and pre-miR-138-1. One hundred micrograms of total RNA from murine brain were blotted and probed with 5'-radiolabeled, LNA-modified oligonucleotide probes complementary to the mature sequence of *miR-138* (m), to the terminal loop of pre-miR-138-2 (p2), and to the terminal loop of pre-miR-138-1 (p1).

FIGURE 2. (A) In situ hybridization experiment on cryo-sections of E17 mouse embryos using LNA-modified probes that recognize mature miR-138 (left panel) or pre-miR-138-2 (right panel). The probe against mature miR-138 shows a strong expression in neuronal tissues (brain, CNS) and also in fetal liver, while the probe hybridizing to pre-miR-138-2 gives a strong signal in almost all tissues of the embryo. The seemingly low level of pre-miR-138-2 in brain can be explained by a low cellular density of the brain region as well as to a short exposure time used to prevent overexposure of tissues with higher cellular density. Magnifications of the fetal brain region can be seen in the Supplemental Material, where exposure time has been optimized to visualize expression of both mature miR-138 and pre-miR-138-2. (B) In situ hybridization experiments on cryo-sections of adult mouse brain using LNA-modified oligonucleotide probes that recognize mature miR-138 (upper panel) or pre-miR-138-2 (lower panel). miR-138 is primarily located to specific regions of the neocortex, most neurons in the hippocampus, and granule and purkinje cells of the cerebellum, while pre-miR-138-2 is essentially uniformly distributed. Magnifications of the cerebellum can be seen in the Supplemental Material. (C) Northern blot analysis of miR-138 expression in fetal and adult brain and liver. One hundred micrograms of total RNA prepared from the indicated tissues were blotted and probed for miR-138. In the mouse embryo, miR-138 is expressed in both brain and liver, which is in agreement with the findings in the in situ hybridization experiments. In the adult, miR-138 is restricted to brain. (D) Northern blot of total HeLa RNA and cytoplasmic and nuclear RNA fractions from HeLa cells probed against pre-miR-138-2. The band at 69 nt corresponds to pre-miR-138-2 that is mainly found in the cytoplasmic fraction, indicating nuclear export of the precursor.

RNA from adult tissues, we prepared a probe recognizing the loop region of the pre-miR-138-2 hairpin. Like Northern blots, in situ hybridizations showed a broad expression of pre-miR-138-2 in most organs of the embryo (Fig. 2A, right panel) as well as in adult brain (Fig. 2B, lower panel). To summarize, these data (Northern blots and in situ hybridizations) suggest that pre-miR-138-2 processing is both temporally and spatially regulated.

How is this differential processing achieved? A tempting hypothesis is that the export of pre-miR-138-2 is impaired in all tissues except brain, thus preventing cleavage by Dicer in the cytoplasm. To evaluate this possibility, we isolated cytoplasmic and nuclear RNA from HeLa cells, where pre-miR-138-2 is not processed, and examined its subcellular distribution. Northern blot analysis showed that the precursor is effectively exported to the cytoplasm (Fig. 2D), indicating that cleavage by Dicer could be the regulated step.

We envision that tissue-specific expression could be achieved either by an activator present in tissues that express *miR-138* or, alternatively, by an inhibitor acting on tissues that lack expression of *miR-138*. We were able to show that recombinant Dicer enzyme is able to convert pre-miR-138-2 into mature miR-138, a result that rules out the activator model (Fig. 3A). Thus, we favor the presence of an inhibitory factor, which binds pre-miR-138-2 and prevents its conversion into a mature miR-138 by Dicer in all tissues not expressing *miR-138*. This hypothesis is fostered by in vitro experiments, where the addition of increasing amounts of HeLa cytoplasmic extracts readily abolished processing of pre-miR-138-2 by recombinant Dicer (Fig. 3B). Importantly, processing of pre-miR-19a, a miRNA that is normally expressed in HeLa cells (Lagos-Quintana et al. 2001), was not abolished by titrating increasing amounts of HeLa cytoplasmic extracts (Fig.

FIGURE 4. Model depicting how differential processing of an otherwise ubiquitously expressed pre-miRNA mediates tissue- and/or developmental stage-specific expression of the mature miRNA. In both tissues, expressing or not expressing the mature miR-138, its gene is transcribed into pri-miR-138-2, which is cleaved in the nucleus by Drosha into pre-miR-138-2 and then exported to the cytoplasm. Here, the presence or the absence of an inhibitory factor determines whether pre-miR-138-2 becomes processed or not, thus leading to differential expression of mature miR-138.

Cell culture

HeLa human cervix-carcinoma cells as well as N2A murine neuroblastoma cells were grown in Dulbecco's modified Eagle's medium (DMEM) (Invitrogen) supplemented with 10% fetal calf serum (FCS) (Invitrogen), 100 u/mL penicillin (Sigma-Aldrich), 100 µg/mL penicillin/streptomycin (Sigma-Aldrich), and 20 mM HEPES (pH 7.3) at 37°C in an atmosphere containing 5% CO₂ in 15-cm dishes up to 90%–95% confluency.

HeLa cells were transfected with the indicated amounts of pSUPER.neo+GFP plasmid (Oligoengine), where the sequences of pre-miR-124a or pre-miR-138-2 were previously inserted between the BglII and the HindIII restriction sites. For pre-miR-124a, we used 124a-fwd, 5'-GATCCCCGTGTTACACAGCGGACCTTGATT TAAATGTCCATACAATTAAGGCACGCGGTGAATGCCATTTT TA-3', and 124a-rev, 5'-AGCTTAAAAATGGCATTACCCGCGT GCCTTAATTGTATGGACATTTAAATCAAGGTCCGCTGTGAA CACGGG-3'. For pre-miR-138-2, we used 138-2-fwd, 5'-GATCC CCAGCTGGTGTGTAATCAGGCCGACGAGCAGCGCATCC TCTTACCCGGCTATTTACGACACCAGGGTTTTTA-3', and 138-2-rev, 5'-AGCTTAAAAACCCTGGTGTCTGTAATAGCCG GGTAAGAGGATGCGCTGCTCGTCGGCCTGATTACACAACAC CAGCTGGG-3'. Transfections were performed in 6-well plates (Nunc) using the Lipofectamine 2000 reagent (Invitrogen) with concentrations according to the product manual. Total RNA was isolated 6 h after the start of transfection as described above.

Preparation of cellular extracts

HeLa or N2A cells were briefly washed with 1× PBS and harvested in 1 mL of a buffer containing 100 mM KCl, 5 mM MgCl₂, 10% glycerol, 30 mM HEPES (pH 7.4), 0.1 mM AEBSF, and 0.5 mM DTT by scraping with a rubber policeman. Cells were effectively

lysed by sonication. The remaining cell debris was removed by centrifugation.

Nuclear and cytoplasmic extracts were prepared as previously described (Lehnertz et al. 2003). Briefly, to prepare cytoplasmic extracts, cells were harvested by trypsinization, washed, and pelleted by centrifugation. The pellet was resuspended in 1× PBS and pelleted again by spinning. PBS was removed, and the cell pellet was resuspended in cold buffer A (10 mM HEPES at pH 7.9, 10 mM KCl, 0.1 mM EDTA, 0.1 mM EGTA, 1 mM DTT, 0.5 mM PMSF) by gentle pipetting. After swelling, a 10% solution of Nonidet NP-40 (Fluka) was added and the tube was vigorously vortexed. The homogenate was centrifuged, and the supernatant containing the cytoplasm and the cytoplasmic RNA was transferred to a fresh tube and subsequently used for RNA extraction and Northern blot analysis. Nuclei were prepared by centrifugation of the cells through a cushion of nuclear isolation buffer (20% [v/v] Ficoll-Paque Pharmacia, 80 mM Tris/HCl at pH 7.4, 8 mM MgCl₂, 8 mM CaCl₂, 1.6% [v/v] NP-40, 1.3% [v/v] Triton X-100, and 0.001% [v/v] DMSO). Nuclear pellets were washed once in ice-cold PBS, snap-frozen in liquid nitrogen, and subsequently used for the preparation of nuclear RNA. To obtain nuclear extracts, nuclei were resuspended in immunoprecipitation (IP) buffer (45 mM HEPES/NaOH at pH 7.5, 0.45 M NaCl, 0.9 mM EDTA, 0.9% [v/v] NP-40, 8.7% [v/v] glycerol, 1 mM NaF, 10 mM β-glycerophosphate, 1 tablet/50 mL protease inhibitor cocktail [Roche, no. 1836145]) and sonicated. After centrifugation, the supernatant was snap-frozen in liquid nitrogen and subsequently used for RNA extraction and Northern blot analysis.

In vitro processing assays

For in vitro processing assays, DNA templates encoding the sequences of various miRNAs were transcribed by T7 polymerase (MEGashortscript T7, Ambion) in the presence of ³²P-α-UTP (Amersham), thus generating radiolabeled, synthetic precursor miRNAs. The DNA templates were annealed to a T7 runoff DNA oligonucleotide (AATTTAATACGACTCACTATAGG) that spans the T7 promoter site. The sequences used were as follows: pre-miR-19a, 5'-TCAGTTTTGCATAGATTTGCACAACACTACATTCT TCTTGTAGTGCAACTATGCAAAACCTATAGTGAGTCGTATT AA-3'; pre-miR-138-2, 5'-AACCCCTGGTGTCTGTAATAGCC GGGTAAGAGGATGCGCTGCTCGTCGGCCTGATTACACAACA CCAGCCTATAGTGAGTCGTATTAA-3'. These synthetic precursors were folded into their hairpin-shaped structure by heating for 1 min at 95°C and cooling slowly to room temperature. Processing with recombinant Dicer was performed as previously described (Zhang et al. 2002). Precursors were used at a concentration of 10 nM and were pretreated for 10 min at 30°C with increasing amounts of HeLa cytoplasmic extract (2, 4, 6, 8, 16, 24, and 32 µg protein for pre-miR-138-2; 2, 6, 16, and 32 µg protein for pre-miR-19a). After 2 min incubation at 37°C, the reaction products were separated on a 15% denaturing PAGE and visualized by autoradiography.

In situ hybridizations

All mice were maintained at the animal facility at the IMP, under pathogen-free conditions. C57BL/6 mice were mated to generate embryos for analyses and the morning of the vaginal plug was considered as E0.5. Embryos, livers, and brains were post-fixed

in 4% paraformaldehyde, cryoprotected (30% sucrose in PBS), embedded in Tissue-Tek OCT compound, and cryosectioned. Ten-micron cryosections were pretreated, hybridized with LNA digoxigenin-labeled probes (Exiqon), and washed according to Schaeren-Wiemers and Gerfin-Moser (1993), with some modifications. Probe sequences were as follows: miR-122a, 5'-ACAAAC ACCATTGTCACACTCCA-3'; miR-138, 5'-CGGCCTGATTCA CAACACCAGCT-3'; and pre-miR-138-2, 5'-GGTAAGAGGATG CGCTGCTCGT-3'. Briefly, sections were fixed in 4% paraformaldehyde for 10 min, acetylated, and treated with 5 µg/mL proteinase K (Roche) in PBS for 5 min, washed, and prehybridized for 4 h at room temperature. Hybridization with 22-nt LNA probes ($T_m \sim 80^\circ\text{C}$) was performed at 55°C overnight. Slides were then washed at 60°C and incubated with alkaline phosphatase-conjugated goat anti-DIG Fab fragments (1:2000 [Roche]) at 4°C overnight. Fluorescent detection was performed using a Fast Red reaction (Dako Cytomation) for 1 h at room temperature. Sections were analyzed with a Zeiss Axioplan-2 microscope and photographed with a digital camera (Coolsnap HQ, Photometrics). A subset of images was adjusted for levels, brightness, contrast, hue, and saturation with Adobe Acrobat 7.0 imaging software to optimally visualize the expression patterns.

SUPPLEMENTAL MATERIAL

Supplemental Material can be found at <http://www.imba.oew.ac.at/index.php?id=142>.

ACKNOWLEDGMENTS

We thank Stefan Ameres, Stefan Weitzer, and the members of the laboratory for encouragement and suggestions during the completion of this work, and Prof. Renée Schroeder for critically reading the manuscript. J.M. is a Junior Group Leader at IMBA. P.J.F.L. is funded by the Boehringer Ingelheim Fonds Ph.D. Scholarship. M.A. was supported by an EMBO long-term fellowship and a Marie Curie fellowship.

Received December 12, 2005; accepted April 24, 2006.

REFERENCES

- Ambros, V., Lee, R.C., Lavanway, A., Williams, P.T., and Jewell, D. 2003. MicroRNAs and other tiny endogenous RNAs in *C. elegans*. *Curr. Biol.* **13**: 807–818.
- Babak, T., Zhang, W., Morris, Q., Blencowe, B.J., and Hughes, T.R. 2004. Probing microRNAs with microarrays: Tissue specificity and functional inference. *RNA* **10**: 1813–1819.
- Barad, O., Meiri, E., Avniel, A., Aharonov, R., Barzilai, A., Bentwich, I., Einav, U., Gilad, S., Hurban, P., Karov, Y., et al. 2004. MicroRNA expression detected by oligonucleotide microarrays: System establishment and expression profiling in human tissues. *Genome Res.* **14**: 2486–2494.
- Bartel, D.P. 2004. MicroRNAs: Genomics, biogenesis, mechanism, and function. *Cell* **116**: 281–297.
- Bentwich, I., Avniel, A., Karov, Y., Aharonov, R., Gilad, S., Barad, O., Barzilai, A., Einat, P., Einav, U., Meiri, E., et al. 2005. Identification of hundreds of conserved and nonconserved human microRNAs. *Nat. Genet.* **37**: 766–770.
- Berezikov, E., Guryev, V., van de Belt, J., Wienholds, E., Plasterk, R.H., and Cuppen, E. 2005. Phylogenetic shadowing and computational identification of human microRNA genes. *Cell* **120**: 21–24.
- Bernstein, E., Caudy, A.A., Hammond, S.M., and Hannon, G.J. 2001. Role for a bidentate ribonuclease in the initiation step of RNA interference. *Nature* **409**: 363–366.
- Bohnsack, M.T., Czapinski, K., and Gorlich, D. 2004. Exportin 5 is a RanGTP-dependent dsRNA-binding protein that mediates nuclear export of pre-miRNAs. *RNA* **10**: 185–191.
- Calin, G.A., Liu, C.G., Sevignani, C., Ferracin, M., Felli, N., Dumitru, C.D., Shimizu, M., Cimmino, A., Zupo, S., Dono, M., et al. 2004. MicroRNA profiling reveals distinct signatures in B cell chronic lymphocytic leukemias. *Proc. Natl. Acad. Sci.* **101**: 11755–11760.
- Griffiths-Jones, S. 2004. The microRNA registry. *Nucleic Acids Res.* **32**: D109–D111.
- Hutvagner, G., McLachlan, J., Bálint, É., Tuschl, T., and Zamore, P.D. 2001. A cellular function for the RNA interference enzyme Dicer in small temporal RNA maturation. *Science* **93**: 834–838.
- Ketting, R.F., Fischer, S.E., Bernstein, E., Sijen, T., Hannon, G.J., and Plasterk, R.H. 2001. Dicer functions in RNA interference and in synthesis of small RNA involved in developmental timing in *C. elegans*. *Genes & Dev.* **15**: 2654–2659.
- Khvorova, A., Reynolds, A., and Jayasena, S.D. 2003. Functional siRNAs and miRNAs exhibit strand bias. *Cell* **115**: 209–216.
- Knight, S.W. and Bass, B.L. 2001. A role for the RNase III enzyme DCR-1 in RNA interference and germ line development in *C. elegans*. *Science* **2**: 2.
- Lagos-Quintana, M., Rauhut, R., Lendeckel, W., and Tuschl, T. 2001. Identification of novel genes coding for small expressed RNAs. *Science* **294**: 853–858.
- Lagos-Quintana, M., Rauhut, R., Yalcin, A., Meyer, J., Lendeckel, W., and Tuschl, T. 2002. Identification of tissue-specific microRNAs from mouse. *Curr. Biol.* **12**: 735–739.
- Lee, Y., Ahn, C., Han, J., Choi, H., Kim, J., Yim, J., Lee, J., Provost, P., Radmark, O., Kim, S., et al. 2003. The nuclear RNase III Drosha initiates microRNA processing. *Nature* **425**: 415–419.
- Lehnertz, B., Ueda, Y., Derijck, A.A., Braunschweig, U., Perez-Burgos, L., Kubicek, S., Chen, T., Li, E., Jenuwein, T., and Peters, A.H. 2003. Suv39h-mediated histone H3 lysine 9 methylation directs DNA methylation to major satellite repeats at pericentric heterochromatin. *Curr. Biol.* **13**: 1192–1200.
- Lim, L.P., Glasner, M.E., Yekta, S., Burge, C.B., and Bartel, D.P. 2003. Vertebrate microRNA genes. *Science* **299**: 1540.
- Liu, C.G., Calin, G.A., Meloon, B., Gamlie, N., Sevignani, C., Ferracin, M., Dumitru, C.D., Shimizu, M., Zupo, S., Dono, M., et al. 2004. An oligonucleotide microchip for genome-wide microRNA profiling in human and mouse tissues. *Proc. Natl. Acad. Sci.* **101**: 9740–9744.
- Lund, E., Guttinger, S., Calado, A., Dahlberg, J.E., and Kutay, U. 2004. Nuclear export of microRNA precursors. *Science* **303**: 95–98.
- Miska, E.A., Alvarez-Saavedra, E., Townsend, M., Yoshii, A., Sestan, N., Rakic, P., Constantine-Paton, M., and Horvitz, H.R. 2004. Microarray analysis of microRNA expression in the developing mammalian brain. *Genome Biol.* **5**: R68.
- O'Donnell, K.A., Wentzel, E.A., Zeller, K.I., Dang, C.V., and Mendell, J.T. 2005. c-Myc-regulated microRNAs modulate E2F1 expression. *Nature* **435**: 839–843.
- Pillai, R.S. 2005. MicroRNA function: Multiple mechanisms for a tiny RNA? *RNA* **11**: 1753–1761.
- Rodriguez, A., Griffiths-Jones, S., Ashurst, J.L., and Bradley, A. 2004. Identification of mammalian microRNA host genes and transcription units. *Genome Res.* **14**: 1902–1910.
- Schaeren-Wiemers, N. and Gerfin-Moser, A. 1993. A single protocol to detect transcripts of various types and expression levels in neural tissue and cultured cells: In situ hybridization using digoxigenin-labelled cRNA probes. *Histochemistry* **100**: 431–440.
- Schwartz, S., Zhang, Z., Frazer, K.A., Smit, A., Riemer, C., Bouck, J., Gibbs, R., Hardison, R., and Miller, W. 2000. PipMaker-A web server for aligning two genomic DNA sequences. *Genome Res.* **10**: 577–586.

- Schwarz, D.S., Hutvagner, G., Du, T., Xu, Z., Aronin, N., and Zamore, P.D. 2003. Asymmetry in the assembly of the RNAi enzyme complex. *Cell* **115**: 199–208.
- Sempere, L.F., Freemantle, S., Pitha-Rowe, I., Moss, E., Dmitrovsky, E., and Ambros, V. 2004. Expression profiling of mammalian microRNAs uncovers a subset of brain-expressed microRNAs with possible roles in murine and human neuronal differentiation. *Genome Biol.* **5**: R13.
- Valoczi, A., Hornyik, C., Varga, N., Burgyan, J., Kauppinen, S., and Havelda, Z. 2004. Sensitive and specific detection of microRNAs by northern blot analysis using LNA-modified oligonucleotide probes. *Nucleic Acids Res.* **32**: e175.
- Weber, M.J. 2005. New human and mouse microRNA genes found by homology search. *FEBS J.* **272**: 59–73.
- Yi, R., Qin, Y., Macara, I.G., and Cullen, B.R. 2003. Exportin-5 mediates the nuclear export of pre-microRNAs and short hairpin RNAs. *Genes & Dev.* **17**: 3011–3016.
- Zhang, H., Kolb, F.A., Brondani, V., Billy, E., and Filipowicz, W. 2002. Human Dicer preferentially cleaves dsRNAs at their termini without a requirement for ATP. *EMBO J.* **21**: 5875–5885.
- Zhao, Y., Samal, E., and Srivastava, D. 2005. Serum response factor regulates a muscle-specific microRNA that targets Hand2 during cardiogenesis. *Nature* **436**: 214–220.

RNA

A PUBLICATION OF THE RNA SOCIETY

Post-transcriptional regulation of microRNA expression

Gregor Obernosterer, Philipp J.F. Leuschner, Mattias Alenius, et al.

RNA 2006 12: 1161-1167

References This article cites 34 articles, 14 of which can be accessed free at:
<http://rnajournal.cshlp.org/content/12/7/1161.full.html#ref-list-1>

Open Access Freely available online through the *RNA* Open Access option.

License Freely available online through the open access option.

Email Alerting Service Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).

To subscribe to *RNA* go to:
<http://rnajournal.cshlp.org/subscriptions>