REPORT

Down-regulation of endogenes mediated by a transitive silencing signal

ANNICK BLEYS, HELENA VAN HOUDT,¹ and ANNA DEPICKER

Department of Plant Systems Biology, Flanders Interuniversity Institute for Biotechnology (VIB), Ghent University, B-9052 Gent, Belgium

ABSTRACT

Some RNA silencing systems in plants, nematodes, and fungi show spreading of silencing along target sequences, termed transitive silencing. Here, we address the question of whether endogenous targets can be silenced by a transitive silencing signal in plants. In transgenic *Arabidopsis thaliana* plants that harbored a silencing-inducing locus and a transgenic chimeric primary target, silencing of a secondary transgenic target occurred and the expression of the endogenous catalase genes was down-regulated, coinciding with a knock-down phenotype. Strikingly, the efficiency of the catalase silencing appeared to be correlated with the zygoty of the primary target locus and, to a lesser extent, with that of the silencing-inducing locus. These data suggest that silencing of an endogene induced by transgenic secondary small interfering RNAs (siRNAs) might depend on the amount of primary target transcripts that can act as template for the production of an efficient transitive silencing signal.

Keywords: RNA silencing; transitivity; inverted repeat locus; gene family; *Arabidopsis thaliana*

INTRODUCTION

RNA silencing is a general term used for epigenetic regulatory pathways in which small RNAs of 21–28 nucleotides (nt) mediate either transcriptional gene silencing by changes in chromatin state or post-transcriptional gene silencing (PTGS) through target degradation or translational repression (for review, see Baulcombe 2004). Since the accidental discovery that the introduction of a sense overexpressing transgene can lead to cosuppression of both the transgene and the endogene (Napoli et al. 1990), RNA silencing has been widely used as a tool to down-regulate gene expression. Transgenic RNAs are recruited in the silencing pathway by the production of double-stranded RNA (dsRNA), which is processed by an RNase III-like enzyme (Dicer) into “small interfering” RNAs (siRNAs) that mediate the degradation of target RNAs and possibly also DNA methylation by RNA-induced silencing complex (RISC)-like effector complexes (Mette et al. 1999; Hamilton et al. 2002; Zilberman et al. 2003).

In plants, nematodes, and fungi, not only can trigger-derived siRNAs be detected, but also siRNAs that corre-

spond to regions upstream from the region targeted by the dsRNA trigger. Moreover, in plants and fungi, siRNAs can be found corresponding to the downstream region as well (Sijen et al. 2001; Braunstein et al. 2002; Han and Grierson 2002; Klahre et al. 2002; Sanders et al. 2002; Vaistij et al. 2002; Nicolás et al. 2003; Van Houdt et al. 2003). Secondary siRNAs have been shown to induce silencing of targets without homology with the silencing inducer (for review, see Bleys et al. 2006). This phenomenon, designated transitive silencing, involves an amplification process catalyzed by the RNA-dependent RNA polymerase RDR6 that synthesizes dsRNA by using target RNAs as template, either by priming of primary siRNAs (Lipardi et al. 2001; Sijen et al. 2001) or by a primer-independent mechanism that starts at the 3' end of the target RNAs (Schiebel et al. 1993; Vaistij et al. 2002; Tang et al. 2003; Petersen and Albrechtsen 2005).

In *Caenorhabditis elegans* and *Mucor circinelloides*, transitivity occurs along transcripts derived from both transgenes and endogenes (Sijen et al. 2001; Alder et al. 2003; Nicolás et al. 2003). In plants, transgenes and endogenes seem to behave differently in relation to transitivity. Many studies have demonstrated spreading of silencing along transgene sequences, but failed to do so for endogenous transcripts (Vaistij et al. 2002; Himber et al. 2003; Kościńska et al. 2005; Miki et al. 2005; Petersen and Albrechtsen 2005). The ability of endogenes to serve as template for the RDR6-mediated production of secondary siRNAs has been described just once (Sanders et al. 2002). Because all these transitivity studies only investigated the ability of secondary siRNAs to

¹Present address: Sint-Lieven, Gildestraat 17, B-9000 Gent, Belgium.

Reprint requests to: Anna Depicker, Department of Plant Systems Biology, Flanders Interuniversity Institute for Biotechnology (VIB), Ghent University, Technologiepark 927, B-9052 Gent, Belgium; e-mail: ann.depicker@psb.ugent.be; fax: 32-9-3313809.

Article and publication are at <http://www.rnajournal.org/cgi/doi/10.1261/rna.108106>.

induce silencing of transgenic secondary targets, it is not known whether they can target plant endogenes as well.

Previously, we have established that in tobacco (*Nicotiana tabacum*) a post-transcriptionally silenced inverted repeat (IR) transgene locus (X) is able to silence a nonhomologous transgene (Z), when a stepwise homology is created by introducing a chimeric primary target (Y) with one region homologous to the silencing inducer X and a second upstream region homologous to the secondary target Z (Van Houdt et al. 2003). A similar XYZ system has been developed in *Arabidopsis thaliana* (L.) Heynh., allowing us to demonstrate that a transitive silencing signal produced by an amplification mechanism is able to trigger silencing of endogenous targets, and that the efficiency of silencing correlates with the zygoty of the primary target locus, suggesting a connection with the transcript abundance.

RESULTS AND DISCUSSION

Development of a transitive silencing system in *Arabidopsis*

First, we checked whether a particular isolated IR transgene locus X_{21} (P35S-*nptII*-3'chs; Fig. 1a) in *Arabidopsis* is able to transmit its in *trans* silencing capacity to a primary target

FIGURE 1. Schematic outline of the T-DNA constructs (drawn to scale), present in locus X_{21} , primary targets Y_7 and Y_{11} , and secondary target Z_c . (a) Loci X_{21} , Y_7 , and Z_c used to assess transitive silencing of transgenes in *A. thaliana*; (b) locus Y_{11} used to assess transitive silencing of endogenes. Silenced locus X_{21} contains two copies of the DEchs278 T-DNA that are arranged as an IR around the right border. Locus Z_c is a single-copy locus. The copy number of the Y_7 and Y_{11} loci is unknown. 3'chs, 3'-UTR of the chalcone synthase gene of *Anthriscum majus*; 3'g7, 3'-UTR of the *Agrobacterium tumefaciens* octopine T-DNA gene 7; 3'nos, 3'-UTR of the nopaline synthase gene; bar, bialaphos acetyltransferase-coding sequence conferring phosphinothricin resistance; CAT2 800, last 800 nt of the *A. thaliana* CAT2 sequence; gus c.s., gus-coding sequence; hpt, hygromycin phosphotransferase-coding sequence; I, artificial intron; nptII c.s., neomycin phosphotransferase II-coding sequence; P35S, CaMV 35S promoter; Pnos, nopaline synthase promoter; Pss, promoter of the small subunit of ribulose-1,5-bisphosphate carboxylase; LB, left T-DNA border; RB, right T-DNA border.

Y_7 (P35S-*gus-npt3'chs*), resulting in transitive silencing of a secondary target transgene Z_c (P35S-*gus-npt3'nos*), similar to the transitive XYZ system in tobacco (Van Houdt et al. 2003). For details of constructs, plant material, and crosses, we refer to Materials and Methods. Genotype and zygoty of the transgenic loci in the different *Arabidopsis* plants were schematically annotated as follows: slashes separate the loci X, Y, and Z; XX, YY, or ZZ represent homozygosity for the respective loci; hyphens indicate hemizygoty, double hyphens azygoty, and an asterisk unknown zygoty with the locus either homo- or hemizygous.

The occurrence of transitive silencing in X_{21} -/ Y_7 -/ Z_c hybrid plants was analyzed through measurement of the GUS activity, because both Y_7 and Z_c encode a functional β -glucuronidase (GUS) protein. A 130- to 140-fold reduction in GUS activity was observed in 11 X_{21} -/ Y_7 -/ Z_c hybrid plants (Fig. 2a, A) compared to the progeny of the parental plants (17 --/ Y_7^* -/ Z_c and 10 $X_{21}X_{21}$ -/ Z_c plants; Fig. 2a, B and C, respectively). These data indicate that not only in tobacco (Van Houdt et al. 2003), but also in *Arabidopsis*, in *trans*-silenced Y transcripts can produce a transitive silencing signal leading to spreading of silencing.

In *C. elegans*, transitive RNA interference (RNAi) is not limited to transgenic targets (Sijen et al. 2001). In plants, however, secondary siRNAs have been shown to target transgenic and viral sequences (Sanders et al. 2002; Vaistij et al. 2002; Van Houdt et al. 2003), but it has never been investigated whether endogenous sequences can be silenced by a transitive silencing signal. Therefore, we constructed Y_{11} (P35S-CAT2 800-*npt3'chs*; Fig. 1b), which harbors an expression cassette with the last 800 nt of the CAT2-coding sequence. The objective was to determine whether the CAT genes in *A. thaliana* can be transitively silenced by combining the X_{21} and Y_{11} loci. The *Arabidopsis* catalase (CAT) multigene family consists of three members (CAT1, CAT2, and CAT3), with a nucleotide and amino acid sequence identity of 70%–72% and 75%–84%, respectively (Frugoli et al. 1996).

All wild-type (Fig. 2b, D) and parental plants ($X_{21}X_{21}$ -/ $CATCAT$ and --/ $Y_{11}Y_{11}$ -/ $CATCAT$; Fig. 2b, B and C, respectively) retained a normal CAT activity, whereas the X_{21} -/ Y_{11} -/ $CATCAT$ F1 hybrid plants (Fig. 2b, A; see Materials and Methods) had an average total CAT activity of 36%. This level of silencing corresponds with that observed in the hairpin RNA (hpRNA)-silenced line CAT2HP1 (Fig. 2b, E; 33% total CAT activity; Vandenaabeele et al. 2004), but is less strong than the reduction seen in the CAT2HP2 line (Fig. 2b, F; 3% total CAT activity; Vandenaabeele et al. 2004).

To analyze the silencing at the RNA level, we performed a quantitative real-time PCR with a CAT2-specific primer pair (see Materials and Methods). Total RNA was prepared from leaf material, and first-strand cDNA was generated

FIGURE 2. Protein and RNA levels in F1 hybrid plants. (a) GUS activity levels in protein extracts of leaf tissue harvested from transgenic *Arabidopsis* plants containing different combinations of loci X_{21} , Y_7 and Z_c . SD is given by error bars. A, $X_{21}/Y_7/Z_c$; B, $-/Y_7^*/Z_c$; C, $X_{21}X_{21}/-/Z_c$; n, total number of plants tested; TSP, total soluble protein. (b) Average total CAT activity in wild-type and transgenic *Arabidopsis* plants containing different combinations of loci X_{21} and Y_{11} or containing a CAT2 hairpin construct (Vandenabeele et al. 2004). CAT activity in protein extracts of leaf tissue was measured with a spectrophotometric assay. The average total CAT activity of the different transgenic plants is relative to the average CAT activity of four wild-type (Col-0, D) plants, which is set at 100%. SD is given by error bars. A, $X_{21}/Y_{11}/CATCAT$; B, $X_{21}X_{21}/-/CATCAT$; C, $-/Y_{11}Y_{11}/CATCAT$; D, Col-0; E, CAT2HP1; F, CAT2HP2; n, total number of plants tested. (c) CAT2 mRNA abundance. A quantitative real-time PCR was performed using CAT2-specific primers to determine the CAT2 RNA level relative to that in a wild-type background (100%). A1, A2, and A3, RNA levels in three $X_{21}/Y_{11}/CATCAT$ plants; B, RNA level in one $X_{21}X_{21}/-/CATCAT$ plant; C, RNA level in one $-/Y_{11}Y_{11}/CATCAT$ plant; D, RNA levels in one wild-type Col-0; E, RNA level in one CAT2HP1 plant; F, RNA level in one CAT2HP2 plant.

with an oligo(dT)₁₂₋₁₈ primer. Figure 2c clearly shows a two- to fivefold lower CAT2 transcript level in $X_{21}/Y_{11}/CATCAT$ plants (A1, A2, and A3) than that of the wild-type plant (D). The reduction was similar in the CAT2HP1 plant (Fig. 2c, E), but 20-fold lower in the CAT2HP2 plant (Fig. 2c, F). The parental plants ($X_{21}X_{21}/-/CATCAT$ and $-/Y_{11}Y_{11}/CATCAT$; Fig. 2c, B and C, respectively) showed a slight increase in the CAT2 RNA level. We can conclude that an in *trans*-silenced transgenic primary target produces transitive silencing signals that are able to down-regulate the expression of endogenes. This capacity does not depend on the potential of the primary target transcripts to be translated into a functional protein, because the Y_{11} construct contains no translation start codon.

Influence of the zygosity of the silencing inducer and the primary target on the efficiency of transitive silencing of an endogene

To investigate the stability of endogene suppression via a transitive signal and to determine whether the efficiency of silencing is affected by the homozygous condition of either the silencing-inducing locus X_{21} or the primary target locus Y_{11} , seeds from a self-fertilized F1 hybrid plant ($X_{21}/Y_{11}/CATCAT$) were grown on medium selective for both X_{21} and Y_{11} . The two loci segregated independently, and these F2 progeny plants were expected to have genotypes $X_{21}X_{21}/Y_{11}Y_{11}/CATCAT$, $X_{21}/Y_{11}Y_{11}/CATCAT$, $X_{21}X_{21}/Y_{11}/CATCAT$, or $X_{21}/Y_{11}/CATCAT$ at a 1:2:2:4 ratio. Thirty-six selected F2 progeny plants were analyzed and ranked according to their total CAT activity (Fig. 3a). Segregation analysis of the 36 plants showed a correlation between the zygosity of loci X_{21} and Y_{11} and the efficiency of transitive CAT silencing. The eight plants exhibiting very efficient silencing (<3% total CAT activity; Fig. 3a, A) were all plants homozygous for locus Y_{11} ($X_{21}X_{21}/Y_{11}Y_{11}/CATCAT$ and $X_{21}/Y_{11}Y_{11}/CATCAT$), whereas the seven plants with the lowest degree of reduction (between 42% and 59% total CAT activity; Fig. 3a, C) were hemizygous for locus Y_{11} , but homozygous for locus X_{21} ($X_{21}X_{21}/Y_{11}/CATCAT$). The 21 plants with intermediate silencing efficiencies (between 13% and 34% total CAT activity; Fig. 3a, B) were hemizygous for both loci ($X_{21}/Y_{11}/CATCAT$). These results suggest that the zygosity of the primary target locus Y_{11} and to a lesser extent that of the silencing-inducing locus X_{21} influence the efficiency of transitive silencing of the CAT genes. We cannot exclude the possibility that some of the observed CAT silencing results from the activation of cosuppression induced by the sense construct Y_{11} , because sense-induced PTGS can occur stochastically in transformed plants. However, the Y_{11} locus did not induce silencing in any of the 24 $-/Y_{11}Y_{11}/CATCAT$ plants (Fig. 2b, C; Fig. 3a, D), making it is very likely that the efficient silencing observed in $X_{21}X_{21}/Y_{11}Y_{11}/CATCAT$ plants results from transitive signals produced by Y_{11} .

The positive correlation between the silencing efficiency and the zygosity of the primary target locus suggests that the amount of primary transcripts serving as template for RDR6 might determine the level of secondary siRNAs produced, thus resulting in a certain degree of silencing of the secondary target. Analyses in the transitive XYZ system in tobacco have demonstrated that the accumulation level of secondary siRNAs depends on the zygosity of the primary target locus from which they are derived (García-Pérez et al. 2004). The negative correlation in plants hemizygous for Y_{11} between the level of silencing and the zygosity of locus X_{21} indicates that the primary silencing signal also plays a role. Perhaps a higher dose of primary siRNAs might be produced by locus X_{21} in the homozygous condition, which could trigger a higher rate of

FIGURE 3. Protein levels, siRNA accumulation, and RNA levels in F2 progeny plants and parental plants. (a) Total CAT activity in F2 progeny plants with segregating X_{21} and Y_{11} loci and parental $-/-Y_{11}Y_{11}/CATCAT$ plants. The total CAT activity of the different transgenic plants is relative to the average CAT activity of five wild-type plants (100%; not shown). A, $X_{21}X_{21}/Y_{11}Y_{11}/CATCAT$; B, $X_{21}X_{21}/Y_{11}Y_{11}/CATCAT$; C, $X_{21}X_{21}/Y_{11}Y_{11}/CATCAT$; D, $-/-Y_{11}Y_{11}/CATCAT$; n, total number of plants per group. (b) Accumulation of $CAT2$ -specific siRNAs in two silenced $X_{21}X_{21}/Y_{11}Y_{11}/CATCAT$ plants (A₁ and A₂) and absence of siRNAs in one nonsilenced $-/-Y_{11}Y_{11}/CATCAT$ plant (B). Low molecular weight RNA was extracted from leaves of 4-wk-old plants. RNA oligomers of 21 nt and 24 nt were used as molecular markers. siRNAs were detected with a hydrolyzed ^{32}P -labeled probe comprising 274 bp in the 3' end of the $CAT2$ sequence. With this probe it is impossible to discriminate between siRNAs originating from the transgene Y_{11} or from the endogenous sequences. The predominant ethidium bromide (EtBr)-stained species in the low molecular weight RNA fraction are shown as loading controls. (c) $CAT1$, $CAT2$, and $CAT3$ mRNA abundancies. Quantitative real-time PCRs were performed with gene-specific primers to determine the RNA levels of $CAT1$ (shaded bars), $CAT2$ (black bars), and $CAT3$ (dotted bars) relative to those in a wild-type background (100%). A, RNA levels in one $X_{21}X_{21}/Y_{11}Y_{11}/CATCAT$ plant; B1, B2, and B3, RNA levels in three $X_{21}X_{21}/Y_{11}Y_{11}/CATCAT$ plants; C, RNA levels in one $X_{21}X_{21}/Y_{11}Y_{11}/CATCAT$ plant; D, RNA levels in wild-type Col-0.

RISC-mediated cleavage of Y_{11} templates. This hypothesis is supported by the fact that a hairpin construct controlled by the strong 35S promoter induces a stronger silencing phenotype than the same construct controlled by the weak nopaline synthase promoter (Chuang and Meyerowitz 2000). In this case, the production of more hpRNAs and thus siRNAs leads to a more efficient degradation and silencing of the endogenous target RNAs.

We also looked at the accumulation of siRNAs that are the key molecules of PTGS. An RNA gel blot with a hydrolyzed ^{32}P -labeled probe comprising 274 bp in the 3' end of the $CAT2$ sequence revealed the presence of $CAT2$ -specific siRNAs in $X_{21}X_{21}/Y_{11}Y_{11}/CATCAT$ (Fig. 3b, A₁ and A₂) plants, whereas in $-/-Y_{11}Y_{11}/CATCAT$ plants (Fig. 3b, B) no such molecules were detected.

To assess whether the most efficiently silenced $X_{21}X_{21}/Y_{11}Y_{11}/CATCAT$ progeny plants displayed the characteristic CAT-deficient necrosis phenotype, we performed a high-light (HL) treatment (see Materials and Methods). HL irradiation induces photorespiration that leads to accumulation of H_2O_2 in CAT-deficient plants, followed by active cell death (Chamngongpol et al. 1998; Dat et al. 2003; Vandenabeele et al. 2004). All $X_{21}X_{21}/Y_{11}Y_{11}/CATCAT$ progeny plants had visible signs of cell death within 8 h of HL irradiation, whereas wild-type control plants did not, even after 48 h of HL (Fig. 4). In conclusion, transitive silencing elicits the same knock-down phenotype as hairpin-induced silencing.

Differential susceptibility of different members of the CAT gene family to transitive silencing signals

The *Arabidopsis* $CAT2$ gene targeted by the Y_{11} construct is part of a gene family. All three members are highly expressed in inflorescences, but in leaves only the $CAT2$ and $CAT3$ genes are highly expressed (Frugoli et al. 1996). With the CAT activity assay used above, it is not possible to discriminate between the three CAT proteins, because the total CAT activity is measured, implying that possible differential silencing efficiencies of $CAT1$, $CAT2$, and $CAT3$ cannot be determined in this way. To assay to what extent the three CAT genes were silenced in F2 progeny plants, we performed a quantitative real-time PCR with gene-specific primer pairs. In wild-type samples, the $CAT1$

FIGURE 4. Phenotypes of wild-type and $X_{21}X_{21}/Y_{11}Y_{11}/CATCAT$ plants after exposure to high-light (HL) irradiation ($1000 \mu\text{mol m}^{-2} \text{sec}^{-1}$). Cell death was visible in CAT-deficient $X_{21}X_{21}/Y_{11}Y_{11}/CATCAT$ plants within 8 h of HL irradiation, while in wild-type control plants this phenotype did not occur, even not after 48 h of HL.

and *CAT3* mRNAs were ~500- to 600-fold and 20- to 25-fold less abundant than those of *CAT2*, respectively (data not shown).

The transcript levels of *CAT1*, *CAT2*, and *CAT3* in leaf tissue of F2 progeny plants relative to those of a wild-type plant are given in Figure 3c. In this graph, the mRNA abundancies of *CAT1*, *CAT2*, and *CAT3* in the wild-type plant (Fig. 3c, D) are all three set at 100%. The *CAT2* RNA level was sevenfold reduced in a $X_{21}X_{21}/Y_{11}Y_{11}/CATCAT$ sample (Fig. 3c, A), 2.3-fold in three $X_{21}/Y_{11}/CATCAT$ samples (Fig. 3c, B1, B2, B3), and not significantly in a $X_{21}X_{21}/Y_{11}/CATCAT$ sample (Fig. 3c, C), indicating that the reduction in *CAT2* RNA levels also correlated with the zygosity of loci X_{21} and Y_{11} . The transcript levels of *CAT3* seemed to be reduced to the same extent as those of *CAT2*. This observation is very intriguing, because a sevenfold reduction in *CAT2* RNAs in the $X_{21}X_{21}/Y_{11}Y_{11}/CATCAT$ sample results in a residual *CAT2* RNA level that is higher than the nonsilenced *CAT3* steady-state RNA level. Still, the *CAT3* RNAs are targeted for degradation to the same degree as those of *CAT2*, depending on the zygosity of both loci X_{21} and Y_{11} . We cannot explain this result, but it shows that there are factors other than just the RNA abundance that determine the susceptibility of particular transcripts for siRNA-mediated turnover, for instance, the spatial and temporal gene expression patterns. Similarly, Kerschen et al. (2004) found that 25 different endogenous target sequences possess an inherent degree of susceptibility to RNA silencing induced by hairpin constructs, which was independent from the gene expression levels.

The degree of reduction in *CAT2* and *CAT3* RNAs is consistent with the dosage dependency observed at the protein level. However, the reduction at the RNA level is not as pronounced as that at the protein level; for instance, in $X_{21}X_{21}/Y_{11}Y_{11}/CATCAT$ plants the *CAT2* and *CAT3* RNA levels are reduced to sevenfold, whereas the total CAT activity is reduced to 44-fold. This supports previous data (Van Houdt et al. 1997) and implies that the silencing signal can also act at the translational level, either because the detected mRNAs might not be translatable anymore (e.g., no 5' cap structure) or might be sequestered in cytoplasmic processing bodies (P-bodies) that are found in yeast and mammalian cells (Rossi 2005).

In contrast to *CAT2* and *CAT3*, the *CAT1* mRNA abundance is not significantly reduced in plants that are hemizygous for locus Y_{11} (Fig. 3c, B1,B2,B3), but it is in Y_{11} homozygous plants (Fig. 3c, A). This difference in susceptibility to silencing between *CAT1* and *CAT3* is most probably not the consequence of a different degree in homology, because both the sequence identity of the *CAT1* and *CAT3* genes with the Y_{11} transcript (78% and 75%, respectively) as well as the number of stretches of at least 21 nt with full complementarity are very similar. One of the contributing factors might be the different accumulation levels of the *CAT1* and *CAT3* genes, the former being about 25-fold less abundant than the latter.

Conclusions

We demonstrate that secondary siRNAs originating from a transgenic primary target are able to down-regulate the expression of the endogenous *CAT* gene family in *A. thaliana*, resulting in a knock-down phenotype. Our experimental setup suggests that the expression of genes in gene families is determined by a subtle equilibrium between the gene expression level and the amount of siRNAs and could reflect an epigenetic gene regulation mechanism mediated by endogenous small RNAs.

MATERIALS AND METHODS

Constructs

The T-DNA-derived plant transformation vector pDEchs287 (referred to as P35S-*nptII*-3'-*chs*) carried a neomycin phosphotransferase II (*nptII*) gene (conferring kanamycin resistance) fused to the 3'-untranslated region (UTR) of the *Antirrhinum majus* chalcone synthase gene (3'-*chs*, 287 nt) and transcribed from the cauliflower mosaic virus 35S promoter (P35S) toward the right T-DNA border. After floral dip transformation with the DEchs287 T-DNA, plants were obtained harboring the IR locus X_{21} (see below; Fig. 1a). The T-DNA vector pPs35SGUSnpt3'-*chs* (referred to as P35S-*gus*-*npt3'*-*chs*) contained two chimeric genes between the T-DNA borders: the bialaphos acetyltransferase-coding (*bar*) sequence conferring phosphinothricin resistance, fused to the 3'-UTR of the *Agrobacterium tumefaciens* octopine T-DNA gene 7 (3'-*g7*) and under the control of the promoter of the small subunit of ribulose-1,5-bisphosphate carboxylase (*Pss*) and a P35S-driven *gus* transgene fused to the last 122 nt of the *nptII* gene and the *A. majus* *chs* terminator sequence (*npt3'-chs*). After floral dip transformation with the Ps35SGUSnpt3'-*chs* T-DNA, plants were obtained harboring the single locus Y_7 (Fig. 1a). A high-throughput Gateway vector, pPs35SGW+npt3'-*chs*, was constructed by inserting the reading frame A (rfA) sequence (1700 nt EcoRV fragment of the vector pGemGWrfA; Karimi et al. 2002) into the vector pPs35Snpt3'-*chs* (cut with *PacI* and blunted with T4-DNA polymerase). The orientation of the inserted *rfA* fragment was checked by sequencing. The T-DNA vector pPs35SCAT800npt3'-*chs* (referred to as P35S-CAT2 800-*npt3'*-*chs*) was obtained through an LR clonase reaction between the vector pPs35SGW+npt3'-*chs* and the entry clone pENTR3'-CAT800 (obtained by a BP clonase reaction between pDONR and an *attB*-flanked PCR product containing the last 800 nt of the *A. thaliana* *CAT2*-coding sequence). After floral dip transformation with the Ps35SCAT800npt3'-*chs* T-DNA, plants harboring the single locus Y_{11} were obtained (see below; Fig. 1b). The T-DNA H610 (H T-DNA in De Buck et al. 1998, 2001) contained a *gus* gene fused to the 3'-UTR of the nopaline synthase gene (3'-*nos*) under control of P35S and the hygromycin acetyl transferase (*hpt*) gene conferring resistance to hygromycin. Plants homozygous for the Z_c locus (Fig. 1a) with a single copy of the T-DNA H610, were already available. All the T-DNA vectors were inserted into the *Agrobacterium tumefaciens* strain C58C1Rif^R (pMP90).

Plant material, crosses, and high-light treatment

All transformants were obtained via the floral dip procedure, performed as described by Clough and Bent (1998). Either

wild-type *A. thaliana* (L.) Heyhn. (ecotype Columbia) or transgenic plants homozygous for the Z_c locus ($-/-Z_cZ_c$ plants) containing a single copy of the T-DNA H610, were used for the dipping experiments. Seeds of the dipped plants were harvested and sown on K1 medium supplemented with kanamycin (50 mg L^{-1}) for selection of the DEchs287 T-DNA or phosphinothricin (10 mg L^{-1}) for selection of Ps35SGUSnpt3'chs and Ps35SCAT800npt3'chs. To obtain plants with a candidate X locus, $-/-Z_cZ_c$ plants were supertransformed with the DEchs287 T-DNA. Primary transformants ($X/-Z_cZ_c$) were retained with low amounts of the NPTII protein, indicative of silencing of the *nptII* gene(s) in the X locus, and with a high GUS activity, excluding transcriptional silencing of the *gus* gene in locus Z_c because of 35S promoter homology. These plants were screened for harboring a single locus and were further characterized through DNA gel blot analysis to determine the T-DNA integration pattern. Because the efficient in *trans*-silencing locus X in the tobacco XYZ system contained an IR about the right T-DNA border (Van Houdt et al. 2003), a supertransformant with the T-DNAs in an IR configuration (locus X_{21} , Fig. 1a), namely $X_{21}/-Z_cZ_c$, was chosen for further handling. To obtain $X_{21}/-/-$ plants, seeds from a self-fertilized $X_{21}/-Z_cZ_c$ plant (obtained after crossing $X_{21}/-Z_cZ_c$ with a $-/Y/-$ plant, not discussed here) were sown on K1 medium supplemented with kanamycin. The presence of locus Z_c was checked through a callus induction test on hygromycin-containing (20 mg L^{-1}) medium. Only one plant out of 52 kanamycin-resistant plants was hygromycin sensitive, suggesting that locus X_{21} and locus Z_c were genetically linked. To obtain candidate Y_7 and Y_{11} plants, $-/-Z_cZ_c$ plants were supertransformed with the Ps35SGUSnpt3'chs T-DNA, and wild-type plants were dipped with *A. tumefaciens* containing the T-DNA Ps35SCAT800npt3'chs, respectively. Fluorometric GUS assays and CAT activity assays were performed to select primary transformants with highly expressed Y transgenes, excluding silencing of the *gus* transgene of Z_c or cosuppression of the endogenous CAT genes. Via segregation analysis we ensured the presence of single loci, leading to the selection of transformants $-/Y_7/Z_cZ_c$ (Ps35SGUSnpt3'chs) and $-/Y_{11}/-$ (Ps35SCAT800npt3'chs). Transitive silencing was studied by crossing progeny plants of $X_{21}X_{21}/-Z_cZ_c$ with those of $-/Y_7/Z_cZ_c$ and $-/Y_{11}/-$. The resulting hybrid seeds were grown on medium selective for the presence of X_{21} and Y_7 or Y_{11} . After 2 wk, all plants were transferred to soil and grown under a light regime of 16 h day/8 h night, at 21°C . For high-light (HL) treatment, 4-wk-old plants were exposed to continuous HL irradiation ($\sim 1000 \mu\text{mol photons m}^{-2} \text{ sec}^{-1}$).

NPTII and GUS quantification

Proteins were extracted from two rosette leaves harvested 6 wk after sowing and frozen in liquid nitrogen. The protein concentrations, NPTII concentrations, and GUS activities were determined as described previously (Breyne et al. 1993; Van Houdt et al. 1997).

Catalase activity assay

Protein extracts were prepared by grinding two rosette leaves, frozen in liquid nitrogen, in $120 \mu\text{L}$ extraction buffer (60 mM Tris-HCl at pH 6.9, 1 mM phenylmethylsulfonylfluoride, 10 mM dithiothreitol, 20% glycerol). The homogenate was centrifuged

($15,000g$) twice at 4°C for 10 min to remove insoluble material. The supernatant was used for spectrophotometric catalase analysis according to Clare et al. (1984) after determining the total amount of soluble protein with the BioRad Protein Assay (Bradford 1976) with bovine serum albumin as a standard.

Quantitative real-time PCR analysis

For the quantification of *CAT1*, *CAT2*, and *CAT3* transcript levels by real-time PCR, total RNA from leaf material of 5-wk-old plants was prepared with TRIzol Reagent (Invitrogen). For each sample, $1 \mu\text{g}$ of DNase-treated total RNA was reverse-transcribed with the SuperScript First-Strand Synthesis System for RT-PCR (Invitrogen) according to the instruction manual. First-strand cDNA was generated with an oligo(dT)₁₂₋₁₈ primer. Of the first-strand cDNA, $0.2 \mu\text{L}$ was used as a template in a subsequent quantitative PCR using the qPCRTM Core Kit for SYBR[®] Green I (Eurogentec). The transcripts were amplified with gene-specific primers: 5'-TGCTCCTCTTACGGTTTGGTTTC-3' and 5'-ACCCTTCTTTAAGCGTTTCATTGTC-3' for *CAT1* (At1g20630); 5'-GCTGGCAAGCCGCTCTGAAC-3' and 5'-AGCACAGAAGATCCACATGATGAAG-3' for *CAT2* (At4g35090); 5'-GGCCAATCTCCATAAAGCTCAGT-3' and 5'-GGATTTAACGACCAAGCGATGATAG-3' for *CAT3* (At1g20620); 5'-AACTTGCTCATCTGCCATAGG-3' and 5'-TGATTCTGCGGAAACACCACTTTAG-3' for *actin2* (At3g60830).

Small-RNA analysis

Small RNAs were detected as described by Van Houdt et al. (2003), with minor changes. Low molecular weight RNA ($35 \mu\text{g}$) was loaded on the gel, as well as 21-nt and 24-nt RNA oligomers as size controls. The probe, a PCR fragment of 274 bp amplified with the primers 5'-CCGGTTAATTAATGCTGAGAAGTATCAACT-3' and 5'-CCGGTTAATTAACGGCTTGCCAGCTTCTGTCC-3', was labeled with [α -³²P]dCTP using the Rediprime II random prime labeling system (GE-Healthcare) according to the manufacturer's instructions. After removal of unincorporated [α -³²P]dCTPs with the Bio-Spin p30 Column (Bio-Rad Laboratories), the probe was hydrolyzed into fragments of ~ 50 nt. Hybridization and washes were performed as described at 50°C . Labeled membranes were exposed to a PhosphorImager screen (GE-Healthcare). Predominantly ethidium bromide-stained species of low molecular weight RNA, separated by agarose gel electrophoresis ($1 \mu\text{g}$ per lane) were used as loading controls.

ACKNOWLEDGMENTS

We thank Els Van Lerberge and Ellen Wirawan for technical assistance and Martine De Cock and Karel Spruyt for help with the manuscript and figures, respectively. This work was supported by grants from the European Union Biotech Project (no. QLRT-2000-00078). A.B. is a Research Fellow of the Research Foundation-Flanders and H.V.H. is indebted to the Institute for the Promotion of Innovation by Science and Technology in Flanders for a postdoctoral fellowship.

Received April 7, 2006; accepted June 28, 2006.

REFERENCES

- Alder, M.N., Dames, S., Gaudet, J., and Mango, S.E. 2003. Gene silencing in *Caenorhabditis elegans* by transitive RNA interference. *RNA* **9**: 25–32.
- Baulcombe, D. 2004. RNAi silencing in plants. *Nature* **431**: 356–363.
- Bleys, A., Van Houdt, H., and Depicker, A. 2006. Transitive and systemic RNA silencing: Both involving an RNA amplification mechanism? In *Small RNAs: Analysis and regulatory functions. Nucleic Acids and Molecular Biology* (eds. W. Nellen and C. Hammann), Vol. 17, pp. 119–139. Springer, Berlin.
- Bradford, M.M. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* **72**: 248–254.
- Braunstein, T.H., Moury, B., Johannessen, M., and Albrechtsen, M. 2002. Specific degradation of 3' regions of GUS mRNA in posttranscriptionally silenced tobacco lines may be related to 5'–3' spreading of silencing. *RNA* **8**: 1034–1044.
- Breyne, P., De Loose, M., Dedonder, A., Van Montagu, M., and Depicker, A. 1993. Quantitative kinetic analysis of β -glucuronidase activities using a computer-directed microtiter plate reader. *Plant Mol. Biol. Rep.* **11**: 21–31.
- Chamnongpol, S., Willekens, H., Moeder, W., Langebartels, C., Sandermann Jr., H., Van Montagu, M., Inzé, D., and Van Camp, W. 1998. Defense activation and enhanced pathogen tolerance induced by H₂O₂ in transgenic plants. *Proc. Natl. Acad. Sci.* **95**: 5818–5823.
- Chuang, C.-F. and Meyerowitz, E.M. 2000. Specific and heritable genetic interference by double-stranded RNA in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci.* **97**: 4985–4990.
- Clare, D.A., Duong, M.N., Darr, D., Archibald, F., and Fridovich, I. 1984. Effects of molecular oxygen on detection of superoxide radical with nitroblue tetrazolium and on activity stains for catalase. *Anal. Biochem.* **140**: 532–537.
- Clough, S.J. and Bent, A.F. 1998. Floral dip: A simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant J.* **16**: 735–743.
- Dat, J.F., Pellinen, R., Beeckman, T., Van De Cotte, B., Langebartels, C., Kangasjärvi, J., Inzé, D., and Van Breusegem, F. 2003. Changes in hydrogen peroxide homeostasis trigger an active cell death process in tobacco. *Plant J.* **33**: 621–632.
- De Buck, S., Jacobs, A., Van Montagu, M., and Depicker, A. 1998. *Agrobacterium tumefaciens* transformation and cotransformation frequencies of *Arabidopsis thaliana* root explants and tobacco protoplasts. *Mol. Plant Microbe Interact.* **11**: 449–457.
- De Buck, S., Van Montagu, M., and Depicker, A. 2001. Transgene silencing of invertedly repeated transgenes is released upon deletion of one of the transgenes involved. *Plant Mol. Biol.* **46**: 433–445.
- Frugoli, J.A., Zhong, H.H., Nuccio, M.L., McCourt, P., McPeck, M.A., Thomas, T.L., and McClung, C.R. 1996. Catalase is encoded by a multigene family in *Arabidopsis thaliana* (L.) Heynh. *Plant Physiol.* **112**: 327–336.
- García-Pérez, R.D., Van Houdt, H., and Depicker, A. 2004. Spreading of post-transcriptional gene silencing along the target gene promotes systemic silencing. *Plant J.* **38**: 594–602.
- Hamilton, A., Voinnet, O., Chappell, L., and Baulcombe, D. 2002. Two classes of short interfering RNA in RNA silencing. *EMBO J.* **21**: 4671–4679.
- Han, Y. and Grierson, D. 2002. The influence of inverted repeats on the production of small antisense RNAs involved in gene silencing. *Mol. Genet. Genomics* **267**: 629–635.
- Himber, C., Dunoyer, P., Moissiard, G., Ritzenthaler, C., and Voinnet, O. 2003. Transitivity-dependent and -independent cell-to-cell movement of RNA silencing. *EMBO J.* **22**: 4523–4533.
- Karimi, M., Inzé, D., and Depicker, A. 2002. GATEWAYTM vectors for *Agrobacterium*-mediated plant transformation. *Trends Plant Sci.* **7**: 193–195.
- Kerschen, A., Napoli, C.A., Jorgensen, R.A., and Muller, A.E. 2004. Effectiveness of RNA interference in transgenic plants. *FEBS Lett.* **566**: 223–228.
- Klahre, U., Crété, P., Leuenberger, S.A., Iglesias, V.A., and Meins Jr., F. 2002. High molecular weight RNAs and small interfering RNAs induce systemic posttranscriptional gene silencing in plants. *Proc. Natl. Acad. Sci.* **99**: 11981–11986.
- Kościńska, E., Kalantidis, K., Wypijewski, K., Sadowski, J., and Tabler, M. 2005. Analysis of RNA silencing in agroinfiltrated leaves of *Nicotiana benthamiana* and *Nicotiana tabacum*. *Plant Mol. Biol.* **59**: 647–661.
- Lipardi, C., Wei, Q., and Paterson, B.M. 2001. RNAi as random degradative PCR: siRNA primers convert mRNA into dsRNAs that are degraded to generate new siRNAs. *Cell* **107**: 297–307.
- Mette, M.F., van der Winden, J., Matzke, M.A., and Matzke, A.J.M. 1999. Production of aberrant promoter transcripts contributes to methylation and silencing of unlinked homologous promoters. *in trans. EMBO J.* **18**: 241–248.
- Miki, D., Itoh, R., and Shimamoto, K. 2005. RNA silencing of single and multiple members in a gene family of rice. *Plant Physiol.* **138**: 1903–1913.
- Napoli, C., Lemieux, C., and Jorgensen, R. 1990. Introduction of a chimeric chalcone synthase gene into petunia results in reversible co-suppression of homologous genes *in trans*. *Plant Cell* **2**: 279–289.
- Nicolás, F.E., Torres-Martínez, S., and Ruiz-Vázquez, R.M. 2003. Two classes of small antisense RNAs in fungal RNA silencing triggered by non-integrative transgenes. *EMBO J.* **22**: 3983–3991.
- Petersen, B.O. and Albrechtsen, M. 2005. Evidence implying only unprimed RdRP activity during transitive gene silencing in plants. *Plant Mol. Biol.* **58**: 575–583.
- Rossi, J.J. 2005. RNAi and the P-body connection. *Nat. Cell Biol.* **7**: 643–644.
- Sanders, M., Maddelein, W., Depicker, A., Van Montagu, M., Cornelissen, M., and Jacobs, J. 2002. An active role for endogenous β -1,3-glucanase genes in transgene-mediated co-suppression in tobacco. *EMBO J.* **21**: 5824–5832.
- Schiebel, W., Haas, B., Marinković, S., Klanner, A., and Sanger, H.L. 1993. RNA-directed RNA polymerase from tomato leaves. II. Catalytic *in vitro* properties. *J. Biol. Chem.* **268**: 11858–11867.
- Sijen, T., Fleenor, J., Simmer, F., Thijssen, K.L., Parrish, S., Timmons, L., Plasterk, R.H.A., and Fire, A. 2001. On the role of RNA amplification in dsRNA-triggered gene silencing. *Cell* **107**: 465–476.
- Tang, G., Reinhart, B.J., Bartel, D.P., and Zamore, P.D. 2003. A biochemical framework for RNA silencing in plants. *Genes & Dev.* **17**: 49–63.
- Vaistij, F.E., Jones, L., and Baulcombe, D.C. 2002. Spreading of RNA targeting and DNA methylation in RNA silencing requires transcription of the target gene and a putative RNA-dependent RNA polymerase. *Plant Cell* **14**: 857–867.
- Van Houdt, H., Ingelbrecht, I., Van Montagu, M., and Depicker, A. 1997. Post-transcriptional silencing of a neomycin phosphotransferase II transgene correlates with the accumulation of unproductive RNAs and with increased cytosine methylation of 3' flanking regions. *Plant J.* **12**: 379–392.
- Van Houdt, H., Bleys, A., and Depicker, A. 2003. RNA target sequences promote spreading of RNA silencing. *Plant Physiol.* **131**: 245–253.
- Vandenabeele, S., Vanderauwera, S., Vuylsteke, M., Rombauts, S., Langebartels, C., Seidlitz, H.K., Zabeau, M., Van Montagu, M., Inzé, D., and Van Breusegem, F. 2004. Catalase deficiency drastically affects gene expression induced by high light in *Arabidopsis thaliana*. *Plant J.* **39**: 45–58.
- Zilberman, D., Cao, X., and Jacobsen, S.E. 2003. ARGONAUTE4 control of locus-specific siRNA accumulation and DNA and histone methylation. *Science* **299**: 716–719.

RNA

A PUBLICATION OF THE RNA SOCIETY

Down-regulation of endogenes mediated by a transitive silencing signal

Annick Bleys, Helena Van Houdt and Anna Depicker

RNA 2006 12: 1633-1639

References

This article cites 38 articles, 17 of which can be accessed free at:
<http://rnajournal.cshlp.org/content/12/9/1633.full.html#ref-list-1>

License

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).

To subscribe to RNA go to:
<http://rnajournal.cshlp.org/subscriptions>