REPORT

The *Saccharomyces cerevisiae* Pus2 protein encoded by *YGL063w* ORF is a mitochondrial tRNA:Ψ^{27/28}-synthase

ISABELLE BEHM-ANSMANT, CHRISTIANE BRANLANT, and YURI MOTORIN

Laboratoire de Maturation des ARN et Enzymologie Moléculaire, UMR 7567 CNRS-UHP Nancy I, Faculté des Sciences, BP 239, 54506 Vandoeuvre-les-Nancy Cedex, France

ABSTRACT

The RNA:pseudouridine (Ψ)-synthase family is one of the most complex families of RNA modification enzymes. Ten genes encoding putative RNA:Ψ-synthases have been identified in *S. cerevisiae*. Most of the encoded enzymes have been characterized experimentally. Only the putative RNA:Ψ-synthase Pus2p (encoded by the *YGL063w* ORF) had no identified substrate. Here, we analyzed Ψ residues in cytoplasmic and mitochondrial tRNAs extracted from *S. cerevisiae* strains, carrying disruptions in the *PUS1* and/or *PUS2* ORFs. Our results demonstrate that Pus2p is a mitochondrial-specific tRNA:Ψ-synthase acting at positions 27 and 28 in tRNAs. The importance of the Asp56 residue in the conserved ARTD motif of the Pus2p catalytic site is demonstrated in vivo. Interestingly, in spite of the absence of a characteristic N-terminal targeting signal, our data strongly suggest an efficient and rapid targeting of Pus2p in yeast mitochondria. In contradiction with the commonly held idea that a unique nuclear gene encodes the enzyme required for both cytoplasmic and mitochondrial tRNA modifications, here we show the existence of an enzyme specifically dedicated to mitochondrial tRNA modification (Pus2p), the corresponding modification in cytoplasmic tRNAs being catalyzed by another protein (Pus1p).

Keywords: *S. cerevisiae*; pseudouridine; RNA:pseudouridine-synthase; tRNAs; mitochondria

INTRODUCTION

In all organisms, pseudouridine (Ψ) and 2'-O-methylated residues are the most frequent modifications in RNAs (see The RNA Modification Database, URL <http://medstat.med.utah.edu/RNAmods/>). The Ψ residues are formed by base isomerization of the transcriptionally incorporated uridine residues (Kammen et al. 1988; Huang et al. 1997). This reaction is catalyzed by RNA:Ψ-synthases, which do not require cofactors or energy for catalysis. RNA:Ψ-synthases represent a broad group of proteins that are divided into five distinct families (TruA, TruB, RluA, RsuA, and TruD) on the basis of amino acid sequence homology (see Koonin 1996; Conrad et al. 1998; Kaya and Ofengand 2003). Based

on computer analysis of bacterial and yeast genomes, nine genes encoding putative RNA:Ψ-synthases have been identified in *Saccharomyces cerevisiae* (Koonin 1996). The RNA:Ψ-synthase activity of the products of eight of them has been identified. Only the substrate specificity of the *PUS2* gene product has not yet been characterized. RNA:Ψ-synthases Pus3p, Pus4p, and Pus6p are tRNA-specific enzymes acting on both cytoplasmic and mitochondrial tRNAs (Becker et al. 1997; Lecoite et al. 1998; Ansmant et al. 2001). Pus5p is responsible for the modification of the mitochondrial 21S rRNA (Ansmant et al. 2000), and Cbf5p is associated with the H/ACA snoRNA guides and catalyzes Ψ formation in cytoplasmic rRNAs and U2 snRNA (Lafontaine et al. 1998; Kiss et al. 2004; Ma et al. 2005). Pus7p was not identified using sequence homology but by genome-wide screening of GST-tagged yeast ORFs. It was initially characterized as a U2 snRNA-specific Ψ-synthase (Ma et al. 2003), but it also catalyzes tRNA modification at positions 13 and 35 (Behm-Ansmant et al. 2003). We found that two distinct enzymes, Rib2p/Pus8p and Pus9p, are required for Ψ³² formation in tRNAs, depending on their cytoplasmic or mitochondrial localization, and cytoplasmic Rib2p/Pus8p contains both a tRNA:Ψ³²-synthase domain and a DRAP-deaminase domain (Behm-Ansmant et al.

Abbreviations: 2D, bidimensional; CMCT, 1-cyclohexyl-3-[2-morpholinoethyl] carbodiimide metho-*p*-toluenesulfonate; NMP, nucleotide monophosphate; Ψ, pseudouridine; ORF, open reading frame; PCR, polymerase chain reaction; RT, reverse transcriptase; TLC, thin layer chromatography; U, uridine; WT, wild-type.

Reprint requests to: Yuri Motorin, Laboratoire Maturation des ARN et Enzymologie Moléculaire, UMR 7567 CNRS-UHP Nancy I, Faculté des Sciences, BP 23954506 Vandoeuvre-les-Nancy Cedex, France; e-mail: Yuri.Motorin@maem.uhp-nancy.fr; Fax 33-3-83-68-43-07.

Article published online ahead of print. Article and publication date are at <http://www.rnajournal.org/cgi/doi/10.1261/rna.605607>.

2004). Pus1p is an RNA: Ψ -synthase with a multisite specificity (tRNA positions 1, 26, 27, 28, 34, 36, 65, and 67) (Motorin et al. 1998; Behm-Ansmant et al. 2006). In addition, it also modifies U2 snRNA at position 44 (Massenet et al. 1999). In contrast to yeast cytoplasmic tRNAs, their mitochondrial counterparts are significantly less modified; only uridines at positions 27, 28, 31, 32, 38, 39, 55, and 72 may be converted to Ψ s. Based on the assumption that a unique enzyme is generally used to modify both cytoplasmic and mitochondrial tRNAs, Pus1p was also expected to be responsible for the formation of residues Ψ 27 and Ψ 28 commonly present in mitochondrial tRNAs. However, its activity in yeast mitochondria was never tested experimentally. The only remaining RNA: Ψ -synthase without identified substrates (Pus2p) shows strong sequence homology with Pus1p, especially in motifs II and IIa (Fig. 1), hinting at a similar catalytic activity for these two enzymes. Furthermore, in spite of the absence of a characteristic motif for mitochondrial localization in Pus2p, its preferential mitochondrial localization is clearly predicted by the NNPSL software (Reinhardt and Hubbard 1998). Thus, we hypothesized that Pus2p may be the mitochondrial counterpart of Pus1p. Our interest in the discovery of the yeast RNA: Ψ -synthase responsible for modification of mitochondrial tRNAs at positions 27 and 28 was reinforced by the discovery of a direct link between Ψ 27 formation in human mitochondrial tRNA and a rare mitochondrial disorder called mitochondrial myopathy and sideroblastic anemia (MLASA) (Patton et al. 2005).

To test this hypothesis, we analyzed the effects of individual or simultaneous deletions of *PUS1* and *PUS2* ORFs on the in vivo tRNA pseudouridylation pattern. The loss of *PUS1* did not affect pseudouridylation at positions 27 and 28 in mitochondrial tRNAs, while the deletion of *PUS2* abolished Ψ 27 and Ψ 28 formation in these species. Since the enzyme responsible for Ψ 72 formation in the

mitochondrial tRNA^{Met}_i(CAU) has not yet been identified, we also tested the possible activity of Pus2p at position 72 of this tRNA. The data presented demonstrate that Pus2p is a yeast mitochondrial RNA: Ψ -synthase acting at positions 27 and 28, but not at position 72.

RESULTS AND DISCUSSION

Mitochondrial tRNA modifications are affected by *PUS2* ORF disruption

To investigate whether Ψ formation at positions 27 and 28 in mitochondrial tRNAs is catalyzed by Pus1p or Pus2p, we tested the effect of individual or simultaneous *PUS1* and *PUS2* ORF deletions on mitochondrial tRNA pseudouridylation. The haploid *S. cerevisiae* strain RS453, whose *PUS1* ORF was disrupted by insertion of the *HIS3* gene (*pus1 Δ ::HIS3*, denoted Δ *pus1*) (Simos et al. 1996), was used to test for the effect of the absence of Pus1p expression. To analyze the consequences of the absence of Pus2p expression, we used the haploid *S. cerevisiae* strain BY4742, whose *PUS2* ORF is disrupted by a *Kan^r* gene replacement (*pus2 Δ ::Kan^r*, denoted Δ *pus2*; EUROSCARF). The effects of the simultaneous *PUS1* and *PUS2* ORF disruptions were tested in the haploid Δ *pus1*/ Δ *pus2* strain, which we isolated by mating of the *pus2 Δ ::Kan^r* and the *pus1 Δ ::HIS3* strains and dissecting the resulting diploids.

Six out of the 17 sequenced mitochondrial tRNAs in *S. cerevisiae* bear a Ψ residue at position 27. However, four of them are not suitable for analysis by the CMCT/RT approach because of the presence of a m²G26 or/and m¹G37 residue(s), which block(s) primer extension analysis. To avoid this difficulty when testing the Pus1p and Pus2p activity at position 27 we used tRNA^{Arg}(UCU), which does not contain these modifications at positions 26 and 37. Pseudouridine at position 28 is found in only

FIGURE 1. Alignment of the amino acid sequences of the *Escherichia coli* TruA and *S. cerevisiae* Pus1p, Pus2p, and Pus3p tRNA: Ψ -synthases of the TruA family. (A) Proteins are schematically represented as bars, whose lengths are proportional to lengths of the polypeptide chains. Motifs I, II, IIa, III, and IIIa of RNA: Ψ -synthases (Del Campo et al. 2004) are represented by black or gray rectangles. (B) The alignments of the amino acid sequences of the conserved blocks I to IIIa are shown. Identical and semiconserved residues are boxed. The thickness of the line is an indication of the level of sequence conservation. The conserved residues (D, Y, I, R, L) in the tRNA: Ψ -synthases active sites are indicated by asterisks.

one mitochondrial tRNA, the tRNA^{Lys}(UUU), and fortunately, Ψ 28 formation in this tRNA could also be analyzed by the CMCT/RT approach. Disruption of the *PUS1* ORF affected neither Ψ 27 formation in the mitochondrial tRNA^{Arg}(UCU) (Fig. 2A, lanes 5,6) nor Ψ 28 formation in mitochondrial tRNA^{Lys}(UUU) (Fig. 2B, lanes 3,4). In contrast, after deletion of *PUS2* ORF, both of these pseudouridylations were abolished (Fig. 2A,B, lanes 7, 8 in both panels). Accordingly, similar results were obtained with the $\Delta pus1/\Delta pus2$ strain carrying both deletions (Fig. 2A,B, panels A and B, lanes 9,10 and 11,12, respectively). Transformation of this strain with a plasmid expressing the WT protein restored Ψ 27 formation in tRNA^{Arg}(UCU),

while no modification was detected in the strain expressing a variant protein with a D56A substitution in Motif II (Fig. 3, lanes 7–10).

Since Ψ 72 in tRNA^{Met}_i(CAU) is located too close to the tRNA 3'-extremity to be analyzed by the CMCT/RT approach, the possible implication of Pus2p in Ψ 72 formation was only tested in vitro by the nearest-neighbor approach. To this end, the tRNA: Ψ 72-synthase activity of cell-free extracts prepared from different strains was tested on a transcript of the mitochondrial tRNA^{Met}_i(CAU) labeled by [α -³²P]ATP incorporation. As evidenced by thin layer chromatography (TLC) analysis of the labeled RNase T2 digestion products, none of the gene disruptions affected the activity of the extracts (Table 1, see also Behm-Ansmant et al. 2006). Thus, Pus1p and Pus2p are probably not involved in Ψ formation at position 72.

Other pseudouridines in cytoplasmic yeast tRNA normally formed by Pus1p (positions 1, 34, 36, 65, 67) are absent in the mitochondrial compartment; thus, positions 27 and 28 are the only possible targets for Pus2p in mitochondria.

Pus2p is unable to insure cytoplasmic tRNA modification

Since Pus1p and Pus2p have redundant activity toward Ψ 27 and Ψ 28, we tested if Pus2p could complement the absence of active Pus1p for cytoplasmic tRNA modification. Previous studies have already demonstrated the absence of tRNA: Ψ 27-synthase activity in extracts from the $\Delta pus1$ strain (Motorin et al. 1998). However, this apparent absence of in vitro Pus2p activity could result from protein inactivation or from inappropriate reaction conditions. We therefore verified the absence of in vivo Pus2p activity on cytoplasmic tRNAs by CMCT/RT analysis of the cytoplasmic tRNA^{Met}(CAU) extracted from the $\Delta pus1$ strain. As shown in Figure 4, lanes 7, 8,15, and 16, no Ψ residue was detected at position 27 in this tRNA, whereas it was detected in tRNA^{Met}(CAU) extracted from the $\Delta pus2$ strain (Fig. 4A, lanes 11,12). Formation of Ψ 31, which was used as a control, remained unaffected. Thus, Pus2p is not able to complement the absence of Pus1p activity in vivo.

How to explain the inability of Pus2p to modify cytoplasmic tRNAs? Either Pus2p cannot modify cytoplasmic tRNAs due to the absence of the required recognition determinants in these tRNAs or, immediately after protein synthesis, Pus2p is rapidly and predominantly targeted to mitochondria and thus its potential substrates in the cytoplasm escape modification. The first hypothesis seems unlikely since in contrast to their vertebrates counterparts, yeast cytoplasmic and mitochondrial tRNAs have rather similar 2D structures (Wesolowski and Fukuhara 1979; Leehey et al. 1995).

To validate the second hypothesis, we compared the in vitro tRNA: Ψ 27-synthase activities of extracts from the WT, $\Delta pus1$, $\Delta pus2$, and $\Delta pus1/\Delta pus2$ cells on a mitochondrial

FIGURE 2. Disruption of the *PUS2* gene leads to the absence of Ψ 27 residues in the mitochondrial tRNA^{Arg}(UCU) (A) and Ψ 28 in the mitochondrial tRNA^{Lys}(UUU) (B). Total RNAs extracted from the WT, $\Delta pus1$, $\Delta pus2$, and $\Delta pus1/\Delta pus2$ strain were modified by CMCT, for 2, 10, and 20 min with (+) or without (–) subsequent alkaline treatment (OH[–]). A control experiment was performed in the absence of CMCT treatment. Lanes U, G, C, and A correspond to the sequencing ladders obtained with the same oligonucleotide. Primer extension analyses were performed on the mitochondrial tRNA^{Arg}(UCU) (A) and tRNA^{Lys}(UUU) (B) using specific oligonucleotide primers. The reverse transcription stops, corresponding to residue Ψ 27 in tRNA^{Arg}(UCU) and Ψ 28 in tRNA^{Lys}(UUU), are indicated by arrows.

FIGURE 3. The aspartic acid residue D56 in the Pus2p active site is essential for the RNA:Ψ-synthase activity. The yeast *S. cerevisiae* $\Delta pus1/\Delta pus2$ strain was transformed with recombinant p416GalS plasmids, expressing the WT (D56) or mutated (D56A) PUS2 ORF. Total RNAs were extracted from the WT, the untransformed $\Delta pus1/\Delta pus2$, and the two transformed $\Delta pus1/\Delta pus2$ strains. The mitochondrial tRNA^{Arg}(UCU) was analyzed by the CMCT/RT approach (same legend as in Fig. 2). The reverse transcription stop, corresponding to residue Ψ27 is indicated by an arrow.

tRNA known to be modified at position 27 (tRNA^{His}(GUG)) and on a cytoplasmic tRNA that has Ψ residues at both positions 27 and 55 (tRNA^{Val}(UAC)). The analysis was performed using the nearest neighbor approach. To this end, the mitochondrial tRNA^{His}(GUG) and the cytoplasmic tRNA^{Val}(UAC) were labeled by [α -³²P]ATP and [α -³²P]CTP incorporation, respectively. The results obtained are given in Table 1. Formation of Ψ residues were detected at positions 27 and 55 in the cytoplasmic tRNA^{Val}(UAC) and at position 27 in the mitochondrial tRNA^{His}(GUG), after incubation with the WT and the $\Delta pus2$ cell extracts. In contrast, residue Ψ27 was not formed in any of these two tRNAs after incubation in both the $\Delta pus1$ and the $\Delta pus1/\Delta pus2$ cell extracts. The following interpretation can be proposed: (1) Pus1p in the extract has

an intrinsic capability of modifying mitochondrial as well as cytoplasmic tRNAs, and (2) the absence of tRNA:Ψ27-synthase activity on both the cytoplasmic and the mitochondrial tRNA transcripts in the $\Delta pus1$ extract is due to the absence of Pus2p in the cellular extract.

The tight association of Pus2p with the insoluble fraction of disrupted yeast cells, which probably contain the mitochondria, can explain its almost complete loss during preparation of cell-free extract and the observed absence of Pus2p activity in these extracts. Indeed, by expression of an HA-tagged Pus2p in the *S. cerevisiae* strain 26-G-7 (Open BioSystems), we showed by Western Blot analysis that the S10 cell-free soluble extract is devoid of HA-tagged Pus2p (Fig. 4B).

Taken together, our results show that Pus1p is able to modify both the cytoplasmic and mitochondrial tRNAs in vitro. However, it has a strict cytoplasmic/nuclear localization and therefore only modifies cytoplasmic tRNAs in vivo. In contrast, Pus2p is very rapidly targeted to mitochondria and is not present in a yeast soluble cell fraction. This rapid transport to mitochondria likely explains its absence of activity on cytoplasmic tRNAs.

Nonconventional mitochondrial localization signal in Pus2p

Taken together, our data suggested that Pus2p is efficiently targeted to mitochondria after its synthesis. This was rather surprising since no clear mitochondrial signal peptide was found at the N-terminus of Pus2p. Moreover, among the *S. cerevisiae* RNA:Ψ-synthases acting in mitochondria (Pus3p, Pus4p, Pus5p, and Pus9p), only Ydl036p (Pus9p) has a characteristic N-terminal targeting signal (Mason 1998; Behm-Ansmant et al. 2004). All other mitochondrial yeast tRNA:Ψ-synthases, although efficiently transported to this compartment, have no characteristic N-terminal signals (Becker et al. 1997; Lecointe et al. 1998; Ansmant et al. 2000, 2001). Nevertheless, based on their amino acid sequences, all of them are predicted to have a predominant mitochondrial localization by the PSORTII software.

On the other hand, a new mechanism was recently found to allow the efficient targeting of proteins without characteristic N-terminal signals to mitochondria. It is based on the presence of a sequence element in the 3'-UTR of

TABLE 1. Pseudouridine formation in selected yeast tRNA transcripts catalyzed by extracts of the $\Delta pus1$, $\Delta pus2$, and $\Delta pus1/\Delta pus2$ yeast strains

Position tested in tRNA	tRNA transcript/labeling	WT strain	$\Delta pus1$ strain	$\Delta pus2$ strain	$\Delta pus1/\Delta pus2$ strain
27 + 55	Yeast cytoplasmic tRNA ^{Val} (UAC)/CTP	1.36	0.71	1.42	0.80
27	Yeast mitochondrial tRNA ^{His} (GUG)/ATP	0.61	0.01	0.57	0.02
72	Yeast mitochondrial tRNA ^{Met} (CAU)/ATP	0.78	0.83	0.72	0.78

Radiolabeled transcripts were incubated with yeast extract for 60 min at 37°C, and the Ψ content was analyzed by 2D TLC after RNase T2 digestion.

FIGURE 4. Formation of residue Ψ 27 in the cytoplasmic tRNA^{Met} (CAU) is not affected by disruption of the *PUS2* ORF. (A) Total RNAs extracted from the WT, Δ *pus1*, Δ *pus2*, and Δ *pus1/Δpus2* strains were subjected to CMCT modification (same legend as in Fig. 2). The presence of residues Ψ 27 and Ψ 31 in tRNA^{Met}(CAU) was tested by primer extension using the oligonucleotide described in Materials and Methods. A sequence ladder was made with the same oligonucleotide. Arrows indicate the reverse transcription stops, corresponding to residues Ψ 27 and Ψ 31. (B) Western blot analysis of HA-tagged *Pus2p* in yeast *S. cerevisiae* strain 26-G-7 (Open BioSystems). Yeast cells lysed by SDS (SDS lysate), insoluble fraction (P10), and S10 extract (S10) were loaded onto 10% PAAG, and separated proteins were transferred to nitrocellulose membrane. Staining was done using monoclonal anti-HA antibodies (12CA5, Roche) and ECL Plus kit (GE Healthcare). Molecular weights of marker proteins are indicated on the left of the panel.

mRNAs that allows mRNA concentration and translation in close vicinity of mitochondria (for review, see Jensen and Johnson 2001; Tekotte and Davis 2002). We can exclude the utilization of this mechanism for rapid transport of *Pus2p* to mitochondria. Indeed, the *PUS2* ORF that we expressed in the Δ *pus2* cells was followed by a foreign 260-base pair (bp) DNA sequence (the CYC1-terminator of p416GalS plasmid), instead of the authentic 3'-UTR of the *PUS2* gene. Despite this, the expressed *Pus2p* efficiently complemented the mitochondrial tRNA: Ψ 27-synthase activity in Δ *pus2* strain. This indicates that *Pus2p* probably uses another mechanism for efficient transport into mitochondria.

Two distinct RNA: Ψ -synthases act at positions 27 and 28 in tRNAs

Based on several previous studies, up to now it has been accepted that the modification of a given position in tRNAs is ensured by the product(s) of a unique gene in both the cytoplasmic and mitochondrial compartments (Martin and Hopper 1994; Rose et al. 1995; Becker et al. 1997; Lecointe et al. 1998; Ansmant et al. 2001). Either a unique protein is expressed from the gene and it has a dual cytoplasmic and mitochondrial localization (Becker et al. 1997; Lecointe et al. 1998; Ansmant et al. 2001), or two distinct enzymes differing by the presence of an N-terminal mitochondrial targeting signal in one of them are produced from the unique gene by the alternative utilization of AUG initiation codons (Martin and Hopper 1994; Rose et al. 1995). The only reported exception is the Rib2p/*Pus8p* and *Pus9p* pair of tRNA: Ψ -synthases: Rib2p/*Pus8p* is a cytoplasmic tRNA: Ψ 32-synthase, while *Pus9p* can act both in the cytoplasm and mitochondrial compartments with a predominant activity in mitochondria (Behm-Ansmant et al. 2004). Here we show that another pair of RNA: Ψ -synthases is involved in the modification of tRNA positions 27 and 28. Nevertheless, the situation is quite different from the Rib2p/*Pus8p*-*Pus9p* pair, since the *Pus1p* activity is strictly cytoplasmic, while *Pus2p* activity is strictly mitochondrial.

One possible explanation for the presence of a specific mitochondrial enzyme in addition to *Pus1p* may be a preferential localization of *Pus1p* in the nucleus, where it modifies the tRNA^{Ile}(UAU) precursor (Simos et al. 1996; Motorin et al. 1998) and U2 snRNA (Massenet et al. 1999). In addition, *Pus1p* participates in tRNA export from the nucleus in the cytoplasm (Simos et al. 1996; Grosshans et al. 2001). These various functions of *Pus1p* may be incompatible with its additional mitochondrial localization and may explain the necessity of a dedicated mitochondrial enzyme with a similar substrate specificity (*Pus2p*).

Although conserved in yeast, the *PUS1* and *PUS2* genes may be deleted without marked growth phenotype

Another question is to know whether the existence of two redundant tRNA: Ψ -synthases with different sites of action is evolutionarily conserved. A search in genomic databanks of genes that may encode *Pus1p* and *Pus2p* homologs revealed the occurrence of genes encoding these two proteins only in yeasts that belong to the *Saccharomycetoidae* and the *Schizosaccharomycetoidae* families. They have a strong similarity with both *Pus1p* and *Pus2p*. However, only one gene that may encode a *Pus1p*/*Pus2p* homolog was found in vertebrates and other eukaryotes (Chen and Patton 1999, 2000). In relation to this observation, it must be pointed out that in yeasts, modification at position 44 in U2 snRNA is catalyzed by *Pus1p* (Massenet et al. 1999), whereas

a scaRNA guided system is used in vertebrates (Ma et al. 2005). The emergence of the scaRNA guided modification system for UsnRNAs probably had a strong influence on the activity of members of the RNA:Ψ-synthase families. The striking similarity between Pus1p and Pus2p suggests that the presence of two distinct enzymes may have arisen from gene duplication, followed by acquisition of the appropriate subcellular targeting signals.

Despite our observation of nonoverlapping functions of Pus1p and Pus2p, individual and even simultaneous disruptions of their genes had no marked effect on cell viability and growth, even in growth conditions allowing detection of mitochondrial respiratory defects (medium with glycerol as the only carbon source, data not shown). The absence of marked growth phenotype is a rather general property of strains carrying disruption in a gene encoding an RNA modification enzyme. When growth phenotypes were detected, they were almost always due to the lack of an additional protein function distinct from the RNA modification activity.

MATERIALS AND METHODS

Yeast strains

The haploid *S. cerevisiae* BY4742 strain (denoted WT in this article) and the haploid strain carrying a disruption of the *PUS2* gene (*pus2Δ::Kan^r*, denoted Δ *pus2*), were obtained from the EUROSCARF collection (Germany). The haploid Δ *pus1* yeast strain was a generous gift of G. Simos (University of Larissa, Larissa, Greece). It was obtained by interruption of the *PUS1* gene by the *HIS3* auxotrophic marker (*pus1Δ::HIS3*) in the genomic DNA of the RS453 *S. cerevisiae* strain (Simos et al. 1996). Yeast *S. cerevisiae* strain 26-G-7 expressing 3× HA-tagged Pus2p (insertion position 312) was obtained from Open BioSystems. A haploid strain with simultaneous deletions of the *PUS1* and *PUS2* genes was generated by mating of the *pus2Δ::Kan^r* and the *pus1Δ::HIS3* strains. After sporulation of the obtained diploid mutants, asci were separated by microdissection.

Complementation of the disrupted strains

To generate a *S. cerevisiae* plasmid expressing Pus2p, the coding region of the *PUS2* ORF together with the upstream 300-bp region were amplified from the genomic DNA of the *S. cerevisiae* BY4742 strain. The oligonucleotides used generated a *NheI* and an *EcoRI* restriction site at the 5'- and 3'-extremities, respectively. The amplified DNA fragment was inserted at the *SmaI* site of plasmid pUC18. The resulting construct (pUC18-PUS2) was cleaved by the *NheI* and *EcoRI* restriction endonucleases, and the fragment encoding Pus2p was inserted between the *XbaI* and *EcoRI* sites of plasmid p416GalS (Mumberg et al. 1994). A point mutation in the putative RNA:Ψ-synthase active site of Pus2p was introduced by PCR-mediated site-directed mutagenesis (D56A, GAC→GCC). All the recombinant genes produced were entirely sequenced. Yeast transformation was performed with the standard lithium acetate procedure (Gietz et al. 1992; Adams et al. 1997).

CMCT/RT mapping of Ψ residues in RNA

Preparation of total RNA from yeast strains and CMCT-RT mapping of Ψ residues were performed as described previously (Bakin and Ofengand 1993; Massenet et al. 1999; Ansmant et al. 2000). The presence of residue Ψ27 in mitochondrial tRNA^{Arg}(UCU), Ψ28 in mitochondrial tRNA^{Lys}(UUU), and Ψ27 in cytoplasmic tRNA^{Met}(CAU) was analyzed using oligonucleotides complementary to residues 46–62, residues 43–72, and residues 40–57, respectively.

Analysis of Ψ residue formation by the nearest neighbor approach

The coding sequences of the mitochondrial tRNA^{His}(GUG) and tRNA^{Met}i(CAU) were PCR amplified from the genomic DNA of the BY4742 *S. cerevisiae* strain and inserted downstream from a T7 RNA polymerase promoter at the *SmaI* site of plasmid pUC18. The pTFM-Val plasmid used for in vitro transcription of cytoplasmic tRNA^{Val}(UAC) was kindly provided by C. Florentz (IBMC, Strasbourg, France). Cell-free S10 extracts preparation and tests of their in vitro RNA:Ψ-synthase activities were performed as previously described (Auxilien et al. 1996; Behm-Ansmant et al. 2003).

Sequence analysis

The NNPSL software (Reinhardt and Hubbard 1998, <http://predict.sanger.ac.uk/nnpsl/>, Web site) was used for prediction of the Pus2p subcellular localization. This program takes into account the amino acid composition of the whole protein, its predicted secondary structure, and various characteristic features, such as the presence of a mitochondrial targeting sequence or of a potential signal peptidase cleavage site.

ACKNOWLEDGMENTS

We thank H. Grosjean (LEBS, Gif-sur-Yvette, France) for helpful discussions, R. Giegé and C. Florentz (CNRS, Strasbourg, France) for giving us the plasmid containing the yeast tRNA^{Val}(UAC) gene, A. Lescure (CNRS, Strasbourg, France) for help in yeast dissection, and G. Simos (University of Larissa, Larissa, Greece) for the gift of the *pus1Δ::HIS3* yeast strain and helpful discussions. This work was supported by laboratory funds from the “Ministère de l'Éducation Nationale, de l'Enseignement Supérieur et de la Recherche” (MENESR) and the French “Centre National de la Recherche Scientifique”. I.B.-A. was a predoctoral fellow supported by a fellowship from the MENESR.

Received April 4, 2007; accepted July 10, 2007.

REFERENCES

- Adams, A., Gottschling, D.E., Kaiser, C.A., and Stearns, T. 1997. *Methods in yeast genetics*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Ansmant, I., Massenet, S., Grosjean, H., Motorin, Y., and Branlant, C. 2000. Identification of the *Saccharomyces cerevisiae* RNA:pseudouridine synthase responsible for formation of Ψ₂₈₁₉ in 21S mitochondrial ribosomal RNA. *Nucleic Acids Res.* **28**: 1941–1946.

- Ansmant, I., Motorin, Y., Massenet, S., Grosjean, H., and Branlant, C. 2001. Identification and characterization of the tRNA: Ψ ₃₁-synthase (Pus6p) of *Saccharomyces cerevisiae*. *J. Biol. Chem.* **276**: 34934–34940.
- Auxilien, S., Crain, P.F., Trewyn, R.W., and Grosjean, H. 1996. Mechanism, specificity and general properties of the yeast enzyme catalysing the formation of inosine 34 in the anticodon of transfer RNA. *J. Mol. Biol.* **262**: 437–458.
- Bakin, A. and Ofengand, J. 1993. Four newly located pseudouridylate residues in *Escherichia coli* 23S ribosomal RNA are all at the peptidyltransferase center—Analysis by the application of a new sequencing technique. *Biochemistry* **32**: 9754–9762.
- Becker, H.F., Motorin, Y., Planta, R.J., and Grosjean, H. 1997. The yeast gene YNL292w encodes a pseudouridine synthase (Pus4) catalysing the formation of Ψ ₅₅ in both mitochondrial and cytoplasmic tRNAs. *Nucl. Acids Res.* **25**: 4493–4499.
- Behm-Ansmant, I., Urban, A., Ma, X., Yu, Y.T., Motorin, Y., and Branlant, C. 2003. The *Saccharomyces cerevisiae* U2 snRNA:pseudouridine-synthase Pus7p is a novel multisite-multisubstrate RNA: Ψ -synthase also acting on tRNAs. *RNA* **9**: 1371–1382.
- Behm-Ansmant, I., Grosjean, H., Massenet, S., Motorin, Y., and Branlant, C. 2004. Pseudouridylation at position 32 of mitochondrial and cytoplasmic tRNAs requires two distinct enzymes in *Saccharomyces cerevisiae*. *J. Biol. Chem.* **279**: 52998–53006.
- Behm-Ansmant, I., Massenet, S., Immel, F., Patton, J.R., Motorin, Y., and Branlant, C. 2006. A previously unidentified activity of yeast and mouse RNA:pseudouridine synthases 1 (Pus1p) on tRNAs. *RNA* **12**: 1583–1593.
- Chen, J. and Patton, J.R. 1999. Cloning and characterization of a mammalian pseudouridine synthase. *RNA* **5**: 409–419.
- Chen, J. and Patton, J.R. 2000. Pseudouridine synthase 3 from mouse modifies the anticodon loop of tRNA. *Biochemistry* **39**: 12723–12730.
- Conrad, J., Sun, D., Englund, N., and Ofengand, J. 1998. The *rhuC* gene of *Escherichia coli* codes for a pseudouridine synthase that is solely responsible for synthesis of pseudouridine at positions 955, 2504, and 2580 in 23 S ribosomal RNA. *J. Biol. Chem.* **273**: 18562–18566.
- Del Campo, M., Ofengand, J., and Malhotra, A. 2004. Crystal structure of the catalytic domain of RluD, the only rRNA pseudouridine synthase required for normal growth of *Escherichia coli*. *RNA* **10**: 231–239.
- Gietz, D., St. Jean, A., Woods, R.A., and Schiestl, R.H. 1992. Improved method for high efficiency transformation of intact yeast cells. *Nucleic Acids Res.* **20**: 1425.
- Grosshans, H., Lecointe, F., Grosjean, H., Hurt, E., and Simos, G. 2001. Pus1p-dependent tRNA pseudouridinylation becomes essential when tRNA biogenesis is compromised in yeast. *J. Biol. Chem.* **276**: 46333–46339.
- Huang, L., Pookanjanatavip, M., and Santi, D.V. 1997. Catalytic mechanism of RNA pseudouridine synthase. *FASEB J.* **11**: A1320.
- Jensen, R.E. and Johnson, A.E. 2001. Opening the door to mitochondrial protein import. *Nat. Struct. Biol.* **8**: 1008–1010.
- Kammen, H.O., Marvel, C.C., Hardy, L., and Penhoet, E.E. 1988. Purification, structure, and properties of *Escherichia coli* tRNA pseudouridine synthase I. *J. Biol. Chem.* **263**: 2255–2263.
- Kaya, Y. and Ofengand, J. 2003. A novel unanticipated type of pseudouridine synthase with homologs in bacteria, archaea, and eukarya. *RNA* **9**: 711–721.
- Kiss, A.M., Jady, B.E., Bertrand, E., and Kiss, T. 2004. Human box H/ACA pseudouridylation guide RNA machinery. *Mol. Cell. Biol.* **24**: 5797–5807.
- Koonin, E.V. 1996. Pseudouridine synthases: four families of enzymes containing a putative uridine-binding motif also conserved in dUTPases and dCTP deaminases. *Nucleic Acids Res.* **24**: 2411–2415.
- Lafontaine, D.L.J., Bousquet-Antonelli, C., Henry, Y., Caizergues-Ferrer, M., and Tollervey, D. 1998. The box H+ACA snoRNAs carry Cbf5p, the putative rRNA pseudouridine synthase. *Genes & Dev.* **12**: 527–537.
- Lecointe, F., Simos, G., Sauer, A., Hurt, E.C., Motorin, Y., and Grosjean, H. 1998. Characterization of yeast protein Deg1 as Pseudouridine synthase (Pus 3) catalysing the formation of Ψ ₃₈ and Ψ ₃₉ in tRNA anticodon loop. *J. Biol. Chem.* **273**: 1316–1323.
- Leehey, M.A., Squassoni, C.A., Friederich, M.W., Mills, J.B., and Hagerman, P.J. 1995. A noncanonical tertiary conformation of a human mitochondrial transfer RNA. *Biochemistry* **34**: 16235–16239.
- Ma, X., Zhao, X., and Yu, Y.T. 2003. Pseudouridylation (Ψ) of U2 snRNA in *S. cerevisiae* is catalyzed by an RNA-independent mechanism. *EMBO J.* **22**: 1889–1897.
- Ma, X., Yang, C., Alexandrov, A., Grayhack, E.J., Behm-Ansmant, I., and Yu, Y.T. 2005. Pseudouridylation of yeast U2 snRNA is catalyzed by either an RNA-guided or RNA-independent mechanism. *EMBO J.* **24**: 2403–2413.
- Martin, N.C. and Hopper, A.K. 1994. How single genes provide tRNA processing enzymes to mitochondria, nuclei, and the cytosol. *Biochimie* **76**: 1161–1167.
- Mason, T.L. 1998. Functional aspects of the three modified nucleotides in yeast mitochondrial large-subunit rRNA. In *The modification and editing of RNA* (eds. H. Grosjean and R. Benne). pp. 273–280. ASM Press, Washington, DC.
- Massenet, S., Motorin, Y., Lafontaine, D.L.J., Hurt, E.C., Grosjean, H., and Branlant, C. 1999. Pseudouridine mapping in the *Saccharomyces cerevisiae* spliceosomal UsnRNAs reveals that pseudouridine synthase Pus1p exhibits a dual substrate specificity for U2 snRNA and tRNA. *Mol. Cell. Biol.* **19**: 2142–2154.
- Motorin, Y., Keith, G., Simon, C., Foiret, D., Simos, G., Hurt, E., and Grosjean, H. 1998. The yeast tRNA:pseudouridine synthase Pus1p displays a multisite substrate specificity. *RNA* **4**: 856–869.
- Mumberg, D., Muller, R., and Funk, M. 1994. Regulatable promoters of *Saccharomyces cerevisiae*: Comparison of transcriptional activity and their use for heterologous expression. *Nucl. Acids Res.* **22**: 5767–5768.
- Patton, J.R., Bykhovskaya, Y., Mengesha, E., Bertolotto, C., and Fischel-Ghodsian, N. 2005. Mitochondrial myopathy and sideroblastic anemia (MLASA): Missense mutation in the pseudouridine synthase 1 (PUS1) gene is associated with the loss of tRNA pseudouridylation. *J. Biol. Chem.* **280**: 19823–19828.
- Reinhardt, A. and Hubbard, T. 1998. Using neural networks for prediction of the subcellular location of proteins. *Nucleic Acids Res.* **26**: 2230–2236.
- Rose, A.M., Belford, H.G., Shen, W.C., Greer, C.L., Hopper, A.K., and Martin, N.C. 1995. Location of N²,N²-dimethylguanosine-specific tRNA methyltransferase. *Biochimie* **77**: 45–53.
- Simos, G., Tekotte, H., Grosjean, H., Segref, A., Sharma, K., Tollervey, D., and Hurt, E.C. 1996. Nuclear pore proteins are involved in the biogenesis of functional tRNA. *EMBO J.* **15**: 2270–2284.
- Tekotte, H. and Davis, I. 2002. Intracellular mRNA localization: Motors move messages. *Trends Genet.* **18**: 636–642.
- Wesolowski, M. and Fukuhara, H. 1979. The genetic map of transfer RNA genes of yeast mitochondria: Correction and extension. *Mol. Gen. Genet.* **170**: 261–275.

RNA

A PUBLICATION OF THE RNA SOCIETY

The *Saccharomyces cerevisiae* Pus2 protein encoded by YGL063w ORF is a mitochondrial tRNA: Ψ 27/28-synthase

Isabelle Behm-Ansmant, Christiane Branlant and Yuri Motorin

RNA 2007 13: 1641-1647 originally published online August 7, 2007

Access the most recent version at doi:[10.1261/ma.605607](https://doi.org/10.1261/ma.605607)

References

This article cites 35 articles, 17 of which can be accessed free at:
<http://rnajournal.cshlp.org/content/13/10/1641.full.html#ref-list-1>

License

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).

To subscribe to RNA go to:
<http://rnajournal.cshlp.org/subscriptions>