Human DNMT2 methylates tRNA^{Asp} molecules using a DNA methyltransferase-like catalytic mechanism

TOMASZ P. JURKOWSKI,¹ MADELEINE MEUSBURGER,² SAMEER PHALKE,³ MARK HELM,⁴ WOLFGANG NELLEN,⁵ GUNTER REUTER,³ and ALBERT JELTSCH¹

¹Biochemistry Laboratory, School of Engineering and Science, Jacobs University Bremen, 28759 Bremen, Germany

²Department of Epigenetics, German Cancer Research Center, 69120 Heidelberg, Germany

³Institute of Genetics, Martin Luther University Halle-Wittenberg, 06120 Halle (Saale), Germany

⁴Institute of Pharmacy and Molecular Biotechnology, Department of Chemistry, Ruprecht-Karls Universität Heidelberg, 69120 Heidelberg, Germany

⁵Abteilung für Genetik, CINSaT, Universität Kassel, 34132 Kassel, Germany

ABSTRACT

Although their amino acid sequences and structure closely resemble DNA methyltransferases, Dnmt2 proteins were recently shown by Goll and colleagues to function as RNA methyltransferases transferring a methyl group to the C5 position of C38 in tRNA^{Asp}. We observe that human DNMT2 methylates tRNA isolated from Dnmt2 knock-out *Drosophila melanogaster* and *Dictyostelium discoideum*. RNA extracted from wild type *D. melanogaster* was methylated to a lower degree, but in the case of *Dictyostelium*, there was no difference in the methylation of RNA isolated from wild-type and Dnmt2 knock-out strains. Methylation of in vitro transcribed tRNA^{Asp} confirms it to be a target of DNMT2. Using site directed mutagenesis, we show here that the enzyme has a DNA methyltransferase-like mechanism, because similar residues from motifs IV, VI, and VIII are involved in catalysis as identified in DNA methyltransferases. In addition, exchange of C292, which is located in a CFT motif conserved among Dnmt2 proteins, strongly reduced the catalytic activity of DNMT2. Dnmt2 represents the first example of an RNA methyltransferase using a DNA methyltransferase type of mechanism.

Keywords: Dnmt2; catalytic mechanism; RNA methylation; tRNA^{Asp}

INTRODUCTION

Dnmt2 was initially assigned a member of the DNA methyltransferase family on the basis of its extensive homology with eukaryotic and prokaryotic DNA-(cytosine C5)-methyltransferases (Yoder and Bestor 1998). However, in the apparent absence of a phenotype in *dnmt2* knock-out cells, Dnmt2's possible biological function remained unknown (Okano et al. 1998), even though Dnmt2 is strongly conserved and it is found in species ranging from *Schizosaccharomyces pombe* to human. Later very weak, residual DNA methylation activity was found with enzymes from different species (Hermann et al. 2003; Kunert et al. 2003; Tang et al. 2003; Fisher et al. 2004; Kuhlmann et al. 2005). The finding that Dnmt2 is an active RNA methyl-

transferase capable of methylating the C38 position of the tRNA^{Asp} came as a surprise (Goll et al. 2006). However, still no cellular function of the tRNA^{Asp} methylation has been found, although in Zebrafish Dnmt2 knock-down caused a developmental phenotype (Rai et al. 2007). It is very intriguing that an enzyme that looks like a DNA methyltransferase can methylate RNA, in particular since the RNA and DNA specific m⁵C methyltransferases use different catalytic residues and a different mechanism for the methyl transfer reaction (Liu and Santi 2000).

Although, the cofactor S-adenosyl-L-methionine (AdoMet) is a very effective donor of methyl groups, methylation of cytosines at position 5 is not a trivial reaction, because cytosine is an electron-poor heterocyclic aromatic ring system and the carbon 5 of cytosine is not capable of making a nucleophilic attack on the methyl group of AdoMet. Therefore, the reactions catalyzed by RNA and DNA m⁵C methyltransferases follow the reaction pathway of a Michael addition. The catalytic mechanism of DNA m⁵C methyltransferases was first suggested by Santi (Santi et al. 1984) and later refined by Wu and Santi (1987).

Reprint requests to: Albert Jeltsch, Biochemistry Laboratory, School of Engineering and Science, Jacobs University Bremen, Campus Ring 1, 28759 Bremen, Germany; e-mail: a.jeltsch@jacobs-university.de; fax: +49 421 200 3249.

Article published online ahead of print. Article and publication date are at <http://www.rnajournal.org/cgi/doi/10.1261/rna.970408>.

According to this mechanism (for review, see Jeltsch 2002), DNA methylation is initiated by a nucleophilic attack of an SH group from a catalytic cysteine residue located in the conserved amino acids sequence motif IV (GPPC) (Kumar et al. 1994) on the C6 position of the target cytosine, yielding a covalent intermediate between the base and the enzyme. Thereby, the C5 position of the cytosine gets activated and becomes capable of performing a nucleophilic attack on the methyl group bound to the cofactor substrate AdoMet. The enzyme facilitates the nucleolytic attack on the C6 atom by a transient protonation of the cytosine ring at the endocyclic nitrogen atom N3 (Chen et al. 1991), which is stabilized by the glutamate residue from a highly conserved motif VI (ENV). The covalent complex between the methylated base and the DNA is resolved by deprotonation at the C5 position, which leads to the

elimination of the cysteinyl group and the reestablishment of aromaticity. Then, the methylated base together with the cofactor product, *S*-adenosyl-*L*-homocysteine, is released. In addition to the residues already mentioned, the second arginine residue in motif VIII (RXR) plays an important role in the catalytic mechanism of DNA ^{m5}C methyltransferases (O'Gara et al. 1996; Gowher et al. 2006).

The fact that RNA and DNA ^{m5}C methyltransferases employ different mechanisms for the methyl transfer was discovered by Liu and Santi (2000) by showing that RNA ^{m5}C methyltransferases do not use the cysteine from motif IV for the initial attack on the base, but rather one located in motif VI (TCS in RNA MTases). Later, King and Redman (2002) provided evidence that also the cysteine in motif IV has a role in catalysis. Moreover, instead of using the glutamate residue located in motif VI of DNA

FIGURE 1. Alignment of Dnmt2 family. (A) Multiple sequence alignment of Dnmt2 proteins from human, mouse, *D. melanogaster*, *E. histolytica*, *D. discoideum*, and *S. pombe*. The regions corresponding to catalytic motifs of DNA MTases are labeled. The residues selected for analysis are marked with arrows and labeled. (B) Structure of human DNMT2 protein (Dong et al. 2001). The residues studied in this work are shown in space fill representation. Note that the loop containing C79A is not ordered in the structure, such that this residue is not visible. (C) Schematic picture of the catalytic mechanisms proposed for DNA MTases and RNA MTases. For details see the text (data adapted from Bujnicki et al. 2004).

FIGURE 2. Protein purification and circular dichroism spectra of the DNMT2 wild type and all variants. (A) Coomassie stained SDS polyacrylamide gel showing the purified wild-type and mutant DNMT2 proteins. (B) Far UV circular dichroism spectra of DNMT2 and its variants recorded using 10 μ M enzyme in 10 mM Tris/HCl (pH 7.5), 200 mM KCl solution. The figure shows a superposition of the spectra measured with the wild-type and mutant proteins.

methyltransferases (ENV), it uses an aspartate residue from motif IV (DAPC) to stabilize the transition state of the reaction (Bujnicki et al. 2004).

Dnmt2 proteins contain all the conserved residues that are used for catalysis by DNA m^5C methyltransferases. However, they also contain a cysteine residue (C292 in human DNMT2-CFT), which is strongly conserved in the Dnmt2 family but absent in the DNA m^5C methyltransferase family. Based on the structure of the human DNMT2 enzyme (Dong et al. 2001), this residue is located near the putative catalytic pocket, such that it could be involved in catalysis. It was the aim of this study to determine if DNMT2 methylates RNA following a DNA or RNA MTase like catalytic mechanism. We conclude from our data that DNMT2 is the first example of an RNA MTase that uses the catalytic mechanism of DNA MTases.

RESULTS AND DISCUSSION

In order to identify the principal catalytic mechanism of the human DNMT2 enzyme for the transfer of methyl groups to RNA, we have created and purified the alanine exchange mutants of residues possibly involved in catalysis. To differentiate the catalytic mechanism of DNMT2 between the m^5C RNA and m^5C DNA types, we have selected nine residues for this mutagenesis study. The C79 (from motif

IV-PCQ) and C292 residues (conserved in the Dnmt2 family and close to the putative active site) were chosen to be studied by mutagenesis. Additionally, we have decided to exchange all other the cysteine residues in the protein for alanines to investigate their hypothetical involvement in catalysis. Moreover, we have selected E119 (from motif VI-ENV), R160 and R162 (from motif VIII-RXR) by their homology with the catalytic residues found in DNA m^5C methyltransferases that were experimentally confirmed in different systems (see Fig. 1; Klimasauskas et al. 1994; O'Gara et al. 1999; Sankpal and Rao 2002; Reither et al. 2003; Gowher et al. 2006; Shieh et al. 2006; Shieh and Reich 2007). We have purified these variants, confirmed their correct folding, and studied their RNA methylation activity and RNA binding.

Cloning, site-directed mutagenesis, protein expression, and purification

Using site-directed mutagenesis, all nine amino acid exchanges were performed and the mutant genes sequenced to confirm introduction of the desired mutation and the lack of additional ones. DNMT2 mutant proteins carrying a C-terminal His₆-Tag along with the wild-type enzyme were expressed in *Escherichia coli* and purified in soluble form to 100–500 μ M concentrations. The purity of DNMT2 wild-type and mutant proteins was >95% as determined by SDS-PAGE electrophoresis and Coomassie staining (Fig. 2).

Since single amino acid exchanges can disrupt the proper folding of a protein, we have determined the far UV circular dichroism (CD) spectra (which reflect the secondary structure composition of the protein) of all the DNMT2 protein mutants and compared them with the spectra obtained with the wild-type protein. As shown in Figure

FIGURE 3. RNA binding by DNMT2 and its variants analyzed by the nitrocellulose filter binding assay. (A) Dot blot analysis of the wild-type DNMT2 binding the ^{32}P labeled in vitro transcribed tRNA^{Asp}. (B) Binding curves of DNMT2 wild type and C79A and E119A mutants to in vitro transcribed and ^{32}P -labeled tRNA^{Asp}. Diamonds denote E119A mutant, triangles wild-type DNMT2, circles C79A. The experimental data points were fitted to a bimolecular binding equilibrium to determine the K_{Ass} values.

TABLE 1. Equilibrium binding constants and catalytic activities of the human DNMT2 and its variants

Enzyme variant	$K_{\text{Ass}} \text{ (M}^{-1}\text{)}$	Relative activity
DNMT2	6.1×10^5	+++
C24A	5.2×10^5	++
C79A	2.4×10^5	—
E119A	2.9×10^5	—
C140A	1.1×10^5	+
R160A	1.0×10^5	—
R162A	6.2×10^5	—
C287A	4.3×10^5	++
C292A	2.7×10^5	(+)

RNA binding was analyzed using in vitro transcribed tRNA^{Asp}. RNA methylation was investigated using total RNA extract prepared from Dnmt2 KO *D. melanogaster* flies. Activity ranges were classified as following: + + +, full activity or less than twofold reduction in activity; + +, 2- to 10-fold reduced activity; +, 10- to 50-fold reduced activity; (+), 50- to 250-fold reduced activity; —, no activity detectable (>5000-fold reduced activity).

2, all the single exchange variants' CD spectra were superimposable with the spectrum of the wild-type protein. This result indicates that all the mutant proteins were properly folded. All spectra were fitted by 32% α -helix and 17% β -strand, which is in good agreement with the secondary structure composition calculated from the crystallographic structure of human DNMT2 (35% α -helix, 15% β -strands).

RNA binding of wild-type Dnmt2 and its variants

We have investigated the RNA binding affinity of the wild-type DNMT2 protein as well as of all the mutants using a nitrocellulose filter binding assay. ³²P-labeled in vitro transcribed tRNA^{Asp} was used as a substrate, since tRNA^{Asp} was identified as a Dnmt2 target (Goll et al. 2006). In these experiments, constant amounts of labeled in vitro transcribed tRNA (0.4 nM) were incubated with increasing amounts of DNMT2 and sucked through a nitrocellulose filter (Fig. 3). The apparent equilibrium binding constant of the wild-type DNMT2 was $6 \times 10^5 \text{ M}^{-1}$ (Table 1). This value is low, when compared with nucleic acid interaction of other DNA or RNA modifying enzymes (for example, the K_{Ass} for M.EcoRV binding to 40 bp DNA substrate is $1.2 \times 10^6 \text{ M}^{-1}$ [Beck et al. 2001], and the K_{Ass} for Trm4p binding to tRNA^{Phe} is $2.2 \times 10^7 \text{ M}^{-1}$ [Walbott et al. 2007]). The weak tRNA binding of DNMT2 may be explained by the fact that the substrate used in the experiment was an in vitro transcribed tRNA^{Asp}, which lacks additional modifications like the mannosylqueosine base (manQ) present at the wobble position (34) of endogenous tRNA^{Asp} (Kuchino et al. 1981; Johnson et al. 1985). Most of the variants had equilibrium RNA binding constants similar to the wild type (Table 1). The weakest binding was observed with C79A and C292A, which both showed

about threefold reductions in binding affinity. The E119A mutant bound stronger to the tRNA, as shown in Figure 3B, which could be explained, because the removal of negative charge from the protein could reduce the electrostatic repulsion of the RNA. We conclude that all variants bind to the RNA to a similar degree as the wild-type DNMT2 does.

RNA methylation activity of the wild-type DNMT2

To detect the tRNA^{Asp} methylation activity of human DNMT2 protein reported by Goll et al. (2006), we have constructed a *Drosophila melanogaster* Dnmt2 null allele strain (Fig. 4). Total RNA was extracted from *D. melanogaster* wild-type flies and the Dnmt2 knock-out strain and used as substrate for in vitro methylation experiments using the purified enzymes. In these experiments the incorporation of tritiated methyl groups from *S*-[methyl-³H]-AdoMet into the RNA was detected. After the methylation reactions, the RNA was ethanol precipitated, dissolved in formamide, heat denatured, and subjected to gel electrophoresis on denaturing polyacrylamide gels. As shown in Figure 5A, the wild-type DNMT2 enzyme methylated RNA molecules extracted from Dnmt2 knock-out cells, which migrated in the gel in the size range of the tRNA fraction. In contrast, the RNA extracted from *D. melanogaster* cells that contained the active *dDnmt2* gene was methylated to a much lower extent (about 0.1% in Fig. 5B), which confirmed the specificity of the assay for DNMT2. The residual methylation observed with RNA isolated from wild-type flies suggests that either the tRNA substrates of Dnmt2 are not fully methylated in the wild-type *D. melanogaster* or

FIGURE 4. Generation of a dDnmt2 knock-out *Drosophila melanogaster* strain. (A) Schematic drawing of the structure of the dDnmt2¹⁴⁹ null allele. A null allele of dDnmt2 was generated after remobilization of the P element GE15695 inserted 128 bp upstream of the dDnmt2 gene. dDnmt2¹⁴⁹ shows an insertion of 59 bp of 5' P element sequences 9 bp downstream of first ATG position within the Dnmt2 ORF introducing an early stop codon. (B) RT-PCR detection of dDnmt2 mRNA. No specific transcript is detected in dDnmt2¹⁴⁹ null embryos. Lamine mRNA was used as internal control.

FIGURE 5. Methylation of total RNA extracted from the *dnmt2* knock-out *D. melanogaster* cells by wild-type DNMT2 and its variants. (A) Human DNMT2 methylates RNA molecules in the size range of tRNA. A DNA molecular weight marker was used and the approximate positions of the DNA fragments of different lengths (in nucleotides) are indicated. (B) Quantification of DNMT2 catalyzed methyl group transfer to RNA. In this experiment, DNA methylated by M.SssI was used to calibrate the RNA methylation by DNMT2. Different amounts of methylated DNA were loaded into the different lanes of the gel. The amount of methyl groups in each lane (in picomoles) is indicated above the corresponding well. In this experiment, 10 μ g of total RNA were loaded into each well.

that the DNMT2 shows relaxed specificity in vitro and methylates targets that are not methylated in vivo.

In order to estimate the amount of methyl groups incorporated into the tRNA, we used a 60mer DNA fragment methylated with M.SssI using the same radiolabeled AdoMet as methyl group donor and loaded in different amounts on the denaturing polyacrylamide gel together with the tritiated RNA samples (Fig. 5B). Quantification of the radioactivity observed in the DNA and in 10 μ g of DNMT2 modified RNA revealed that about 1 pmol of methyl groups was transferred to the RNA. Although this is only a rough estimate, the result fits to the estimated amount of tRNA^{Asp} used in the experiment, if one assumes that tRNA represents about 10% of the total RNA extract and tRNA^{Asp} is about 1/40 of all tRNAs. To measure the

rate of RNA methylation by DNMT2, time courses of the incorporation of radioactivity into the RNA were determined. The data were fitted to a single exponential reaction progress curve, revealing an apparent rate of RNA methylation of 4.4 (± 0.2) h⁻¹ under our experimental conditions (Fig. 6A).

Since tRNA^{Asp} was identified as Dnmt2 target, we used in vitro transcribed tRNA^{Asp} for the methylation assays as well. As shown in Figure 7A, in vitro transcribed tRNA^{Asp} was methylated by DNMT2, albeit at reduced efficiency, as indicated by the fact that the methylation signal was similar to that observed with the whole RNA extract, although 0.6 μ g of the in vitro transcribed tRNA were used, which is about 10 times more than the amount of tRNA^{Asp} present in 20 μ g of total RNA extract. Therefore, either native tRNA^{Asp} is the preferred substrate of DNMT2 or DNMT2 methylated other targets in the whole RNA preparation as well. Preferential methylation of native tRNAs could be due to the presence of additional modifications. Additionally, methylation of in vitro transcribed tRNA^{Asp} was only observed after Mg²⁺ ions were added to the buffer, presumably because this supports native folding of the tRNA. This detail in the experimental procedures, or the lower activity of DNMT2 on in vitro transcribed tRNA^{Asp}, might explain why this activity was not observed before (Goll et al. 2006).

We have also tested whether the human DNMT2 can methylate total RNA isolated from other organisms. To this end, RNA was extracted from *Dictyostelium discoideum* Ax2 wild-type strain and from *dnmA* knock-out cells (Kuhlmann et al. 2005) (DnmA is the homolog of Dnmt2 in *D. discoideum*). As shown in Figure 7B, the RNA isolated from both wild-type and *dnmA* knock-out cells was methylated by the human DNMT2 enzyme in vitro. We conclude that human, mouse, and *Dictyostelium* tRNA all are substrates for human DNMT2. Our results suggest, however, that tRNA methylation in vivo appears to be less complete or even absent in *Dictyostelium*; whether this depends on growth conditions and/or cell cycle remains an open question.

Methylation activity of the DNMT2 variants

In order to investigate the catalytic mechanism of DNMT2, we have performed in vitro methylation reactions of the total RNA extracts from Dnmt2 deficient *D. melanogaster* flies with all the purified DNMT2 mutant enzymes. Four of the variants (namely, C24A, C140A, C287A, and C292A) methylated the RNA to a detectable amount (Fig. 6B). For more detailed analysis, time courses of incorporation of radioactivity were determined, and the level of RNA methylation by mutants compared with the wild-type enzyme (Table 1). Our data show that C292 is important for catalytic activity. However, this residue cannot be directly involved in the catalytic mechanism, because the C292A

FIGURE 6. In vitro methylation of RNA by DNMT2 and its variants. (A) Time course of RNA methylation by wild-type DNMT2. The upper panel shows the time course of incorporation of radioactivity into RNA. The radioactive bands were analyzed quantitatively and the data fit to a single exponential reaction progress curve as shown in the lower panel. (B) Example of RNA methylation by the DNMT2 variants after 3 h reaction time.

variant showed significant residual activity. The C79A, E119A, R160A, and R162A mutants showed no detectable in vitro methylation activity. These inactive mutants did not show any detectable methylation activity even when the concentration of the protein in the methylation reaction was fivefold increased (data not shown). These experiments were done using one preparation of RNA and reproduced with two independent batches. As described above, the lack of activity cannot be explained by loss of RNA binding or misfolding of the proteins. We conclude that C79, E119, R160, and R162 are essential for the catalytic mechanism of DNMT2. These residues correspond to the residues that have most important roles in the catalytic mechanism of DNA MTases, suggesting that DNMT2 methylates RNA with a DNA MTase-like mechanism.

In conclusion, we have shown here that DNMT2 methylates RNA with a mechanism that follows DNA MTases and differs from mechanisms previously established for RNA MTases. This result is in agreement with the high similarity of DNMT2 to DNA MTases, with respect to both primary sequence and structure (Dong et al. 2001). Therefore, DNMT2 is the RNA MTase most similar to DNA MTases, which raises interesting questions of the evolutionary origin of both enzyme families and their substrate interaction.

MATERIALS AND METHODS

Site-directed mutagenesis, protein expression, and protein purification

Human DNMT2 wild type and its C79A variant (both in fusion with a C-terminal His₆-tag) were cloned into the pET28a(+) vector at HindIII-XhoI sites (Hermann et al. 2003).

The variants of human DNMT2 were created by the megaprimer site-directed mutagenesis method as described previously (Jeltsch and Lanio 2002). All created mutants were sequenced to check for absence of undesired additional mutations. The oligonucleotides used for site-directed mutagenesis were purchased from Thermo Hybaid. The human DNMT2-His₆ fusion proteins were expressed in *E. coli* (DE3) *Rosetta2 pLysS* cells. The cells were induced at OD(600 nm) = 0.6 with 1 mM IPTG and harvested 3 h after induction. The protein purification was performed as described previously (Hermann et al. 2003).

Generation of a *D. melanogaster* Dnmt2 null allele

The Dnmt2¹⁴⁹ null allele was generated by remobilization of the P element GE15695 inserted 128 bp upstream of the Dnmt2 ORF using the P[ry⁺ Δ2-3] transposase producing element. Homozygote GE15695 females were mated with SM6, *al² Cy sp²/+*; *Dr P[ry⁺ Δ2-3]/+* males. The resulting GE15695/SM6, *al² Cy sp²/Sco* females and w⁻ *Cy* exceptional progenies selected and analyzed for lesions in the Dnmt2 gene. Dnmt2¹⁴⁹ shows an insertion of 59 bp of 5' P element sequences 9 bp downstream of first ATG position within the Dnmt2 ORF introducing an early stop codon.

For RT-PCR analysis cDNA was prepared from the total RNA isolated from wild-type and Dnmt2¹⁴⁹ embryos. cMT2_{fwd} (TTG GTCGACTCATGCCTTTAATTGTGAG) and cMT2_{rev} (ATG TCGACGTTTTATCGTCAGCAATTTAATA) primers amplifying the complete Dnmt2 ORF were used. Lamine cDNA amplified with the primers Lamine_{fwd} (GAGACCACCAACATCAAGAAC)

FIGURE 7. In vitro methylation of different RNA substrates by DNMT2. (A) In vitro transcribed tRNA^{Asp} can be methylated by DNMT2 wild-type protein. A DNA molecular weight marker was used and the approximate positions of the DNA fragments of different lengths (in nucleotides) are indicated. (B) Methylation of the *D. discoideum* total RNA extract from the Ax2 wild-type cells and the *dnmtA* knock-out cells.

and Lamine_rev (AGGGACTGGATTTTCATCGC) was used as internal control.

RNA substrates

Overnight collections of *D. melanogaster* wild-type and mutant embryos were dechorionated in 50% bleach and flash frozen in liquid nitrogen. Total RNA was prepared using Trizol (Invitrogen) following the manufacturer's instructions. Pellets were dissolved in DEPC treated water.

The DNA template encoding for tRNA^{Asp}, the hammerhead ribozyme and the T7 promoter (TGGCTCCCCGTCGGGGAATC GAACCCCGGTCTCCCGCGTGACAGGCGGGGATACTACCA CTATACTAACGAGGAGACGGTACCGGGTACCGTTTCGTCC TCACGGACTCATCAGTCCTCGTTATCTCCCTATAGTGAGTC GTATT) was amplified in a standard PCR reaction using T7 primer and tRNA^{Asp} primer (TGGCTCCCCGTCGGGGAATCG). For in vitro transcription, 100 μ L of the PCR reaction were incubated with 200 μ L 2 $\times$ transcription buffer (80 mM Tris-HCl at pH 8.1, 2 mM Spermidine, 10 mM DTT, 0.02% Triton-X-100, 60 mM MgCl₂, 4 μ g/mL BSA), 5 mM of each NTP (final concentration), and 10 μ L of T7-Polymerase (Fermentas) in a total volume of 400 μ L for 3 h at 37°C. Transcripts were purified over 12% denaturing PAGE and bands of correct size were excised, eluted in 0.5 M ammonium acetate, and precipitated with two volumes of 100% ethanol. After centrifugation, pellets were washed once with 80% ethanol and then dissolved in water. Radioactively labeled tRNA^{Asp} was prepared by phosphorylating the in vitro transcribed tRNA^{Asp} with polynucleotide kinase (NEB) and γ -³²P-ATP (Amersham) according to the NEB protocol.

In vitro RNA methylation assay

The methylation of the total RNA extracts with the DNMT2 wild-type and its variants was performed similarly as previously described (Goll et al. 2006). Briefly, 20 μ g of *D. melanogaster* total RNA extracts were incubated with 1 μ M of the DNMT2 protein for 3 h in 40 μ L of methylation buffer (100 mM Tris/HCl at pH 7.5, 5% glycerol, 5 mM MgCl₂, 1 mM DTT, and 100 mM NaCl) containing 4.2 μ M labeled [methyl-³H] AdoMet (NEN). After the reaction, samples were phenol extracted, ethanol precipitated, resuspended in formamide, and run on a 7 M urea 12% denaturing polyacrylamide gel. Afterward, the gels were stained with ethidium bromide, fixed with 10% acetic acid, 10% methanol solution, immersed for 1 h in Amplify solution (Amersham), dried, and exposed to Hyperfilm film (Amersham) for 1 d to 1 wk at -70°C. All RNA methylation assays were conducted at least in triplicate. Quantification was done by densitometric analysis of the films. Data were fitted to a single exponential reaction progress curve using the following equation: $\text{intensity}(t) = BL + A \times [1 - \exp(-t \times k)]$, where *BL* is the background signal, *A* is the scaling factor relating densitometric staining and methylation, and *k* is the apparent turnover rate constant of RNA methylation.

Circular dichroism

Far UV circular dichroism spectra were obtained with a Jasco J-815 circular dichroism spectrophotometer, using protein concentrations of 10 μ M DNMT2 in 10 mM Tris/HCl (pH 7.5), 200 mM

KCl solution. The spectra were recorded in a cell with a 0.1-mm path length in the wavelength window between 190 nm and 250 nm with a step size of 0.1 nm and bandwidth of 1 nm. For all spectra, 25 scans were performed and averaged. The buffer baseline was recorded and consequently subtracted from the protein spectra. All experiments were carried out at least twice.

Nitrocellulose filter binding assay with in vitro transcribed tRNA^{Asp}

The tRNA^{Asp} binding of the DNMT2 wild type and the mutants was analyzed by nitrocellulose filter binding assay experiments using the in vitro transcribed and ³²P labeled tRNA^{Asp}. The experiment was carried out using a Bio-Dot apparatus (BioRad). The reaction mixtures contained 0–5 μ M DNMT2 protein in 100 mM Tris/HCl (pH 7.5), 5 mM MgCl₂, 100 mM NaCl, 1 mM DTT, 5 mg/mL BSA, 2 units RNasin ribonuclease inhibitor (Promega), 10 μ M sinefungin, and 0.4 nM in vitro transcribed ³²P labeled tRNA^{Asp}. The reaction mixtures were first preincubated at ambient temperature for 20 min to allow the system to reach equilibrium, then loaded on a prewashed nitrocellulose membrane and immediately sucked through the membrane. The membrane was washed three times with the wash buffer (100 mM Tris/HCl at pH 7.5, 5 mM MgCl₂, 100 mM NaCl, 1 mM DTT) and dried, and the bound radioactivity was quantified using a phosphorimager (FLA-300, Fuji). The data were fitted to a bimolecular binding equilibrium model using the Excel Solver module. All binding experiments were reproduced at least twice.

ACKNOWLEDGMENTS

Thanks are due to Vladimir Maximov for providing total RNA extracts from *Dictyostelium* and Béatrice Gollinelli for providing modified RNA samples. This work has been supported partially by DFG (JE 252/5).

Received December 19, 2007; accepted April 17, 2008.

REFERENCES

- Beck, C., Cranz, S., Solmaz, M., Roth, M., and Jeltsch, A. 2001. How does a DNA interacting enzyme change its specificity during molecular evolution? A site-directed mutagenesis study at the DNA binding site of the DNA-(adenine-N6)-methyltransferase EcoRV. *Biochemistry* **40**: 10956–10965.
- Bujnicki, J.M., Feder, M., Ayres, C.L., and Redman, K.L. 2004. Sequence-structure-function studies of tRNA:m5C methyltransferase Trm4p and its relationship to DNA:m5C and RNA:m5U methyltransferases. *Nucleic Acids Res.* **32**: 2453–2463.
- Chen, L., MacMillan, A.M., Chang, W., Ezaz-Nikpay, K., Lane, W.S., and Verdine, G.L. 1991. Direct identification of the active-site nucleophile in a DNA (cytosine-5)-methyltransferase. *Biochemistry* **30**: 11018–11025.
- Dong, A., Yoder, J.A., Zhang, X., Zhou, L., Bestor, T.H., and Cheng, X. 2001. Structure of human DNMT2, an enigmatic DNA methyltransferase homolog that displays denaturant-resistant binding to DNA. *Nucleic Acids Res.* **29**: 439–448.
- Fisher, O., Siman-Tov, R., and Ankri, S. 2004. Characterization of cytosine methylated regions and 5-cytosine DNA methyltransferase (Ehmeth) in the protozoan parasite *Entamoeba histolytica*. *Nucleic Acids Res.* **32**: 287–297.

- Goll, M.G., Kirpekar, F., Maggert, K.A., Yoder, J.A., Hsieh, C.L., Zhang, X., Golic, K.G., Jacobsen, S.E., and Bestor, T.H. 2006. Methylation of tRNA^{Asp} by the DNA methyltransferase homolog Dnmt2. *Science* **311**: 395–398.
- Gowher, H., Loutchanwoot, P., Vorobjeva, O., Handa, V., Jurkowska, R.Z., Jurkowski, T.P., and Jeltsch, A. 2006. Mutational analysis of the catalytic domain of the murine Dnmt3a DNA-(cytosine C5)-methyltransferase. *J. Mol. Biol.* **357**: 928–941.
- Hermann, A., Schmitt, S., and Jeltsch, A. 2003. The human Dnmt2 has residual DNA-(cytosine-C5) methyltransferase activity. *J. Biol. Chem.* **278**: 31717–31721.
- Jeltsch, A. 2002. Beyond Watson and Crick: DNA methylation and molecular enzymology of DNA methyltransferases. *ChemBioChem* **3**: 274–293.
- Jeltsch, A. and Lanio, T. 2002. Site-directed mutagenesis by polymerase chain reaction. *Methods Mol. Biol.* **182**: 85–94.
- Johnson, G.D., Pirtle, I.L., and Pirtle, R.M. 1985. The nucleotide sequence of tyrosine tRNA^Q ψ A from bovine liver. *Arch. Biochem. Biophys.* **236**: 448–453.
- King, M.Y. and Redman, K.L. 2002. RNA methyltransferases utilize two cysteine residues in the formation of 5-methylcytosine. *Biochemistry* **41**: 11218–11225.
- Klimasauskas, S., Kumar, S., Roberts, R.J., and Cheng, X. 1994. HhaI methyltransferase flips its target base out of the DNA helix. *Cell* **76**: 357–369.
- Kuchino, Y., Shindo-Okada, N., Ando, N., Watanabe, S., and Nishimura, S. 1981. Nucleotide sequences of two aspartic acid tRNAs from rat liver and rat ascites hepatoma. *J. Biol. Chem.* **256**: 9059–9062.
- Kuhlmann, M., Borisova, B.E., Kaller, M., Larsson, P., Stach, D., Na, J., Eichinger, L., Lyko, F., Ambros, V., Soderbom, F., et al. 2005. Silencing of retrotransposons in *Dictyostelium* by DNA methylation and RNAi. *Nucleic Acids Res.* **33**: 6405–6417.
- Kumar, S., Cheng, X., Klimasauskas, S., Mi, S., Posfai, J., Roberts, R.J., and Wilson, G.G. 1994. The DNA (cytosine-5) methyltransferases. *Nucleic Acids Res.* **22**: 1–10.
- Kunert, N., Marhold, J., Stanke, J., Stach, D., and Lyko, F. 2003. A Dnmt2-like protein mediates DNA methylation in *Drosophila*. *Development* **130**: 5083–5090.
- Liu, Y. and Santi, D.V. 2000. m5C RNA and m5C DNA methyltransferases use different cysteine residues as catalysts. *Proc. Natl. Acad. Sci.* **97**: 8263–8265.
- O’Gara, M., Klimasauskas, S., Roberts, R.J., and Cheng, X. 1996. Enzymatic C5-cytosine methylation of DNA: Mechanistic implications of new crystal structures for HhaI methyltransferase-DNA-AdoHcy complexes. *J. Mol. Biol.* **261**: 634–645.
- O’Gara, M., Zhang, X., Roberts, R.J., and Cheng, X. 1999. Structure of a binary complex of HhaI methyltransferase with S-adenosyl-L-methionine formed in the presence of a short non-specific DNA oligonucleotide. *J. Mol. Biol.* **287**: 201–209.
- Okano, M., Xie, S., and Li, E. 1998. Dnmt2 is not required for de novo and maintenance methylation of viral DNA in embryonic stem cells. *Nucleic Acids Res.* **26**: 2536–2540.
- Rai, K., Chidester, S., Zavala, C.V., Manos, E.J., James, S.R., Karpf, A.R., Jones, D.A., and Cairns, B.R. 2007. Dnmt2 functions in the cytoplasm to promote liver, brain, and retina development in zebrafish. *Genes & Dev.* **21**: 261–266.
- Reither, S., Li, F., Gowher, H., and Jeltsch, A. 2003. Catalytic mechanism of DNA-(cytosine-C5)-methyltransferases revisited: Covalent intermediate formation is not essential for methyl group transfer by the murine Dnmt3a enzyme. *J. Mol. Biol.* **329**: 675–684.
- Sankpal, U.T. and Rao, D.N. 2002. Mutational analysis of conserved residues in HhaI DNA methyltransferase. *Nucleic Acids Res.* **30**: 2628–2638.
- Santi, D.V., Norment, A., and Garrett, C.E. 1984. Covalent bond formation between a DNA-cytosine methyltransferase and DNA containing 5-azacytosine. *Proc. Natl. Acad. Sci.* **81**: 6993–6997.
- Shieh, F.K. and Reich, N.O. 2007. AdoMet-dependent methyl-transfer: Glu119 is essential for DNA C5-cytosine methyltransferase M.HhaI. *J. Mol. Biol.* **373**: 1157–1168.
- Shieh, F.K., Youngblood, B., and Reich, N.O. 2006. The role of Arg165 towards base flipping, base stabilization and catalysis in M.HhaI. *J. Mol. Biol.* **362**: 516–527.
- Tang, L.Y., Reddy, M.N., Rasheva, V., Lee, T.L., Lin, M.J., Hung, M.S., and Shen, C.K. 2003. The eukaryotic DNMT2 genes encode a new class of cytosine-5 DNA methyltransferases. *J. Biol. Chem.* **278**: 33613–33616.
- Walbott, H., Husson, C., Auxilien, S., and Golinelli-Pimpaneau, B. 2007. Cysteine of sequence motif VI is essential for nucleophilic catalysis by yeast tRNA m5C methyltransferase. *RNA* **13**: 967–973.
- Wu, J.C. and Santi, D.V. 1987. Kinetic and catalytic mechanism of HhaI methyltransferase. *J. Biol. Chem.* **262**: 4778–4786.
- Yoder, J.A. and Bestor, T.H. 1998. A candidate mammalian DNA methyltransferase related to pmt1p of fission yeast. *Hum. Mol. Genet.* **7**: 279–284.

RNA

A PUBLICATION OF THE RNA SOCIETY

Human DNMT2 methylates tRNA^{Asp} molecules using a DNA methyltransferase-like catalytic mechanism

Tomasz P. Jurkowski, Madeleine Meusburger, Sameer Phalke, et al.

RNA 2008 14: 1663-1670

References

This article cites 31 articles, 10 of which can be accessed free at:
<http://rnajournal.cshlp.org/content/14/8/1663.full.html#ref-list-1>

License

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).

To subscribe to *RNA* go to:
<http://rnajournal.cshlp.org/subscriptions>