The archaeal COG1901/DUF358 SPOUT-methyltransferase members, together with pseudouridine synthase Pus10, catalyze the formation of 1-methylpseudouridine at position 54 of tRNA

KUNAL CHATTERJEE,^{1,5} IAN K. BLABY,^{2,5,6} PATRICK C. THIIVILLE,² MRINMOYEE MAJUMDER,¹ HENRI GROSJEAN,³ Y. ADAM YUAN,⁴ RAMESH GUPTA,^{1,7} and VALÉRIE DE CRÉCY-LAGARD^{2,7}

¹Department of Biochemistry and Molecular Biology, Southern Illinois University, Carbondale, Illinois 62901-4413, USA

²Department of Microbiology & Cell Science, University of Florida, Gainesville, Florida 32611-0700, USA

³Université Paris11, IGM, CNRS, UMR 8621, Orsay, F 91405, France

⁴Department of Biological Sciences and Temasek Life Sciences Laboratory, National University of Singapore, Singapore, 117543

ABSTRACT

The methylation of pseudouridine (Ψ) at position 54 of tRNA, producing $m^1\Psi$, is a hallmark of many archaeal species, but the specific methylase involved in the formation of this modification had yet to be characterized. A comparative genomics analysis had previously identified COG1901 (DUF358), part of the SPOUT superfamily, as a candidate for this missing methylase family. To test this prediction, the COG1901 encoding gene, *HVO_1989*, was deleted from the *Haloferax volcanii* genome. Analyses of modified base contents indicated that while $m^1\Psi$ was present in tRNA extracted from the wild-type strain, it was absent from tRNA extracted from the mutant strain. Expression of the gene encoding COG1901 from *Halobacterium* sp. NRC-1, VNG1980C, complemented the $m^1\Psi$ minus phenotype of the ΔHVO_1989 strain. This in vivo validation was extended with in vitro tests. Using the COG1901 recombinant enzyme from *Methanocaldococcus jannaschii* (Mj1640), purified enzyme Pus10 from *M. jannaschii* and full-size tRNA transcripts or T Ψ -arm (17-mer) fragments as substrates, the sequential pathway of $m^1\Psi$ 54 formation in Archaea was reconstituted. The methylation reaction is AdoMet dependent. The efficiency of the methylase reaction depended on the identity of the residue at position 55 of the T Ψ -loop. The presence of Ψ 55 allowed the efficient conversion of Ψ 54 to $m^1\Psi$ 54, whereas in the presence of C55, the reaction was rather inefficient and no methylation reaction occurred if a purine was present at this position. These results led to renaming the Archaeal COG1901 members as TrmY proteins.

Keywords: tRNA modification; tRNA methylation; *Haloferax volcanii*; TrmY; Archaea

INTRODUCTION

Transfer RNAs in all three domains of life contain a large variety of characteristic post-transcriptionally modified nucleosides. Among them pseudouridine, an isomer of uridine (abbreviated Ψ) and base or ribose methylations (mN, or Nm respectively; N being any of the four bases A, U, C, or G) are by far the most abundant (all listed in Czerwonic

et al. 2009; Cantara et al. 2011) (for review, see Grosjean 2009; Motorin and Helm 2011). Many of these noncanonical nucleotides are highly conserved at specific sites in functionally important parts of tRNA molecules. For example Ψ , m^5U (riboT), or $m^1\Psi$ are nearly always found at position 54 in the so-called T Ψ -loop at the elbow of the L-shaped tRNA molecules, while Ψ is found at the neighboring position 55. The conservation of these modifications at these two positions is certainly due to essential structural roles. Indeed, the presence of Ψ 55 reinforces tertiary base-pairing with the conserved G18, favors intra-loop stacking with the conserved purine at position 57 and with the neighboring m^5U 54/ $m^1\Psi$ 54 engaged in a reverse-Hoogsteen pair with the conserved A58 (Romby et al. 1987). Together with the conserved tertiary pair C56-G19 between the T Ψ -loop and D-loop, this “ensemble” of conserved interacting

⁵These authors contributed equally to this work.

⁶Present address: Department of Chemistry and Biochemistry, University of California, Los Angeles, CA 90095-1569, USA.

⁷Corresponding authors.

E-mail rgupta@siumed.edu.

E-mail vcrcy@ufl.edu.

Article published online ahead of print. Article and publication date are at <http://www.rnajournal.org/cgi/doi/10.1261/rna.030841.111>.

nucleotides allows crucial interactions that form a stable “tertiary core” and, hence, the canonical tRNA L-shape architecture (Kotlova et al. 2007 and references therein). In agreement with this “locking” function of both Ψ 55 and m^5U 54 (probably also $m^1\Psi$ 54 in archaeal tRNAs, see below), these base modifications are among the earliest to appear during the complex tRNA maturation process (Nishikura and De Robertis 1981; Grosjean et al. 1996).

Even if the modifications at position 54 and 55 are conserved and have critical structural roles, strains lacking these modifications are viable. A mutant of *Escherichia coli* deleted in the *truB* gene (encoding for the Ψ 55 producing enzyme) grew normally on all media tested (Gutgsell et al. 2000). It did exhibit a competitive disadvantage in extended co-culture with its wild-type progenitor and a defect in surviving rapid transfers from 37°C to 50°C (Gutgsell et al. 2000). Moreover, combining *truB* mutations with mutations affecting the catalytic activity of TrmA, the enzyme catalyz-

ing the formation of the adjacent m^5U 54, further increased the temperature sensitivity phenotype (Kinghorn et al. 2002).

In the present work, we focus on the biosynthesis of 1-methylpseudouridine in tRNAs ($m^1\Psi$ 54) (Fig. 1A). This Ψ derivative was first characterized from the bulk tRNA of archaeon *Halococcus morrhuae* (Pang et al. 1982). It has now been found in tRNAs isolated from many Archaea, mainly Euryarchaeota (Table 1). Its location at position 54 of tRNA was inferred from sequence analysis of 41 tRNAs from *H. volcanii* (Gupta 1984, 1986). In contrast, tRNA sequences from *Thermoplasma acidophilum* (two sequences), *Methanobacterium thermoautotrophicum* (one sequence), and *Sulfolobus acidocaldarius* (one sequence) revealed the presence of a nonmethylated Ψ or a ribose-methylated U (Um) in place of $m^1\Psi$ 54 (Cantara et al. 2011; <http://www.uni-bayreuth.de/departments/biochemie/trna/>). In tRNAs from Thermococcales (*Pyrococcus furiosus* and *Pyrococcus abyssi*), m^5U 54 or s^2m^5U 54 are found instead of $m^1\Psi$ 54 (Fig. 1A;

FIGURE 1. Enzymatic post-transcriptional modifications of selected uridines in RNA. (A) Schematic consensus of tRNA secondary structures indicating the U54 target within the highly conserved 7-nt TΨ-loop. The dashed line indicates a reverse Hoogsteen base pair within the loop. In the majority of Archaea, U54 is first isomerized into Ψ54 by a tRNA pseudouridine synthase, aPus10 (symbol “a” preceding the acronym of the enzyme denotes “archaeal,” “b” means “bacterial,” and “e” means “eukaryal”). In certain Archaea, Ψ54 can be further methylated into $m^1\Psi$ 54 by a SPOUT-type, SAM-dependent methyltransferase designated TrmY (this work). In Thermococcales, Bacteria, and Eukarya, U54 is methylated into m^5U 54 (riboT) by a distinct SAM-dependent tRNA-U54 methyltransferase aTrmU54, bTrmA, and eTrm2p, respectively. In thermophilic organisms, m^5U 54 can be further thiolated into s^2m^5U (s^2T) by the heteromeric enzyme TtuA/TtuB. (B) Schematic consensus of a portion of 16S rRNA as part of domain IV, encompassing the highly conserved 8-nt helix 35 Ψ-loop. U914 in 16S rRNA of *M. jannaschii*, corresponding to U1191 in 18S rRNA of *S. cerevisiae*, is first isomerized into Ψ by an archaeal enzyme (or enzymatic system) that remains to be identified. In *S. cerevisiae* this reaction is mediated by the snoRNP complex consisting of the pseudouridine synthase eCbf5 and snR35 guide RNA. The methylation of N1-atom of the uracil ring is further catalyzed by a SPOUT-type and SAM-dependent methyltransferase, Nep1 (also referred as Emg1). Only in some Eukarya (such as yeast, *Drosophila*, HeLa cells), is the $m^1\Psi$ further hypermodified into an $acp^3m^1\Psi$ derivative. On the right part of the figure are the different uridine derivatives. The dashed lines through the structure of $m^1\Psi$ and $acp^3m^1\Psi$ show the axis of base rotation during the isomerization process. The asterisks with small arrows indicate the atoms normally engaged in the reverse Hoogsteen base pair with A58.

TABLE 1. Identification of m¹Ψ or s²T (s²m⁵U) in archaeal tRNAs

KINGDOM Phylum	Modified base	<i>pus10</i>	<i>trmY</i>	<i>trmU54</i>	Reference
<i>Organism</i>					
EURYARCHAEOTA					
Methanosarcinales					
<i>Methanococcoides burtonii</i>	m ¹ Ψ	Yes	Yes	No	1
<i>Methanosarcina barkerii</i>	m ¹ Ψ	Yes	Yes	No	2
Archeoglobales					
<i>Archaeoglobus fulgidus</i>	m ¹ Ψ	Yes	Yes	No	3
Thermoplasmatales					
<i>Thermoplasma acidophilum</i>	Ψ*	Yes	No	No	2, 4, 5
Methanococcales					
<i>Methanococcus vanniellii</i>	m ¹ Ψ	Yes	Yes	No	2, 6
<i>Methanococcus voltae</i>	m ¹ Ψ	Yes	Yes	No	2
<i>Methanocaldococcus jannaschii</i>	m ¹ Ψ	Yes	Yes	No	This work
<i>Methanococcus igneus</i>	m ¹ Ψ	nc	nc	nc	1
Methanobacteriales					
<i>Methanothermobacter fervidus</i>	m ¹ Ψ	nc	nc	nc	3
<i>Methanob. thermotrophicum</i>	Ψ*	Yes	No	No	7
<i>Methanobacterium bryantii</i>	no m ¹ Ψ	nc	nc	nc	2
<i>Methanobrevibacter smithii</i>	no m ¹ Ψ	Yes	No	No	2
<i>Methanomicrobium mobile</i>	m ¹ Ψ	nc	nc	nc	2
Halobacteriales					
<i>Haloferax volcanii</i>	m ¹ Ψ*	Yes	Yes	No	2, 8, 9
<i>Halococcus morrhuae</i>	m ¹ Ψ*	nc	nc	nc	2, 5, 10
<i>Halobacterium cutirubrum</i>	m ¹ Ψ*	nc	nc	nc	11, 12
Thermococcales					
<i>Pyrococcus abyssi</i>	s ² m ⁵ U	Yes	Yes	Yes	13
<i>Pyrococcus furiosus</i>	s ² m ⁵ U	Yes	Yes	Yes	14, 15
<i>Thermococcus-isola MB4</i>	s ² m ⁵ U	Yes	Yes	Yes	3
CRENARCHAEOTA					
Desulfurococcales					
<i>Pyrodictium occultum</i>	no m ¹ Ψ	No	No	No	3
<i>Stetteria hydrogenophila</i>	no m ¹ Ψ	nc	nc	nc	1
<i>Thermoplasma maritimum</i>	no m ¹ Ψ	nc	nc	nc	3
Sulfolobales					
<i>Sulfolobus acidocaldarius</i>	Um*	No ^{&}	No	No	2, 5
<i>Sulfolobus solfataricus</i>	no m ¹ Ψ	No ^{&}	No	No	3
<i>Acidans infernus</i>	no m ¹ Ψ	nc	nc	nc	3
Thermoproteales					
<i>Pyrobaculum islandicum</i>	m ¹ Ψ	Yes	Yes	No	3
<i>Thermoproteus neutrophilus</i>	m ¹ Ψ	Yes	Yes	No	3

Purified bulk tRNAs were analyzed for their nucleoside content after complete digestion by nuclease, usually nuclease P1, followed by dephosphorylation and LC/MS or 2D-TLC after 5'-³²P- post-labeling. An alternative TLC separation method used RNase T2-digests of uniformly labeled tRNAs isolated from ³²P-labeled cells. The original LC-MS method is described in detail in Edmonds et al. (1985) and Crain (1990). For the ³²P-post-labeling technique and TLC analysis, detailed information can be found in Grosjean et al. (2007). Analyses of uniformly labeled tRNAs can be found in Gupta and Woese (1980) and Gupta (1984). m¹Ψ, m⁵U, or s²T have been found to date only at position 54 in tRNA. Their presence in bulk tRNA hydrolysates are therefore diagnostic of their presence at position 54 exclusively. Presence of Ψ alone or Um, or any other U derivatives at position 54 can be deduced only from direct tRNA sequencing (indicated in the table by an asterisk [*] next to the symbol of the modified nucleoside). The indication "no m¹Ψ" does not exclude the possibility of a nonmethylated Ψ in the TΨ-loop. Beside Um found in initiator tRNA^{Met} sequenced from *S. acidocaldarius*, s²U and s²Um were also found in bulk tRNA in some Crenarchaeota (Edmonds et al. 1991). Since no sequence analyses are available, this information was omitted from the table even if these modifications could be found at position 54. The presence or absence of a gene coding for aPus10, aTrmY, and aTrmU54 was derived from the SEED database and from the taxonomic tool of Blink at NCBI. The italicized symbol Yes for TrmY of Thermococcales means that it is not certain whether the *trmY* gene is expressed and/or the gene product is functional, as no m¹Ψ was reported in tRNA of Thermococcales.

The "nc" symbol means "not applicable" because the genome sequence is not available. The "&" symbol refers to the fact that a gene coding for a Pus10 homolog is found in these genomes, but the corresponding proteins do not meet structural criteria for a functioning Pus10 (E Fitzek, R Gupta, and M Geisler, unpubl.). Names for *Halobacterium cutirubrum* have changed with time; *Halobacterium salinarum*, *Halobacterium halobium* are the same species. *Halobacterium* sp. NRC1, a different strain from *H. cutirubrum*, but with a sequenced genome was used to identify the presence of gene *pus10*, *trmY*, and *trmU54*. Numbers correspond to the following references: (1) (Noon et al. 2003); (2) (Gupta and Woese 1980), m¹Ψ is listed as xU in this reference; (3) (Edmonds et al. 1991); (4) (Kilpatrick and Walker 1981); (5) (Kuchino et al. 1982); (6) (Best 1978), m¹Ψ was misidentified as dihydrouridine in this reference, due to their similar mobilities in TLC; (7) (Gu et al. 1984); (8) (Gupta 1984); (9) (Gupta 1986); (10) (Pang et al. 1982); (11) (Gu et al. 1983); (12) (Nicoghossian et al. 1985); (13) (Urbonavicius et al. 2008); (14) (Constantinesco et al. 1999); (15) (Kowalak et al. 1994).

Kowalak et al. 1994; Constantinesco et al. 1999). As previously noted (Gupta and Woese 1980; Pang et al. 1982), methyl groups on the uracil ring at tRNA position 54 in the two cases (N1 in $m^1\Psi$ and C5 in m^5U) bear similar orientations with respect to the ribose and the polynucleotide chain and both methylated products have similar shapes. This could be an example of evolutionary convergence of structures (and probably of function).

Only the first step in $m^1\Psi$ formation in tRNA, the formation of $\Psi54$, has been elucidated with the recent characterization of the tRNA- $\Psi54$ synthase, Pus10 (Gurha and Gupta 2008). The enzyme responsible for the subsequent N1 methylation of the uracil ring of $\Psi54$ and the corresponding gene has yet to be identified. However, experiments with whole-cell extract of *H. volcanii* incubated with ^{32}P -radiolabeled T7-transcript and AdoMet (SAM) showed enzymatic formation of $m^1\Psi54$, indicating that the enzyme of interest was SAM dependent (Grosjean et al. 1995).

Nep1 (Nucleolar Essential Protein1, formerly named Emg1) from *S. cerevisiae* and *Methanocaldococcus jannaschii* was recently identified as a genuine SAM-dependent N1-pseudouridine methyltransferase (Wurm et al. 2010). This enzyme belongs to Cluster of Orthologous Group (COG) 1756 (Tatusov et al. 2003) and converts Ψ into $m^1\Psi$ in small synthetic fragments of 8, 9, or 11 nucleotides in length (GAUUCACGCC, where the second of the two adjacent Us is Ψ) (Wurm et al. 2010). This motif corresponds to the sequence in helix 35 of SSU rRNA of *S. cerevisiae* as well as of *M. jannaschii*, where Nep1 tightly binds (Buchhaupt et al. 2006). Nep1 is responsible for the methylation step in the formation of the hypermodified 1-methyl-3-(3-amino-3-carboxypropyl) pseudouridine ($m^1acp^3\Psi$) found at position 1191 in helix 35 of yeast 18S rRNA and also has a role in ribosome assembly (Fig. 1B; Meyer et al. 2011; Thomas et al. 2011). These functions appear to be conserved in other eukaryotes (Brand et al. 1978; Youvan and Hearst 1981). Despite the obvious sequence analogy of the Nep1 target with the conserved sequence GUUCAANC/U (underlined, cf. target sequences in Fig. 1, A and B) present in nearly all tRNAs sequenced so far (Marck and Grosjean 2002), Nep1 is certainly not the missing tRNA N1-pseudouridine methyltransferase. The reasons are: (1) The genome of *S. acidocaldarius* contains a Nep1 homolog (Saci_0034), but its bulk tRNA lacks $m^1\Psi$, and its only sequenced tRNA harbors U_{m54} (Table 1; Gupta and Woese 1980; Kuchino et al. 1982); (2) conversely, the genome of *H. volcanii* lacks the gene coding for Nep1, while nearly all of its tRNAs harbor $m^1\Psi54$ (Table 1; Gupta 1984, 1986); (3) this absence is consistent with the fact that helix 35 of *H. volcanii* 16S rRNA harbors an acp^3U and not the “hypermodified” $m^1acp^3\Psi$ as in eukaryotes (Kowalak et al. 2000).

A better candidate for the missing $m^1\Psi54$ methyltransferase came from a bioinformatics analysis of a large variety of orphan genes coding for putative AdoMet-dependent methyltransferases in genomes of microorganisms belong-

ing to the three domains of life. This analysis identified one of the methyltransferases belonging to COG1901, encompassing an α/β knot fold (also named SPOUT) superfamily of methyltransferases as a valid candidate (Tkaczuk et al. 2007). This prediction fits with the observation that genes of this family usually cluster with *pus10* in several archaeal genomes (Grosjean et al. 2008a). Lastly, the crystal structure of a COG1901 family member, Mj1640 from *M. jannaschii*, was solved in complex with AdoMet at 1.4 Å resolution (Chen and Yuan 2010). Mj1640 protein shares much structural similarity with the highly conserved eukaryotic nucleolar Emg1/Nep1 protein at its C-terminal half containing the conserved deep trefoil knot fold (Z-score 15.7, RMS 2.5 Å, 163 C α). However, because of significant differences in N-terminal extension domain and overall surface charge distribution, it was suggested that the two proteins Nep1 and Mj1640 target different RNA sequences (Chen and Yuan 2010). This structural data greatly strengthened the prediction that the COG1901 (or DUF358) RNA methylase family was the missing tRNA- $m^1\Psi$ forming enzyme, and we set out to test this experimentally using both in vivo and in vitro assays. Here we show that a *H. volcanii* strain deleted in the COG1901 family gene *HVO_1989* lacks $m^1\Psi$ in tRNA, and that in vitro Mj1640 catalyzes the formation of $m^1\Psi$ at position 54 on tRNA. These results led to the renaming of the Archaeal COG1901 proteins as TrmY (as Y is commonly used to represent Ψ) for tRNA(pseudouridine54-N1)-methyltransferase.

RESULTS

trmY is not essential in *H. volcanii*, but deletion of *trmY* leads to the absence of $m^1\Psi54$ in *H. volcanii* tRNA

The *H. volcanii trmY* (COG1901 homolog, *HVO_1989*) was deleted using previously published methods (Allers et al. 2004). Deletion of *trmY* in this strain (VDC2376) was confirmed by PCR using two primer pairs, one designed to anneal outside the deleted fragment and the other within (Fig. 2A). *trmY* is therefore dispensable in *H. volcanii*. Deletion of *trmY* did not lead to any obvious phenotype when analyzing growth in rich or minimal liquid medium or on solid medium at 20, 30, 37, 44, and 50°C or with total salt concentrations ranging from 12% (1.812 M NaCl, 65.28 mM MgCl₂, 63 mM MgSO₄, and 41.64 mM KCl) to 25% (3.775 M NaCl, 136 mM MgCl₂, 131.25 mM MgSO₄, and 86.75 mM KCl) (compared with the standard YPC media with a final concentration of 18% salt; data not shown). To test whether TrmY was involved in $m^1\Psi$ formation, we first extracted bulk tRNA from VDC2376 ($\Delta trmY$) and H26 (the isogenic parent strain as wild type [WT]). The bulk tRNA preparations were enzymatically hydrolyzed with nuclease P1 and snake venom phosphodiesterase and dephosphorylated with alkaline phosphatase,

FIGURE 2. Construction and phenotype of the *trmY*-deleted strain of *H. volcanii*. (A) Deletion of the *trmY* gene was confirmed by PCR; (left) PCR products using primers designed to anneal outside the gene (HvO_1989_Ext_F and HvO_1989_Ext_R) confirm a genomic rearrangement of correctly predicted sizes in WT and mutant strains; (middle) PCR products using primers designed to anneal within the target gene (HvO_1989_Int_F and HvO_1989_Int_R) show the absence of the *trmY* internal segment in the mutant and its presence in the WT strain; (right). To confirm the presence of *VNG_1980c* in *trans*, primers were designed to anneal to the complementing gene (Hsa1_COG1901_Fwd and Hsa1_COG1901_Rev). The predicted size of the fragment is observed in the rescued strain. (B) LC-MS/MS analysis of tRNA extracted from H26 (wild-type) and VDC2376 ($\Delta trmY$) showing the UV trace at 254nm (top). The 259 *m/z* ion that corresponds to the protonated molecular weight (MH^+) of $m^1\Psi$ was detected by MS at 13.36 min in the WT background, while no 259 *m/z* ion was detected in the $\Delta trmY$ strain (bottom).

and the ribonucleosides analyzed by LC/MS-MS as described previously (de Crécy-Lagard et al. 2010). The 259 *m/z* ion that corresponds to the protonated molecular weight (MH^+) of $m^1\Psi$ was detected by MS at 13.36 min in the WT background, while no 259 *m/z* ion was detected in the $\Delta trmY$ strain (Fig. 2B). The peak observed in the mutant strain and not in the WT strain between 14 and 14.5 min corresponds to background (see the differences in the intensity scales between the right and left panels). As in our set-up, the $m^1\Psi$ peak eluted as a trail of the U peak at 13–13.6 min and these two peaks were difficult to separate with a trailing peak most certainly corresponding to an unidentified modification remaining in the mutant strain,

a two-dimensional thin layer chromatography (2D-TLC) method was used to better detect the $m^1\Psi$ modification. Also, to rescue the effects of the *trmY* deletion, a derivative of the $\Delta trmY$ strain was created (VDC2604), which contained a plasmid-borne copy of the COG1901 encoding gene from *Halobacterium* sp. NRC1, *VNG_1980C* (Fig. 2A). The WT, $\Delta trmY$ (VDC2376), and $\Delta trmY$ +pHTTrmY (VDC2604) strains were in vivo labeled with ^{32}P . Total tRNA was isolated, digested by nuclease P1, and resolved by 2D-TLC. As shown in Figure 3A, radioactive $m^1\Psi$ spot is present in both WT and complemented strains but is absent from the $\Delta trmY$ strain (VDC2376). Because $m^1\Psi$ is present only at position 54 in *H. volcanii* tRNAs (Gupta 1984, 1986), these results show that, at least in halophilic Archaea, COG1901 proteins (TrmY) are required for the formation of $m^1\Psi$ 54 in tRNA.

Lack of TrmY in the $\Delta trmY$ strain only affects methylation and not Ψ formation at position 54 of the tRNA. We used CMCT-primer extension reactions, which specifically allow the identification of Ψ at a particular position in RNA (Bakin and Ofengand 1998) to determine the presence of Ψ 54 in $\Delta trmY$ strain. As seen in Figure 3B, the Ψ -specific band (marked by an arrow) is present at position 54 of tRNA in the $\Delta trmY$ strain, but not in the WT and complemented strains, as expected if $m^1\Psi$ 54 and not Ψ 54 is present in these last two strains. As expected, Ψ 55-specific bands are present in all three strains. These results suggested that Pus10-mediated Ψ 54 (and Ψ 55) formation does not require the presence of TrmY activity.

M. jannaschii TrmY catalyzes the formation of $m^1\Psi$ 54 in tRNA in vitro

The [α - ^{32}P]UTP-labeled tRNA^{Trp} substrate, generated by transcription of a plasmid carrying a synthetic *H. volcanii* tRNA^{Trp} lacking its intron, was first modified by purified recombinant *M. jannaschii* Pus10, thus converting U at both positions 54 and 55 to Ψ , as described before (Gurha and Gupta 2008). This in vitro pseudouridylated tRNA^{Trp} was used as a substrate for further enzymatic methylation by the recombinant *M. jannaschii* TrmY protein mostly

FIGURE 3. Formation of $m^1\Psi$ of *H. volcanii* tRNA is mediated by TrmY. (A) Nuclease P1 digests of uniformly labeled tRNAs were resolved by 2D-TLC. pA, pC, pG, pU, pΨ, and $pm^1\Psi$ indicate 5'-phosphorylated A, C, G, U, Ψ, and $m^1\Psi$, respectively. The radioactive spot corresponding to $pm^1\Psi$ is present in both wild-type (H26) and complemented ($\Delta trmY$ +pHTrmY) strains, but is absent from $\Delta trmY$ strain (middle). (B) CMCT-primer extension analyses to determine the modification status of residue at position 54 of *H. volcanii* elongator tRNA^{Met} were done using primer Met-CCA2 (position marked in Fig. 4A) and total small RNA of wild-type, $\Delta trmY$, and $\Delta trmY$ +pHTrmY strains. RNAs were treated with (+) or without (−) CMCT for the indicated time (in minutes), followed by alkali (OH⁻) treatment (+) or no treatment (−). Positions of tRNA residues 54 and 55 are marked on the side. A dark band in CMCT followed by alkali treatment lanes, with an increased intensity in the 20-min treatment lane, indicates the presence of Ψ at that position. These reactions show that unmethylated Ψs are present at position 55 in all three strains, but at position 54 only in the $\Delta trmY$ strain (the band is marked by an arrow).

prepared under renaturing conditions. The doubly modified RNA was then phenol extracted, digested with nuclease P1, and the resulting digested products analyzed by 2D-TLC. Autoradiographies of the TLC plates revealed the presence of a radiolabeled spot corresponding to $pm^1\Psi$ only when the tRNA^{Trp} transcript was first incubated with Pus10 and then with TrmY (Fig. 4B). When tRNA^{Trp} was

treated with TrmY without pretreatment with Pus10 it did not show any $pm^1\Psi$ (data not shown). Nearest neighbor analyses using RNase T2 digestion on the same radiolabeled tRNA samples confirmed the presence of $m^1\Psi$ only in the tRNA^{Trp} incubated with both Pus10 and TrmY (Fig. 4C), attesting that within tRNA^{Trp} the $m^1\Psi$ residue was located on the 5' side of a ³²P-labeled-uridine residue. RNase T2 produces ribonucleotide 3'-monophosphates (Np), and in the process transfers labeled phosphate on the 5' side of a residue in the RNA to the 3' side of the preceding residue. In *H. volcanii* tRNA^{Trp} there is only one U residue (position 54) that precedes another U residue (position 55) (see Fig. 4A). Therefore, the $m^1\Psi$ observed in Figure 4C is obviously derived from the residue at position 54 of this RNA. Moreover, Ψp is absent in the rightmost panel of Figure 4C, indicating that the methylation reaction was nearly complete under our experimental conditions, i.e., all Ψ54 being converted into $m^1\Psi$ 54. This methylation is AdoMet dependent, as it was not observed when AdoMet was omitted from the reaction mixture (data not shown). Similar methylations were obtained when *H. volcanii* elongator tRNA^{Met} was used as substrate (Fig. 4D) instead of *H. volcanii* tRNA^{Trp}. This tRNA^{Met} contains residues G58 and G60 instead of the conserved A58 and semiconserved pyrimidine-60 at these positions of the TΨ-loop (Fig. 4A).

Initial methylation reactions were done using proteins prepared under native conditions. However, additional AdoMet was not needed for methylation with this protein when the protein was in excess of the RNA. AdoMet had to be added to the reaction for complete conversion of Ψ54 to $m^1\Psi$ 54 when the RNA was equal to or in molar excess of

this protein. This was not unexpected, because the protein prepared under native conditions contains bound AdoMet (Chen and Yuan 2010). Furthermore, methylation by AdoMet-containing TrmY (under excess enzyme concentration) could be inhibited by addition of S-adenosyl-homocysteine, an inhibitor of SAM-dependent methyltransferases (data not shown). Similarity of results using

FIGURE 4. *M. jannaschii* TrmY can convert Ψ54 of tRNA to m¹Ψ54. (A) Sequences of transcripts used in the reactions. Mutations changing U55 of tRNA^{Trp} and U54 of tRNA^{Met} are indicated. The arrow in tRNA^{Met} sequence indicates the position of primer Met-CCA2 used for CMCT-primer extension analyses shown in Figure 3B. (B,C) [α-³²P]UTP-labeled tRNA^{Trp} was first pseudouridylated by *M. jannaschii* Pus10 (Mj-Pus10) and then methylated by *M. jannaschii* TrmY (Mj-TrmY) as described in Materials and Methods. Nuclease P1 or RNase T2 (indicated in panels) digests of purified products were resolved by 2D-TLC on cellulose plates. “p” before or after a nucleoside letter indicates the 5’ or 3’ phosphate of that nucleoside. (Pi) inorganic phosphate. (D) Treatments similar to those in B using [α-³²P]UTP-labeled tRNA^{Met} as substrate. The middle panel here (and in B) is shown to indicate that Ψ in the tRNA is produced by Mj-Pus10. (E) TLC separation of RNase T2 digest of [α-³²P]CTP-labeled tRNA^{Trp} following treatment with Mj-TrmY. (F–I) TLC separation of nuclease P1 digests of [α-³²P]UTP-labeled T-arm-Trp, mutant tRNA^{Trp}-U55A, tRNA^{Trp}-U55G, and tRNA^{Trp}-U55C, respectively (indicated in the panels), following treatment with Mj-TrmY.

proteins prepared under native and renaturing conditions suggested that AdoMet is not required for the stability of the recombinant protein.

M. jannaschii TrmY converts only tRNA Ψ 54 and not Ψ 55 to $m^1\Psi$. This is shown by TLC analyses of RNase T2 digestion of [α - 32 P]CTP-labeled tRNA^{Trp} first incubated with Pus10 then with TrmY (Fig. 4E). Autoradiography of the TLC plate reveals the presence of labeled Ψ p and the absence of $m^1\Psi$ p, indicating that Ψ 55 (preceding C56) is not converted to $m^1\Psi$ in this tRNA.

***M. jannaschii* TrmY can produce $m^1\Psi$ in just the T Ψ -arm (stem-loop) of tRNA**

Previously it was shown that *M. jannaschii* Pus10 can produce Ψ in the T-arm-Trp (Fig. 4A), a 17-base fragment of tRNA^{Trp}, at positions that correspond to positions 54 and 55 of tRNA (Gurha and Gupta 2008). The same pseudouridylated fragment and the equivalent fragment from tRNA^{Met} (Fig. 4A) are both substrates for the *M. jannaschii* TrmY (Fig. 4F; data not shown). Methylation of Ψ in Pus10-treated T-arm-Trp occurs only at positions equivalent to 54 and not 55 of the tRNA (data not shown).

***M. jannaschii* TrmY requires Ψ 55 (or possibly U55) in the tRNA to produce $m^1\Psi$ 54**

In a mutant tRNA^{Trp}, where U55 is changed to any other residue (C, A, or G), the U54 can still be converted to Ψ 54 by *M. jannaschii* Pus10 (Gurha and Gupta 2008). However, Ψ 54 of a mutant tRNA^{Trp}, where the U55 is changed to A55 (Fig. 4A), is not a substrate for *M. jannaschii* TrmY. Nuclease P1 digest of [α - 32 P]UTP-labeled transcript of this mutant tRNA does not show any $pm^1\Psi$ after treatment with TrmY (Fig. 4G). Similarly, a U-to-G change at position 55 did not lead to any $m^1\Psi$ formation (Fig. 4H), while a trace amount of $pm^1\Psi$ was observed when U55 was changed to C55 (Fig. 4I). These experiments suggest that the presence of Ψ 55 (or possibly U55) in the tRNA is required for enzymatic methylation of the adjacent Ψ 54 by TrmY to produce $m^1\Psi$ 54. However, at present we cannot distinguish whether an unmodified U55 can substitute for the requirement of Ψ 55, because Pus10 treatment of RNA converts both U54 and U55 in our substrates to Ψ (Gurha and Gupta 2008).

DISCUSSION

In the present work, by combining genetic and biochemical experiments, we showed that in at least two Euryarchaeota, *H. volcanii*, and *M. jannaschii* COG1901, genes encode the missing tRNA(pseudouridine54-N1)-methyltransferase or TrmY. This newly identified TrmY enzyme belongs to the SAM-dependent SPOUT super family of dimeric enzymes (Tkaczuk et al. 2007). The closest COG1901 homologs among SPOUT methyltransferase of known catalytic func-

tion is the newly identified dimeric Nep1 (COG1756) catalyzing the N1-methylation of Ψ ($m^1\Psi$) in helix 35 of 18S rRNA of Eukarya and Archaea (Fig. 1B; Wurm et al. 2010). Hence, these two families represent yet another example of two closely related SPOUT subfamilies that target different RNA molecules (an rRNA and a tRNA).

M. jannaschii TrmY can produce $m^1\Psi$ 54 in full-sized tRNAs as well as at the equivalent position in T Ψ -stem-loop substrates (17-mer T Ψ -arm of tRNA). Likewise, the tRNA: m^5 U54 methyltransferases of *E. coli* (TrmA or RumT), of *S. cerevisiae* (Trm2p), and of *P. abyssi* (PAB_0719, aTrmU54) can produce m^5 U54 in similar full-sized or truncated RNAs (Gu et al. 1996; Becker et al. 1997; Urbonavicius et al. 2008), suggesting a possibility of similar substrate recognition modes as described for *E. coli* TrmA (Alian et al. 2008) and archaeal TrmU54 (Walbott et al. 2008). However, archaeal TrmY can methylate Ψ 54 in tRNA harboring G58 and G60 (Fig. 4A,D). This differs from the yeast Trm2p enzyme, which requires the presence of both A58 and a pyrimidine at position 60 to methylate its substrate (Becker et al. 1997). This specificity of yeast Trm2p probably depends on the formation of a reverse intra-loop Hoogsteen pair between U54 and A58 in the tRNAs (Becker et al. 1997). The *M. jannaschii* TrmY requires a Ψ (or possibly a U) at position 55 to efficiently methylate Ψ 54. This is reminiscent of the requirements of the yeast Trm2p enzyme, where U55 cannot be replaced by any other residue (Becker et al. 1997). We could not determine whether the residue 3' adjacent to the target Ψ residue needs to be Ψ or an unmodified U because of our experimental design, but in the case of the archaeal Nep1, it was shown that the 5' adjacent residue to the target Ψ residue can be an unmodified U (Wurm et al. 2010).

In addition to *M. jannaschii* TrmY, high-resolution crystal structures of other SAM-dependent RNA methyltransferases, such as Nep1 of *M. jannaschii* and *S. cerevisiae*, TrmA and RumA both of *E. coli*, are known (Lee et al. 2005; Alian et al. 2008; Leulliot et al. 2008; Taylor et al. 2008; Thomas et al. 2011). Nep1 is an rRNA N1- Ψ methyltransferase, TrmA is a tRNA C5- m^5 U methyltransferase, whereas RumA is an rRNA C5- m^5 U methyltransferase. Structural analysis of these RNA methyltransferases reveals that TrmY and Nep1 belong to SPOUT family, whereas TrmA and RumA belong to Rossmann (α/β) fold. These two groups share fewer structural similarities, except for a somehow similar α/β fold for AdoMet and RNA substrate binding. Although TrmY and Nep1 share the same characteristic deep trefoil knot core, different structural domains/motifs extended from the deep trefoil knot core are observed. These extra domains/motifs are proposed to facilitate RNA substrate binding and/or selection. Similarly, TrmA and RumA share close structural similarity both in RNA binding and catalytic domains. However, significant structural differences of the extended loop linking the RNA-binding domain and the catalytic domain

are observed, suggesting the functional role of this loop in substrate selection.

TrmY homologs are found in the majority of Euryarchaeota and in a few Crenarchaeota. All Archaea that contain a TrmY homolog contain a Pus10 homolog (see the COG1901 Subsystem in the SEED database; <http://pubseed.theseed.org/SubsysEditor.cgi?page=SubsystemOverview>) consistent with our finding that the presence of Ψ at position 54 is a prerequisite to the formation of m¹Ψ. The presence or absence of m¹Ψ in archaeal tRNAs has been reported by different research groups over many years (Table 1). The presence of Ψ or m¹Ψ correlates well with the presence and absence of Pus10 and TrmY, with the exception of the Thermococcales (Table 1). Despite the presence of both *pus10* and *trmY* genes in their genomes (Table 1) (COG1901 Subsystem in the SEED database), m⁵U or s²T and not m¹Ψ were formally identified in bulk tRNAs from Thermococcales (Fig. 1A; Table 1). Likewise, when a tRNA transcript specific for Phe, Asp, or Ile, lacking modifications, was incubated with AdoMet and cell extracts from *P. furiosus* or *P. abyssi*, m⁵U was almost quantitatively formed at position 54 of the TΨ-loop (Constantinesco et al. 1999; Urbonavicius et al. 2008). Under the same experimental conditions, U54 in the tRNA^{Ile} transcript was shown to be “doubly” modified into m¹Ψ when incubated with cell extracts from *H. volcanii* (Grosjean et al. 1995). The enzyme responsible for the formation of m⁵U54 in *P. abyssi* was recently identified as PAB_0719 (aTrmU54) (Urbonavicius et al. 2008; Auxilien et al. 2011). This protein is a member of the COG2265 SAM-dependent, Rossmann-like RNA m⁵U methyltransferase family. Phylogenetic analysis showed that it has been acquired by lateral gene transfer of a bacterial RlmD gene responsible for the site-specific formation of m⁵U in 23S rRNA (position 1939 in *E. coli*). Thus, during evolution, the RlmD-like protein in *P. abyssi* (PAB_0719, aTrmU54) has changed target specificity from rRNA to tRNA. The question of the role of TrmY proteins in Thermococcales remains open. Are these still active as a tRNA m¹Ψ methyltransferase, but in specific conditions that have escaped detection, or have they acquired another function? Further experimental work is required to address this issue.

Only a few sequenced Crenarchaeota encode both genes *pus10* and *trmY* (COG1901 subsystem in the SEED database), these include *Ignococcus hospitalis* KIN4_1, *Aeropyrum pernix* K1, the *Pyrobaculum* species, and a few other (but not all) Thermoproteales. Indeed, m¹Ψ54 was detected in bulk tRNAs extracted from two thermoproteales (*Pyrobaculum islandicum* and *Thermoproteus neutrophilus*) fitting with the genomic data (Table 1). The only other experimental data available for a Crenarchaeota is for *S. acidocaldarius*, where 2'-O-methyluridine at tRNA position 54 (Um54) was identified (Gupta and Woese 1980; Kuchino et al. 1982), there again fitting with the absence of both active Pus10 proteins and TrmY in Sulfolobales (Table 1).

While COG1901 is primarily found in Archaea, homologs are surprisingly found in most sequenced *Vibrio* genomes, in two *Shewanella baltica* strains (OS195 and OS185), two *Photobacterium profundum* strains (SS9 and 3TCK) and in *Alcanivorax borkumensis* SK2 (Fig. 5A) (COG1901 subsystem in the SEED database). m¹Ψ54 is typically an archaeal tRNA signature modification (Gupta and Woese 1980; Pang et al. 1982) and has never been identified in any bacterial tRNA sequenced so far (for review, see Grosjean et al. 2008b)). Therefore, the presence of TrmY was not expected in bacterial genomes. In addition, the gene coding for the enzyme TrmA responsible for the formation of the canonical m⁵U54 in the T-loop of bacterial tRNA (Ny and Björk 1980) is found in all sequenced *Vibrio* genomes (VV1_1171 in *Vibrio vulnificus*, see COG1901 subsystem in the SEED database). The presence of COG1901 homologs in just a few bacteria could suggest that the gene family arose from a recent horizontal gene transfer event from Archaea. However, the phylogenetic analysis of the COG1901 family (Fig. 5A) shows that bacterial and archaeal COG1901 proteins form two distinct monophyletic groups, which is not easily compatible with the above hypothesis. Although the overall fold of VV2_1434 (COG1901 homolog in *V. vulnificus*) shares close structural similarity to that of Mj1640 (Z-score 26.0, RMS 1.6 Å, 181 Cα), significant differences for local conserved motifs and overall charge distributions are observed between these two structures (Fig. 5B). In general, VV2_1434 has a more compact structure compared with Mj1640, suggesting that the bacterial and archaeal homologs may fulfill different functions. For example, Mj1640 displays short β-strands with flexible loops at the middle part of the β-sheet compared with those of VV2_1434, whereas VV2_1434 displays an extra α-helix at its “head” of the overall “butterfly” shape (Fig. 5B). Further experimental work is required to establish the function of the bacterial COG1901 members.

This work generally reinforces the difficulty of transferring functional annotations of RNA methylases. TrmY are homologs of Nep1 but carry the same reaction (methylation of a Ψ residue) in different substrates, one tRNA the other rRNA. Also, even if we have functionally characterized the function of COG1901 as TrmY in a few Archaea, some of the COG1901 homologs (for example, in Thermococcales or Bacteria) might have another function. Experimental validation of judiciously chosen members is the only solution to correctly annotating RNA methylase superfamilies.

MATERIALS AND METHODS

Bioinformatics

Sequences and distribution of all genes/proteins analyzed in this work are available through the “COG1901 sub-system” on the public SEED server (<http://pubseed.theseed.org/SubsysEditor.cgi?page=SubsystemOverview>) (Overbeek et al. 2005). We also used

ers. The amplicon was inserted, after digestion with the compatible enzymes, between the *Nde*I and *Blp*I sites of pJAM202 to generate pIKB421 (pHTrmY). This plasmid was verified by PCR, restriction digestion, and sequencing, and then passed through *E. coli* (INV110, Invitrogen) before transformation into the *H. volcanii* strain VDC2376 (Δ trmY), creating complemented strain VDC2604. Transformants were selected for by plating onto HYP-PC containing novobiocin (0.3 μ g/mL).

Detection of m¹Ψ residues in tRNAs

For analysis by liquid chromatography–tandem mass spectrometry (LC–MS/MS), *H. volcanii* bulk tRNA was prepared, hydrolyzed, and analyzed as described previously (de Crécy-Lagard et al. 2010). tRNA purifications and analyses were performed at least twice independently. For analysis by 2D–TLC, uniformly ³²P-labeled total tRNA was prepared as described before (Gupta 1984; Joardar et al. 2008). Labeled tRNAs were digested with nuclease P1 and digests were resolved on cellulose plates (EM Science) using isobutyric acid/0.5 N NH₄OH (5:3, v/v) in the first dimension and isopropanol/HCl/H₂O (70:15:15, v/v/v) in the second dimension (Gupta 1984). Radioactivity in the plates was revealed by phosphorimaging. Presence of Ψ at a specific position in tRNA was analyzed by the 1-cyclohexyl-3-(2-morpholinoethyl) carbodiimide metho-*p*-toluenesulfonate (CMCT) modification technique (Ofengand et al. 2001; Motorin et al. 2007) using total small RNA. For this, initially total RNA was prepared from the cells using TRI reagent (Molecular Research Center, Inc). Small RNA fraction from the total RNA was extracted by suspending total RNA in 1 M NaCl by vigorous vortexing. The high-salt soluble fraction was separated from insoluble material by centrifugation. RNA was precipitated from the supernatant by ethanol. A total of 5 μ g of this RNA was treated with CMCT for 2, 10, and 20 min each at 37°C, while an untreated sample was simply incubated at 37°C for 20 min as control. After precipitation of the RNA, alkali treatment was done for 3 h at 37°C for all samples except the ones treated with CMCT for 2 min. Again, after precipitation the RNA was used for primer extension with [5′-³²P]-labeled primers that hybridized close to and on the 3′ side of the position to be mapped for Ψ. AMV reverse transcriptase (Promega) was used for primer extension according to the manufacturer's protocol, except that the reverse-transcription reaction was done at 50°C for 1 h with 0.75 mM dNTP each. The extension stops at one residue before the CMCT modified Ψ. A dark band in CMCT, followed by alkali treatment lanes, with an increased intensity in the 20-min treatment lane indicates the presence of Ψ at that position. The position of the band in the gel that corresponds to Ψ is determined by its distance from the end of the primer and sometimes the presence of darker bands for U and G in the 2-min CMCT, but no alkali treatment lane, and correlation with the known sequence of the RNA.

Expression and purification of TrmY of *M. jannaschii*

Recombinant TrmY protein of *M. jannaschii* (MJ1640) was purified as described before (Chen and Yuan 2010). These procedures are considered native conditions in this work. The protein was also prepared under renaturation conditions. For this, harvested cells were resuspended in a buffer (10 mM Na₂HPO₄, 2 mM KH₂PO₄, 2.7 mM KCl, 500 mM NaCl) containing 8 M urea, and lysed by sonication. The supernatant from a 25-min spin at 13,000 rpm at 4°C was loaded onto a Ni²⁺ column at room temperature equilibrated with

the same buffer. The protein was renatured on the column in a gradient of 8 M urea to no urea in the same buffer containing 2.5 mM imidazole. After a brief wash with 25 mM imidazole containing buffer, the protein was eluted using 250 mM imidazole in the same buffer. Pooled fractions were then dialyzed using 500 mM NaCl, 10 mM DTT, 50 mM Tris–Cl (pH 8.0), and 20% Glycerol.

In vitro tRNA methylation assays

Transcripts for *H. volcanii* tRNA^{Trp} and elongator tRNA^{Met} (lacking their introns), T-arm-Trp, and T-arm-Met were generated as described previously (Gurha et al. 2007). Mutants of tRNA^{Trp} were prepared by independently changing U55 to A, C, and G, and of tRNA^{Met} by changing U54 to A as described before (Gurha et al. 2007). Sequences of these transcripts are shown in Figure 4A. Appropriate labeled transcripts were prepared using relevant [α -³²P]NTP. Recombinant *M. jannaschii* Pus10 was used first to convert U54 and U55 equivalents of these transcripts into Ψ as described previously (Gurha and Gupta 2008). *M. jannaschii* TrmY efficiently works under a range of salt concentrations (100–500 mM NaCl). Our standard 20- μ L methylation reaction contained 100–150 nM pseudouridylated RNA and 1 μ M renatured *M. jannaschii* TrmY in 20 mM Tris–Cl (pH 7.0), 150 mM NaCl, 0.75 mM DTT, 1.5 mM MgCl₂, 0.1 mM EDTA, 0.1 mM AdoMet, and 5% Glycerol. The reaction mixtures were incubated at 68°C for 1 h, RNA was purified by phenol/chloroform extraction and ethanol precipitation, and digested with RNase T2 or nuclease P1. The digests were resolved by TLC as described before for uniformly labeled tRNA. For methylation inhibition reactions, proteins isolated under native conditions were used and 0.1 mM of S-adenosylhomocysteine, instead of AdoMet, was used in the reactions.

SUPPLEMENTAL MATERIAL

Supplemental material is available for this article.

ACKNOWLEDGMENTS

H. volcanii H26 was received from Julie Maupin-Furlow (University of Florida) as well as from Thorsten Allers (University of Nottingham, UK). *Halobacterium* sp. NRC-1 genomic DNA was prepared by Gabriela Phillips (University of Florida). Plasmid pJAM202 was a gift from Julie Maupin-Furlow. We thank Archi Joardar for providing *M. jannaschii* Pus10 and Sophie Alvarez for performing the LC–MS analyses, and Elisha Fitzek for analysis of archaeal Pus10 using structural criteria for functional proteins. H.G. holds a position of Emeritus Scientist at the CNRS, Centre de Génétique Moléculaire, in Gif-sur-Yvette. This work was funded in part by NIGMS (Grant no. 1RC2GM092602–01 sub-contract to V.d.C.-L.), NSF (Grant no. MCB-05169448 to V.d.C.-L.), and NIH (Grant no. GM055945 to R.G.).

Received October 10, 2011; accepted December 2, 2011.

REFERENCES

Alian A, Lee TT, Griner SL, Stroud RM, Finer-Moore J. 2008. Structure of a TrmA–RNA complex: A consensus RNA fold contributes to

- substrate selectivity and catalysis in m⁵U methyltransferases. *Proc Natl Acad Sci* **105**: 6876–6881.
- Allers T, Ngo H, Mevarech M, Lloyd RG. 2004. Development of additional selectable markers for the halophilic archaeon *Haloferax volcanii* based on the *leuB* and *trpA* genes. *Appl Environ Microbiol* **70**: 943–953.
- Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* **25**: 3389–3402.
- Auxilien S, Rasmussen A, Rose S, Brochier-Armanet C, Husson C, Fourmy D, Grosjean H, Douthwaite S. 2011. Specificity shifts in the rRNA and tRNA nucleotide targets of archaeal and bacterial m⁵U methyltransferases. *RNA* **17**: 45–53.
- Bakin AV, Ofengand J. 1998. Mapping of pseudouridine residues in RNA to nucleotide resolution. *Methods Mol Biol* **77**: 297–309.
- Becker HF, Motorin Y, Sissler M, Florentz C, Grosjean H. 1997. Major identity determinants for enzymatic formation of ribothymidine and pseudouridine in the T ψ -loop of yeast tRNAs. *J Mol Biol* **274**: 505–518.
- Best AN. 1978. Composition and characterization of tRNA from *Methanococcus vannielii*. *J Bacteriol* **133**: 240–250.
- Blaby IK, Phillips G, Blaby-Haas CE, Gulig KS, El Yacoubi B, de Crécy-Lagard V. 2010. Towards a systems approach in the genetic analysis of archaea: accelerating mutant construction and phenotypic analysis in *Haloferax volcanii*. *Archaea* **2010**: 426239. doi: 10.1155/2010/426239.
- Brand RC, Klootwijk J, Planta RJ, Maden BE. 1978. Biosynthesis of a hypermodified nucleotide in *Saccharomyces carlsbergensis* 17S and HeLa-cell 18S ribosomal ribonucleic acid. *Biochem J* **169**: 71–77.
- Buchhaupt M, Meyer B, Kötter P, Entian K-D. 2006. Genetic evidence for 18S rRNA binding and an Rps19p assembly function of yeast nucleolar protein Nep1p. *Mol Genet Genomics* **276**: 273–284.
- Cantara WA, Crain PF, Rozenski J, McCloskey JA, Harris KA, Zhang X, Vendeix FAP, Fabris D, Agris PF. 2011. The RNA modification database, RNAMDB: 2011 update. *Nucleic Acids Res* **39**: D195–D201.
- Chen H-Y, Yuan YA. 2010. Crystal structure of Mj1640/DUF358 protein reveals a putative SPOUT-class RNA methyltransferase. *J Mol Cell Biol* **2**: 366–374.
- Chenna R, Sugawara H, Koike T, Lopez R, Gibson TJ, Higgins DG, Thompson JD. 2003. Multiple sequence alignment with the Clustal series of programs. *Nucleic Acids Res* **31**: 3497–3500.
- Constantinesco F, Motorin Y, Grosjean H. 1999. Transfer RNA modification enzymes from *Pyrococcus furiosus*: detection of the enzymatic activities *in vitro*. *Nucleic Acids Res* **27**: 1308–1315.
- Crain PF. 1990. Preparation and enzymatic hydrolysis of DNA and RNA for mass spectrometry. *Methods Enzymol* **193**: 782–790.
- Czerwoniec A, Dunin-Horkawicz S, Purta E, Kaminska KH, Kasprzak JM, Bujnicki JM, Grosjean H, Rother K. 2009. MODOMICS: a database of RNA modification pathways. 2008 update. *Nucleic Acids Res* **37**: D118–D121.
- de Crécy-Lagard V, Brochier-Armanet C, Urbonavicius J, Fernandez B, Phillips G, Lyons B, Noma A, Alvarez S, Droogmans L, Armengaud J, et al. 2010. Biosynthesis of wyosine derivatives in tRNA: an ancient and highly diverse pathway in Archaea. *Mol Biol Evol* **27**: 2062–2077.
- Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* **32**: 1792–1797.
- Edmonds CG, Vestal ML, McCloskey JA. 1985. Thermospray liquid chromatography-mass spectrometry of nucleosides and of enzymatic hydrolysates of nucleic acids. *Nucleic Acids Res* **13**: 8197–8206.
- Edmonds CG, Crain PF, Gupta R, Hashizume T, Hocart CH, Kowalak JA, Pomerantz SC, Stetter KO, McCloskey JA. 1991. Posttranscriptional modification of tRNA in thermophilic archaea (Archaeobacteria). *J Bacteriol* **173**: 3138–3148.
- Grosjean H. 2009. Nucleic Acids are not boring long polymers of only four types of nucleotides. In *DNA and RNA modification enzymes: Structure, mechanism, function and evolution* (ed. H Grosjean), pp. 1–18. Landes Bioscience, Austin, TX.
- Grosjean H, Constantinesco F, Foiret D, Benachenhou N. 1995. A novel enzymatic pathway leading to 1-methylinosine modification in *Haloferax volcanii* tRNA. *Nucleic Acids Res* **23**: 4312–4319.
- Grosjean H, Edqvist J, Straby KB, Giegé R. 1996. Enzymatic formation of modified nucleosides in tRNA: dependence on tRNA architecture. *J Mol Biol* **255**: 67–85.
- Grosjean H, Droogmans L, Roovers M, Keith G. 2007. Detection of enzymatic activity of transfer RNA modification enzymes using radiolabeled tRNA substrates. *Methods Enzymol* **425**: 55–101.
- Grosjean H, Gaspin C, Marck C, Decatur WA, de Crécy-Lagard V. 2008a. RNomics and Modomics in the halophilic archaea *Haloferax volcanii*: identification of RNA modification genes. *BMC Genomics* **9**: 470. doi: 10.1186/1471-2164-9-470.
- Grosjean H, Gupta R, Maxwell ES. 2008b. Modified nucleotides in Archaeal RNAs. In *Archaea: New models for prokaryotic biology* (ed. P Blum), pp. 171–196. Horizon Scientific Press, Caister Academic Press, Cambridge, UK.
- Gu X-R, Nicoghossian K, Cedergren RJ, Wong JT-F. 1983. Sequences of halobacterial tRNAs and the paucity of U in the first position of their anticodons. *Nucleic Acids Res* **11**: 5433–5442.
- Gu X-R, Nicoghossian K, Cedergren RJ. 1984. Glycine and asparagine tRNA sequences from the archaeobacterium, *Methanobacterium thermoautotrophicum*. *FEBS Lett* **176**: 462–466.
- Gu X, Ivanetich KM, Santi DV. 1996. Recognition of the T-arm of tRNA by tRNA (m⁵U54)-methyltransferase is not sequence specific. *Biochemistry* **35**: 11652–11659.
- Gupta R. 1984. *Halobacterium volcanii* tRNAs: identification of 41 tRNAs covering all amino acids, and the sequences of 33 Class I tRNAs. *J Biol Chem* **259**: 9461–9471.
- Gupta R. 1986. Transfer RNAs of *Halobacterium volcanii*: sequences of five leucine and three serine tRNAs. *Syst Appl Microbiol* **7**: 102–105.
- Gupta R, Woese C. 1980. Unusual modification patterns in the transfer ribonucleic acids of archaebacteria. *Curr Microbiol* **4**: 245–249.
- Gurha P, Gupta R. 2008. Archaeal Pus10 proteins can produce both pseudouridine 54 and 55 in tRNA. *RNA* **14**: 2521–2527.
- Gurha P, Joardar A, Chaurasia P, Gupta R. 2007. Differential roles of archaeal box H/ACA proteins in guide RNA-dependent and independent pseudouridine formation. *RNA Biol* **4**: 101–109.
- Gutgsell N, Englund N, Niu L, Kaya Y, Lane BG, Ofengand J. 2000. Deletion of the *Escherichia coli* pseudouridine synthase gene *truB* blocks formation of pseudouridine 55 in tRNA *in vivo*, does not affect exponential growth, but confers a strong selective disadvantage in competition with wild-type cells. *RNA* **6**: 1870–1881.
- Joardar A, Gurha P, Skariah G, Gupta R. 2008. Box C/D RNA-guided 2'-O methylations and the intron of tRNA^{Trp} are not essential for the viability of *Haloferax volcanii*. *J Bacteriol* **190**: 7308–7313.
- Kilpatrick MW, Walker RT. 1981. The nucleotide sequence of the tRNA^{Met} from the archaebacterium *Thermoplasma acidophilum*. *Nucleic Acids Res* **9**: 4387–4390.
- Kinghorn SM, O'Byrne CP, Booth IR, Stansfield I. 2002. Physiological analysis of the role of *truB* in *Escherichia coli*: a role for tRNA modification in extreme temperature resistance. *Microbiology* **148**: 3511–3520.
- Kotlova N, Ishii TM, Zagryadskaya EI, Steinberg SV. 2007. Active suppressor tRNAs with a double helix between the D- and T-loops. *J Mol Biol* **373**: 462–475.
- Kowalak JA, Dalluge JJ, McCloskey JA, Stetter KO. 1994. The role of posttranscriptional modification in stabilization of transfer RNA from hyperthermophiles. *Biochemistry* **33**: 7869–7876.
- Kowalak JA, Bruenger E, Crain PF, McCloskey JA. 2000. Identities and phylogenetic comparisons of posttranscriptional modifications in 16S ribosomal RNA from *Haloferax volcanii*. *J Biol Chem* **275**: 24484–24489.

- Kuchino Y, Ihara M, Yabusaki Y, Nishimura S. 1982. Initiator tRNAs from archaeobacteria show common unique sequence characteristics. *Nature* **298**: 684–685.
- Lee TT, Agarwalla S, Stroud RM. 2005. A unique RNA fold in the RumA-RNA-cofactor ternary complex contributes to substrate selectivity and enzymatic function. *Cell* **120**: 599–611.
- Leulliot N, Bohnsack MT, Graille M, Tollervey D, Van Tilbeurgh H. 2008. The yeast ribosome synthesis factor Emg1 is a novel member of the superfamily of α/β knot fold methyltransferases. *Nucleic Acids Res* **36**: 629–639.
- Marck C, Grosjean H. 2002. tRNomics: analysis of tRNA genes from 50 genomes of Eukarya, Archaea, and Bacteria reveals anticodon-sparring strategies and domain-specific features. *RNA* **8**: 1189–1232.
- Meyer B, Wurm JP, Kötter P, Leisegang MS, Schilling V, Buchhaupt M, Held M, Bahr U, Karas M, Heckel A, et al. 2011. The Bowen-Conradi syndrome protein Nep1 (Emg1) has a dual role in eukaryotic ribosome biogenesis, as an essential assembly factor and in the methylation of Ψ 1191 in yeast 18S rRNA. *Nucleic Acids Res* **39**: 1526–1537.
- Motorin Y, Helm M. 2011. RNA nucleotide methylation. *Wiley Interdiscip Rev RNA* **2**: 611–631.
- Motorin Y, Muller S, Behm-Ansmant I, Branlant C. 2007. Identification of modified residues in RNAs by reverse transcription-based methods. *Methods Enzymol* **425**: 21–53.
- Nicoghossian K, Gu X-R, Cedergren R. 1985. *Halobacterium cutirubrum* tRNA sequences. *FEBS Lett* **193**: 255–260.
- Nishikura K, De Robertis EM. 1981. RNA processing in microinjected *Xenopus* oocytes. Sequential addition of base modifications in the spliced transfer RNA. *J Mol Biol* **145**: 405–420.
- Noon KR, Guymon R, Crain PF, McCloskey JA, Thomm M, Lim J, Cavicchioli R. 2003. Influence of temperature on tRNA modification in Archaea: *Methanococcoides burtonii* (Optimum Growth Temperature [T_{opt}], 23°C) and *Stetteria hydrogenophila* (T_{opt} , 95°C). *J Bacteriol* **185**: 5483–5490.
- Ny T, Björk GR. 1980. Cloning and restriction mapping of the *trmA* gene coding for transfer ribonucleic acid (5-methyluridine)-methyltransferase in *Escherichia coli* K-12. *J Bacteriol* **142**: 371–379.
- Ofengand J, Del Campo M, Kaya Y. 2001. Mapping pseudouridines in RNA molecules. *Methods* **25**: 365–373.
- Overbeek R, Begley T, Butler RM, Choudhuri JV, Chuang HY, Cohoon M, de Crécy-Lagard V, Diaz N, Disz T, Edwards R, et al. 2005. The subsystems approach to genome annotation and its use in the project to annotate 1000 genomes. *Nucleic Acids Res Symp Series* **33**: 5691–5702.
- Pang H, Ihara M, Kuchino Y, Nishimura S, Gupta R, Woese CR, McCloskey JA. 1982. Structure of a modified nucleoside in archaeobacterial tRNA which replaces ribosylthymine. 1-Methylpseudouridine. *J Biol Chem* **257**: 3589–3592.
- Romby P, Carbon P, Westhof E, Ehresmann C, Ebel JP, Ehresmann B, Giegé R. 1987. Importance of conserved residues for the conformation of the T-loop in tRNAs. *J Biomol Struct Dyn* **5**: 669–687.
- Ronquist F, Huelsenbeck JP. 2003. MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* **19**: 1572–1574.
- Tatusov R, Fedorova N, Jackson J, Jacobs A, Kiryutin B, Koonin E, Krylov D, Mazumder R, Mekhedov S, Nikolskaya A, et al. 2003. The COG database: an updated version includes eukaryotes. *BMC Bioinformatics* **4**: 41. doi: 10.1186/1471-2105-4-41.
- Taylor AB, Meyer B, Leal BZ, Kötter P, Schirf V, Demeler B, Hart PJ, Entian K-D, Wöhnert J. 2008. The crystal structure of Nep1 reveals an extended SPOUT-class methyltransferase fold and a pre-organized SAM-binding site. *Nucleic Acids Res* **36**: 1542–1554.
- Thomas SR, Keller CA, Szyk A, Cannon JR, LaRonde-LeBlanc NA. 2011. Structural insight into the functional mechanism of Nep1/Emg1 N1-specific pseudouridine methyltransferase in ribosome biogenesis. *Nucleic Acids Res* **39**: 2445–2457.
- Tkaczuk K, Dunin-Horkawicz S, Purta E, Bujnicki J. 2007. Structural and evolutionary bioinformatics of the SPOUT superfamily of methyltransferases. *BMC Bioinformatics* **8**: 73. doi: 10.1186/1471-2105-8-73.
- Urbonavicius J, Auxilien S, Walbott H, Trachana K, Golinelli-Pimpaneau B, Brochier-Armanet C, Grosjean H. 2008. Acquisition of a bacterial RumA-type tRNA(uracil-54, C5)-methyltransferase by Archaea through an ancient horizontal gene transfer. *Mol Microbiol* **67**: 323–335.
- Walbott H, Leulliot N, Grosjean H, Golinelli-Pimpaneau B. 2008. The crystal structure of *Pyrococcus abyssi* tRNA (uracil-54, C5)-methyltransferase provides insights into its tRNA specificity. *Nucleic Acids Res* **36**: 4929–4940.
- Wurm JP, Meyer B, Bahr U, Held M, Frolow O, Kötter P, Engels JW, Heckel A, Karas M, Entian KD, et al. 2010. The ribosome assembly factor Nep1 responsible for Bowen-Conradi syndrome is a pseudouridine-N1-specific methyltransferase. *Nucleic Acids Res* **38**: 2387–2398.
- Youvan DC, Hearst JE. 1981. A sequence from *Drosophila melanogaster* 18S rRNA bearing the conserved hypermodified nucleoside $m^1\Psi$: analysis by reverse transcription and high-performance liquid chromatography. *Nucleic Acids Res* **9**: 1723–1742.

RNA

A PUBLICATION OF THE RNA SOCIETY

The archaeal COG1901/DUF358 SPOUT-methyltransferase members, together with pseudouridine synthase Pus10, catalyze the formation of 1-methylpseudouridine at position 54 of tRNA

Kunal Chatterjee, Ian K. Blaby, Patrick C. Thiaville, et al.

RNA 2012 18: 421-433 originally published online January 24, 2012
Access the most recent version at doi:[10.1261/rna.030841.111](https://doi.org/10.1261/rna.030841.111)

Supplemental Material

<http://rnajournal.cshlp.org/content/suppl/2012/01/05/rna.030841.111.DC1>

Related Content

Identification of the enzyme responsible for N1-methylation of pseudouridine 54 in archaeal tRNAs

Jan Philip Wurm, Marco Giese, Ute Bahr, et al.
[RNA March , 2012 18: 412-420](#)

References

This article cites 63 articles, 15 of which can be accessed free at:
<http://rnajournal.cshlp.org/content/18/3/421.full.html#ref-list-1>

Articles cited in:

<http://rnajournal.cshlp.org/content/18/3/421.full.html#related-urls>

License

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).

To subscribe to *RNA* go to:
<http://rnajournal.cshlp.org/subscriptions>