Novel regulation and functional interaction of polycistronic miRNAs

MARY TRUSCOTT,¹ ABUL B.M.M.K. ISLAM,² and MAXIM V. FROLOV¹

¹Department of Biochemistry and Molecular Genetics, University of Illinois at Chicago, Chicago, Illinois 60607, USA

²Department of Genetic Engineering and Biotechnology, University of Dhaka, Dhaka 1000, Bangladesh

ABSTRACT

The importance of microRNAs in gene expression and disease is well recognized. However, what is less appreciated is that almost half of miRNA genes are organized in polycistronic clusters and are therefore coexpressed. The *mir-11~998* cluster consists of two miRNAs, miR-11 and miR-998. Here, we describe a novel layer of regulation that links the processing and expression of miR-998 to the presence of the *mir-11* gene. We show that the presence of miR-11 in the pri-miRNA is required for processing by Drosha, and deletion of *mir-11* prevents the expression of miR-998. Replacing *mir-11* with an unrelated miRNA rescued miR-998 expression in vivo and in vitro, as did expressing miR-998 from a shorter, more canonical miRNA scaffold. The embedded regulation of miR-998 is functionally important because unchecked miR-998 expression in the absence of miR-11 resulted in pleiotropic developmental defects. This novel regulation of expression of miRNAs within a cluster is not limited to the *mir-11~998* cluster and, thus, likely reflects the more general *cis*-regulation of expression of individual miRNAs. Collectively, our results uncover a novel layer of regulation within miRNA clusters that tempers the functions of the individual miRNAs. Unlinking their expression has the potential to change the expression of multiple miRNA targets and shift a biological response.

Keywords: polycistronic miRNA; miRNA cluster; *cis*-regulation of miRNA expression; miRNA biogenesis

INTRODUCTION

MicroRNAs (miRNAs) are small noncoding RNAs that repress the expression of target protein-coding transcripts. miRNAs are initially expressed as hairpins embedded within a primary transcript. The pri-miRNA structure is recognized by the Drosha/Pasha microprocessor and cleaved to release a pre-miRNA stem-loop, which is exported from the nucleus for further processing into a mature, functional miRNA (Kim et al. 2009). miRNAs regulate a variety of biological processes including development, proliferation, cell death, metabolism, and tissue homeostasis. Although earlier studies identified the stronger and more recognizable phenotypes of a few miRNAs (e.g., *let-7*), the majority of miRNAs exert subtle effects. It has been suggested that this is because miRNAs function to dampen noise arising from the stochastic nature of gene expression or raise the threshold required to elicit a response (Hornstein and Shomron 2006; Ebert and Sharp 2012; Ameres and Zamore 2013; Schmiedel et al. 2015). This may explain why it has been so difficult to assign a clear-cut function to many miRNAs and why inactivation of a particular miRNA rarely gives rise to severe defects.

miRNAs can be expressed from a separate transcript, they can be expressed from a transcript encoding both a miRNA

and a protein-coding gene, and they can be generated from a primary transcript encoding more than one miRNA (polycistronic). This latter case of miRNAs being expressed from clusters accounts for ~40% of miRNAs (Kim et al. 2009). The reason for and implications of such a genomic organization are not thoroughly understood. However, polycistronic clusters were shown to give rise to mature miRNAs that regulate overlapping or complementary sets of targets to elicit a biological response. There is mounting evidence that although the miRNAs from a cluster are processed from a common polycistronic RNA, their biogenesis can be differentially regulated and give rise to changes in relative levels of the mature miRNAs. The mechanism behind such regulation and settings where it is important are active areas of research.

The *mir-11~998* cluster consists of two miRNAs, miR-11 and miR-998, and it is embedded within the *Drosophila* *E2f1* gene. Both miRNAs are coexpressed with the host gene *E2f1* and were shown to modulate the apoptotic function of *E2f1*, albeit by different mechanisms. Here we report a novel layer of regulation between miR-11 and miR-998. Unexpectedly, the processing and expression of miR-998 strongly depends

Corresponding author: mfrolov@uic.edu

Article published online ahead of print. Article and publication date are at <http://www.rnajournal.org/cgi/doi/10.1261/rna.053264.115>.

© 2015 Truscott et al. This article is distributed exclusively by the RNA Society for the first 12 months after the full-issue publication date (see <http://rnajournal.cshlp.org/site/misc/terms.xhtml>). After 12 months, it is available under a Creative Commons License (Attribution-NonCommercial 4.0 International), as described at <http://creativecommons.org/licenses/by-nc/4.0/>.

on the presence of the *mir-11* gene. As a result, pre- and mature miR-998 are undetectable in the absence of *mir-11*. This dependence of miR-998 expression on *mir-11* is functionally important because breaking up this dependence and expressing miR-998 from its endogenous location in the absence of *mir-11* resulted in pleiotropic developmental defects at high frequency. Importantly, this interdependence in the expression of miRNAs within a cluster is not limited to the *mir11~998* cluster and likely reflects the more general *cis*-regulation of expression of individual miRNAs.

RESULTS

The expression of miR-998 is dependent on *mir-11*

To begin to characterize the expression of individual miRNAs from the *mir11~998* cluster we examined the levels of miR-998 and miR-11 in their respective and reciprocal mutant alleles, the *mir-11* deletion, *mir-11 Δ 1*, and *mir-998^{exc222}*, a *mir-998* mutant allele generated by imprecise *P*-element excision (Truscott et al. 2011, 2014). As expected, in both quantitative RT-PCR (Fig. 1A) and Northern blot analysis (Fig. 1B),

FIGURE 1. miR-998 expression is dependent on *mir-11* in vivo. (A) Quantitative RT-PCR was performed on RNA from third instar larvae of the following genotypes: Canton S (wild-type); *mir-11 Δ 1*; *mir-998^{exc222}*; *mir-11 Δ 1/mir-998^{exc222}*. The expression of miR-11 and miR-998 were measured by TaqMan assay and normalized to miR-14. (B) RNA from the samples in A was used in Northern blot analysis. Membranes were probed with ³²P-labeled miR-11, miR-998, or U6 anti-sense probes. (C) S2R+ cells were soaked with EGFP dsRNA or Drosha dsRNA for 4 d, and then transfected with the following Renilla luciferase sensor plasmids: pGL2-hsp70 min-hRluc; pGL2-hsp70 min-hRluc-mir-11~998; pGL2-hsp70 min-hRluc-mir-11 Δ ~998. Cells were harvested 30 h after transfection, and Renilla luciferase activity was measured. Results are presented as the mean of three separate luciferase measurements from each of three biological replicates. Error bars represent standard error.

miR-11 was expressed in wild-type animals, and in the *mir-998^{exc222}* mutant, but not in the *mir-11* deletion, *mir-11Δ1*. Furthermore, miR-11 expression was reduced in the *mir-11Δ1/mir-998^{exc222}* *trans*-heterozygous combination, demonstrating the sensitivity of the measurements.

Surprisingly, miR-998 expression was not detected in *mir-11Δ1* animals by qRT-PCR, even though the sequence of the *mir-998* gene was intact (Fig. 1A). This result was confirmed by Northern blot analysis that showed the absence of the mature miR-998 in *mir-11Δ1* animals (Fig. 1B). Significantly, no precursor miRNA of miR-998 (pre-miR-998) was detected, suggesting that the primary miRNA (pri-miR-998) is not processed in the absence of the *mir-11* gene. Importantly, the expression of miR-11 *in trans* in *mir-11Δ1/ mir-998^{exc222}* *trans*-heterozygous animals did not rescue miR-998 expression. Therefore, *mir-11* is necessary in a *cis* configuration for the expression of miR-998, and the regulation occurs at the level of the pri-miRNA transcript.

A pri-miRNA lacking the *mir-11* gene is not processed by Drosha

Results described above indicate that pri-miR-998 is not processed efficiently in the absence of the adjacent miR-11. Pri-miRNA transcripts are endonucleolytically processed by the Drosha/Pasha microprocessor complex, which binds the pri-miRNA in a manner that is dependent on secondary structure and possibly also on sequence (Lee et al. 2002, 2003; Han et al. 2006). To investigate whether the microprocessor processes miR-998 more efficiently in the presence of miR-11, we inserted the primary miR-11~998 transcript downstream from the luciferase gene in the 3' UTR, and transfected these luciferase sensors in S2R+ cells. If the primary miRNA is processed by the Drosha/Pasha microprocessor, the luciferase transcript is destabilized and degraded, which leads to a decrease in luciferase activity. However, if the primary miRNA is not processed by the microprocessor, the luciferase activity would be comparable to a sensor without a pri-miRNA in its 3' UTR (Fig. 1C).

The luciferase sensor containing the wild-type pri-miR-11~998 in the 3' UTR yielded less luciferase activity than the parental sensor control, which had no pri-miRNA in the 3' UTR (Fig. 1C) indicating that the luciferase assay accurately detects processing of wild-type pri-miR-11~998. In contrast, a luciferase sensor containing *pri-mir-11Δ1 ~998* sequence, which contains only the *mir-998* gene, was expressed at the same level as the control transcript with no pri-miRNA in the 3' UTR. Therefore, the *mir-998* gene on its own had no effect on luciferase transcript stability (Fig. 1C). Inactivating the microprocessor complex by treating the cells with dsRNA against Drosha increased the luciferase activity of the sensor carrying the wild-type *mir-11~998* primary miRNA, which is indicative of reduced processing of the miRNAs present in the 3' UTR. However, the knockdown of Drosha had no effect on the sensor carrying only the *mir-*

998 gene without the adjacent *mir-11*. This suggests that miR-998 is not processed by Drosha unless miR-11 is present. Together with the qRT-PCR and Northern blot analysis these results point to an unusual dependence of the processing of miR-998 on the presence of adjacent miR-11. When *mir-11* was deleted, neither pre- nor mature miR-998 was expressed.

miR-998 expression can be rescued by replacing *mir-11* with a heterologous miRNA

To gain further insight into the mechanism underlying the dependence of miR-998 on miR-11, we generated a series of constructs, expressed them using the baculovirus promoter in S2R+ cells, and followed the expression of mature miR-998 by these constructs using qRT-PCR. In a control experiment, miR-998 was expressed when a wild-type, 1042-bp fragment containing miR-11 and miR-998 was transfected. Similarly to the *in vivo* effects, and in the 3' UTR luciferase sensors (Fig. 1), miR-998 is not expressed when the miR-11 stem-loop is deleted (Fig. 2A). Therefore, the insect cell culture system accurately recapitulates the effects observed in mutant animals, and confirms that a portion of the intron in which *mir-11~998* is embedded contains the *cis*-regulatory sequence that modulates miR-998 expression.

The requirement of *mir-11* for miR-998 expression may be context-, sequence-, or structure-specific. We first asked whether miR-11 was required upstream of *mir-998* for miR-998 expression. *mir-11* and *mir-998* sequences were swapped and miRNA expression was measured. As shown in Figure 2B, miR-998 was expressed regardless whether the *mir-11* gene was upstream of *mir-998* (*mir-11~998*) or downstream from *mir-998* (*mir-998~11*). However, deletion of the *mir-11* gene (*mir-998~11Δ*) was sufficient to block miR-998 expression. Therefore, the relative order of the *mir-11* and *mir-998* genes does not influence the requirement of *mir-11* for miR-998 expression. However, an intact *mir-11* gene is required for miR-998 expression.

To determine whether the dependence of miR-998 expression is specific to the sequences in the *mir-11* gene, *mir-11* was replaced with a chimeric short hairpin miRNA (shmiR), *mir-1/mCherry*. miR-1/mCherry represses mCherry expression and was previously shown to be otherwise biologically inert (Haley et al. 2008). Strikingly, the presence of this heterologous miRNA was sufficient to rescue the expression of miR-998. Furthermore, similarly to the *mir-11* deletion, truncating the mCherry shmiR hairpin also resulted in the loss of expression of miR-998 (Fig. 2A). Therefore, a miRNA is required nearby for the expression of miR-998, but the precise sequence of the miRNA is not important.

One reason that, on its own, *mir-998* is not processed could be that its pre-miRNA scaffold is longer than the average (142 vs. 94 nt, respectively), which may serve as a poor substrate for the Drosha/Pasha microprocessor. To address this possibility, the mature miR-998 was expressed from a chimeric *mir-1/998* shmiR scaffold. In this shorter, more canonical miRNA

FIGURE 2. miR-998 expression is dependent on another miRNA nearby unless expressed from a shorter miRNA scaffold. S2R+ cells were transfected with constructs expressing wild-type and mutant *dE2f1 mir-11~998*-containing intron sequences as shown in the diagram. RNA extracts were made 2 d after transfection and mature miR-998 (A,B) and miR-11 (B) were measured by TaqMan assay and normalized to miR-14 or bantam expression. Expression was normalized to empty vector control (A) or wild-type *mir-11~998* (B). Error bars represent standard error.

scaffold, miR-998 was expressed on its own without the need for a miRNA gene nearby (Fig. 2A). Therefore, the miR-998 gene requires an adjacent miRNA gene in order to be expressed, and this is due, at least in part, to its unusually large backbone, because replacing it with a chimeric miR-1 scaffold was sufficient to rescue miR-998 expression.

Generation of *mir-11~998^{RMCE}* knock-in alleles that express miR-998 in the absence of miR-11

The striking dependence of miR-998 on miR-11 for expression raises the question of the functional significance of such an arrangement within the *mir-11~998* cluster. To address this question, a series of targeted alleles in the *mir-11~998* cluster was generated that express miR-998 in its endogenous location but do not express miR-11. We took advantage of the results obtained in tissue culture cells (Fig. 2) that showed that replacing *mir-11* with a heterologous miRNA (*mCherry shmiR*) or substituting the long scaffold of miR-998 (*mir-1/998*) allows the expression of miR-998 in the absence of *mir-11*.

These alleles were targeted to a chromosome that contains an *attP*-flanked mini-*white* cassette that had been in-

serted into the *mir-11* locus (Ge et al. 2012). Recombinase-mediated cassette exchange (RMCE) was used to replace the *attP*-flanked mini-*white* cassette in the *mir-11* locus (Ge et al. 2012) with either a cassette containing *mir-1/998* (*mir-11~998^{RMCE-1}*) or a cassette containing the *mCherry shmiR* and endogenous *mir-998* (*mir-11~998^{RMCE-2}*) (Fig. 3A). To confirm that miR-998 is expressed in these alleles, RNA was isolated from homozygous mutants and the level of miR-998 expression was determined by qRT-PCR. Although no miR-998 expression was detected in animals homozygous for the *mir-11* deletion (*mir-11^{KO(w-)}*), in *mir-11~998^{RMCE-1}* animals, miR-998 expression was 60% that of wild-type animals, and in *mir-11~998^{RMCE-2}* allele homozygotes, miR-998 was expressed at 30% of wild-type levels (Fig. 3B). Therefore, in agreement with the results in tissue culture cells, in vivo, miR-998 expression can be rescued by replacing the *mir-11* gene with an exogenous shmiR or by expressing miR-998 from the *mir-1* scaffold. Importantly, these alleles express miR-998 at a level similar to the level of the endogenous miR-998.

mir-11~998^{RMCE} alleles suppress E2F-dependent cell death in *rbf* mutant animals

The *mir-11~998^{RMCE}* alleles express miR-998 but do not express miR-11. Therefore, we began by characterizing these alleles in settings that are known to be sensitive to miR-998 expression but insensitive to miR-11 expression. One such context is the apoptotic response of *rbf* mutant cells during eye development. In *rbf* mutant eye discs, E2F activity is repressed, which leads to EGFR-dependent ectopic cell death in a narrow stripe anterior to the morphogenetic furrow (MF) of larval eye discs (Fig. 3C; Moon et al. 2006). Overexpression of miR-998 was previously shown to suppress apoptosis, whereas the loss of *mir-998* enhanced apoptosis in the MF of *rbf* mutant larval eye discs. In contrast, overexpression of miR-11 had no effect on apoptosis (Truscott et al. 2014).

Clones of *mir-11~998^{RMCE-1}* homozygous mutant cells were induced in the *rbf* mutant background and apoptotic cells were visualized with an antibody against active caspase-3. Notably, the high level of apoptosis in the morphogenetic furrow was strongly suppressed in *mir-11~998^{RMCE-1}* mutants (GFP negative) in comparison to the adjacent normal tissue (GFP positive). A similar result was obtained

with other allele *mir-11~998^{RMCE-2}* (Fig. 3C). The absence of cell death in the *mir-11~998^{RMCE}* alleles was not the consequence of changes in the expression of Rbf or E2f1 (data not shown, and Truscott et al. 2011, 2014).

Both *mir-11~998^{RMCE}* alleles express miR-998 in the absence of *mir-11*; therefore, the suppression of apoptosis could

be due to the lack of *mir-11* or due to the presence of miR-998. To distinguish between these two possibilities, clones of *mir-11^{KO(w-)}* mutant cells, which lack expression of both miR-11 and miR-998, were induced in *rbf* mutants, and apoptosis was examined by immunofluorescence. In contrast to strong suppression of apoptosis by the *mir-11~998^{RMCE}*

FIGURE 3. (Legend on next page)

alleles that express miR-998 in the absence of miR-11, the combined loss of miR-998 and miR-11 had no effect on the number of apoptotic cells in the MF of *rbf* mutant eye discs (Fig. 3C). These results strongly argue that the suppression of apoptosis by the *mir-11~998^{RMCE}* alleles is due to the expression of miR-998. Thus, altering the normal balance between miR-11 and miR-998 in the *mir-11~998^{RMCE}* alleles revealed that, in the absence of miR-11, unchecked miR-998 provides a strong prosurvival signal to suppress apoptosis in *rbf* mutants.

Given that miR-998 was shown to limit cell death in *rbf* mutants by increasing EGFR signaling, we performed genetic interaction tests between the *mir-11~998^{RMCE}* and EGFR alleles. The *Ellipse* allele is a gain-of-function allele of EGFR that results in a decrease in the number of ommatidial clusters and therefore a smaller eye. Consistently, the *mir-11~998^{RMCE}* alleles strongly enhanced the *Ellipse* phenotype, whereas *mir-11^{KO(w-)}*, which expresses neither miR-11 nor miR-998, had no effect (Fig. 3D).

Thus, we concluded that in the absence of miR-11, unchecked expression of miR-998 enhances EGFR signaling and protects *rbf* mutant cells from apoptosis. These results are in agreement with the previous analysis of miR-998 transgenic animals, thus providing genetic evidence that the *mir-11~998^{RMCE}* alleles accurately reflect the function of miR-998.

Expression of miR-998 in the absence of miR-11 results in pleiotropic effects and influences multiple phenotypic traits

The results described above suggest that the unchecked expression of miR-998 in the absence of miR-11 elicited the effects in *rbf* mutant cells that were not seen when miR-998 was coexpressed with miR-11. To determine the impact of unchecked miR-998 in normal animals, larval eye discs were dissected from a large cohort of the *mir-11~998^{RMCE}* larvae and stained with ELAV antibodies to visualize differentiating photoreceptors. Surprisingly, we noted a wide range of abnormalities in *mir-11~998^{RMCE}* animals, yet almost no strong defects were observed in animals mutant for *mir-998* only, or both *mir-11* and *mir-998*. The phenotypes associated with the *mir-11~998^{RMCE}* alleles included moderate and significant tissue remodeling (e.g., ectopic morphogenetic

furrow formation, tubular projections), whereas milder defects included changes in composition, spacing, or orientation of ommatidia and delayed morphogenetic furrow movement (summarized in Fig. 4A, with examples in Fig. 4B). Thus, expression of miR-998 in the absence of miR-11 exerts pleiotropic effects that appear to affect various phenotypic traits. Consistent with this interpretation, overexpression of miR-998 using the Gal4/UAS system gave rise to various developmental defects. Expression of miR-998 in the dorsal half of the larval eye disc (*DE > mir-998*) delayed the MF progression and led to projections resembling antennal discs. Similarly, *ptc > mir-998* eye discs led to what appear to be ectopic antennae in >50% of slow-developing animals (Fig. 4C).

To further characterize the *mir-11~998^{RMCE}* mutant animals, we performed RNA sequencing and compared changes in gene expression in the eye discs of the *mir-11~998^{RMCE}* alleles and the *mir-11^{KO(w-)}* allele, which expresses neither miR-11 nor miR-998. Notably, when compared with wild-type, there was a higher number of differentially expressed (DE) genes in the *mir-11~998^{RMCE}* mutants than in the *mir-11^{KO(w-)}* mutants, and more GOBP categories were enriched in the *mir-11~998^{RMCE}* versus WT samples compared with *mir-11^{KO(w-)}* versus WT (Fig. 4D; Supplemental Table S1). GOBP enrichment analysis using the list of DE genes in the *mir-11~998^{RMCE}* alleles that are also predicted miR-998 targets revealed overrepresentation of GOBP categories such as cell fate commitment and polarity. Importantly, such representation was not observed in *mir-998* mutants nor in mutants lacking both miR-11 and miR-998 (Fig. 4D; Truscott et al. 2014). Thus, in agreement with the pleiotropic nature of phenotypic traits associated with *mir-11~998^{RMCE}* alleles, genome-wide expression analysis revealed enrichment for various GOBP categories that may be associated with such phenotypes.

The interdependence of expression in miRNA clusters is not limited to *mir-11~998*

The results described above raise the question whether this type of regulation is specific to *mir-11~998* or whether there are other clusters of coexpressed miRNAs that demonstrate interdependence of miRNA expression. To test this idea, we selected the highly evolutionarily conserved the *mir-*

FIGURE 3. *mir-11~998^{RMCE}* alleles suppresses E2F-dependent cell death in *rbf* mutant animals. (A) Diagrams of the different mutant alleles in the context of the *dE2f1* locus. *mir-11^{KO(w+)}* contains an RMCE-competent cassette, was targeted to, and replaced the *mir-11* gene (see Ge et al. 2012 for more details). *mir-11^{KO(w-)}* is a *mir-11* deletion generated by Cre recombinase-mediated deletion of sequences between *loxP* sites (Ge et al. 2012). *mir-11~998^{RMCE-1}* and *mir-11~998^{RMCE-2}* are the products of RMCE-mediated replacement of the mini-*white* cassette with *E2f1* intron sequences in which the *mir-11* gene was replaced with *mir-1/998* or the *mir-1/mCherry shmiR* with wild-type *mir-998* described in Figure 2. (B) miR-998 expression was measured in adult heads of flies with the indicated genotypes by qRT-PCR and TaqMan assay. miR-998 expression was normalized to *rp49*. Results are expressed relative to wild-type control. Error bars represent standard error. (C) Clones of indicated mutant tissue were generated using *ey-FLP* in *rbf^{d20a}* hemizygous animals. The full genotypes are *rbf^{d20a} ey-FLP/Y; FRT 82B Ubi-GFP/FRT 82B Ubi-GFP, rbf^{d20a} ey-FLP/Y; FRT 82B mir-11~998^{RMCE-1}/FRT 82B Ubi-GFP, rbf^{d20a} ey-FLP/Y; FRT 82B mir-11~998^{RMCE-2}/FRT 82B Ubi-GFP, and rbf^{d20a} ey-FLP/Y; FRT 82B mir-11^{KO(w-)}/FRT 82B Ubi-GFP. Third instar larval eye discs were stained with an antibody recognizing active caspase (casp). The full disc is shown in the left panel, and a higher magnification is shown in panels on the right. A minimum of 10 larvae were analyzed for each genotype, and representative results are shown. (D) Images of adult eyes from wild-type control, *Egfr^{Elp/+}*, *Egfr^{Elp/+}; FRT 82B, mir-11~998^{RMCE-2}*, and *Egfr^{Elp/+}; FRT 82B, mir-11^{KO(w-)}* animals. Images were taken at the same magnification.*

FIGURE 4. Unchecked miR-998 activity leads to pleiotropic defects. (A) Third instar larval eye discs of the genotypes indicated were dissected and stained with anti-ELAV, anti-Ato, or anti-Pnt P1 antibodies. Changes in developmental patterning were assessed and scored as having either no, mild, moderate, or major defect. (B) Examples of defects observed and summarized in A. (C) Expression of miR-998 using the Gal4-UAS system. Third instar larval eye discs were dissected and stained with the antibodies indicated. A minimum of 10 larvae per genotype were assessed, a penetrance of phenotype >50%. (D) Heat map of GOBP enrichment analysis of genes differentially expressed in the indicated genotypes that are also predicted miR-998 targets. Categories associated with pleiotropic defects are shown. Full GOBP enrichment analysis is listed in Supplemental Table S1.

100~let-7~125 miRNA cluster. Notably, like the configuration of *mir-11~998*, *mir-125*, the 3'-most miRNA in the *mir-100~let-7~125* miRNA cluster, is longer than the aver-

age miRNA (109 vs. 94 nt, respectively). We therefore asked whether the expression of miR-125 is dependent on the other miRNAs in the cluster. The *mir-100~let-7~125* cluster was expressed in S2R+ cells, along with deletions of either *mir-100* or *let-7* or of *mir-100*, *let-7*, and the intervening sequence (Fig. 5). The expression of miR-125 did not vary significantly between the intact cluster and the two single miRNA deletions. This was in agreement with previous work investigating expression in vivo in genomic rescue constructs in which individual miRNAs were deleted (Sokol et al. 2008). However, deletion of both *mir-100* and *let-7* resulted in a threefold decrease in miR-125 expression. Therefore, like miR-998, miR-125 expression is enhanced by the presence of additional miRNA genes upstream, although the effect is not as dramatic as the one observed in the *mir-11~998* cluster. Nevertheless, these results suggest that interdependence between miRNAs in clusters extends beyond the *mir-11~998* cluster.

DISCUSSION

Here, we describe a novel layer of post-transcriptional regulation of polycistronic miRNAs, as illustrated by the *mir-11~998* cluster. Most clustered miRNAs are processed from a single polycistronic RNA and, therefore, they are often coexpressed. Unexpectedly, the processing of miR-998 is strongly dependent on the presence of *mir-11*. The *cis*-regulatory role of miR-11 ensures that pre-miR-998 is only expressed if pre-miR-11 is expressed. Therefore, miR-998 expression is kept in check by miR-11.

Although *mir-11* is required for the expression of miR-998, the converse is not true as *mir-998* is not required for miR-11 expression. The asymmetry in this intrinsic regulation may be explained in part by the longer than average length of the miR-998 pre-miRNA, 142 nt. Although *Drosophila* miRNAs range in length from 53 to 214 nt, almost 60%

of miRNAs are within 10 nt of the average length of 94 (Kozomara and Griffiths-Jones 2011). pri-miRNA cleavage by the microprocessor complex is determined by and

FIGURE 5. *Cis*-regulation is not limited to the *mir-11~998* cluster. S2R+ cells were transfected with constructs expressing wild-type and mutant *mir-100~125*, as shown in the diagram. RNA extracts were made 2 d after transfection and mature miR-125, miR-11, miR-998, and bantam were measured by TaqMan assay. miR-125 expression was normalized to miR-11 and miR-998 and to bantam expression. Expression was normalized to wild-type *mir-100~125*. Error bars represent standard error. Restriction sites were introduced flanking miRNAs for the purpose of generating miRNA deletions. Additional mutations were introduced to facilitate construction of gBlocks, and are indicated by tick marks. See Supplemental Figure S1 for full sequences.

measured from the ssRNA stem-loop junction (Han et al. 2006). Examination of the predicted structure of longer miRNAs including miR-998 using the M-fold folding algorithm (Zuker 2003) reveals a number of different conformations with similar free energies (data not shown). It is conceivable that the longer length of these miRNAs renders recognition and cleavage by the microprocessor more challenging. Therefore, longer miRNAs may be dependent on more canonical miRNAs in the vicinity for processing. This idea is supported by the finding that miR-998 expression became independent of the presence of an adjacent miRNA when miR-998 was expressed from the shorter *mir-1* scaffold. Similarly, the *mir-125* miRNA is longer than average, and its expression was significantly compromised when the two other miRNAs in the cluster, *mir-100* and *let-7*, were deleted. This raises a number of questions: How then are long monocistronic miRNAs expressed? What is the consequence of coordinated versus differential expression of miRNAs generated from polycistrons?

We considered addressing the latter question to be particularly important in this work because it would help to under-

stand biological significance of why expression of miR-998 is so tightly linked to expression of miR-11. Strikingly, breaking up the regulation within the *mir-11~998* cluster and expressing unchecked miR-998 in the absence of miR-11 resulted in a high frequency of pleiotropic phenotypes in ~20% of animals. Notably, these phenotypes are not due to the absence of miR-11 because they are suppressed by the loss of miR-998. The reason for such pleiotropy is unclear but it is consistent with the emerging view of miRNAs acting to impose subtle changes in transcript and protein expression, thereby reducing the overall noise in gene expression (Baek et al. 2008; Selbach et al. 2008; Mukherji et al. 2011). It is conceivable that unchecked miR-998 would affect the “precision” of the expression of its target genes that would eventually lead to a wide range of phenotypic effects. This explanation is supported by gene expression profiling that revealed a large number of differentially expressed genes of seemingly unrelated GOBP categories to be enriched in the *mir-11~998^{RMCE}* mutants. Furthermore, overexpression of miR-998 with several Gal4 drivers resulted in highly penetrant developmental defects indicating that, at least in overexpression settings, miR-998 can affect developmental pathways. Collectively

our data argue the functional importance of tightly linking the expression of miR-998 to the presence of miR-11.

Another important finding that we report here is that the intrinsic regulation within the miRNA clusters does not appear to be limited to the *mir-11~998* cluster as the expression of miR-125 from the *mir-100~let-7~125* cluster is similarly dependent on *mir-100* and *let-7*. Transcription of a polycistronic pri-miRNA permits coordinated expression of individual mature miRNAs. However, each step in miRNA biogenesis can be modulated, as can the steady-state level of the mature miRNA (Kim et al. 2009). For example, the individual pre-miRNAs from the *mir-17~92* cluster are differentially regulated by RNA-binding proteins: hnRNP A1 specifically binds pri-*mir-18* and enhances *mir-18* processing by Drosha (Guil and Cáceres 2007). Therefore, like post-transcriptional regulation of protein expression, miRNA expression can be fine-tuned in response to different biological cues, thereby affecting the timing and extent of miRNA target regulation.

Polycistronic miRNAs can give rise to miRNAs from the same or different miRNA families. These miRNAs can share

targets and functions or target genes with opposing functions. For example, the oncogenic *mir-17~92* cluster encodes miRNAs that target both positive and negative cell cycle regulators and pro- and anti-apoptotic proteins (O'Donnell et al. 2005; Hossain et al. 2006; Zhang et al. 2006; Shan et al. 2009; Li et al. 2012). Moreover, miR-92a and miR-17 have antagonistic effects: Although overexpression of miR-92a in mice caused erythroleukemia, coexpression of miR-17 abrogated this effect (Li et al. 2012). Similarly to the context-dependent differences in function of miRNAs from the *mir-17~92* cluster, we show here that miR-11 and miR-998 carry different functions. The functional consequences of coordinated or differential expression of the different individual miRNAs can determine very different biological outcomes. The impact of this layer of regulation on normal or perturbed developmental processes, as well as potential for therapeutic intervention awaits further investigation.

MATERIALS AND METHODS

Fly stocks

All fly crosses were done at 25°C. The following stocks were obtained from Bloomington *Drosophila* Stock Center at Indiana University: *egfr^{EL}*. The following stocks were previously published: *mir-11Δ1* (Truscott et al. 2011); *mir-998^{exc222}* (Truscott et al. 2014); *rbfl^{120a}* *ey-FLP/FM7 Act-GFP; FRT 82B Ubi-GFP* (Moon et al. 2006); *mir-11^{KO(w-)}* and *mir-11^{KO(w+)}* (Ge et al. 2012); *DE-Gal4* (Morrison and Halder 2010).

Construction of cell culture expression vectors

Insect cell culture expression constructs were generated using the pLex-7 Ek/LIC kit (Novagen). Ligation-independent cloning-competent inserts were generated as gBlocks (IDT). Insert sequences are in Supplemental Figure S1.

Cell culture, dsRNA knockdown, transfection, and luciferase assay

S2R+ cells were cultured in Schneider's Insect Medium (Gibco) + 10% FBS. dsRNA knockdown was done as described in Ambrus et al. (2009). Three days after initial soaking, cells were split and the following day, cells were transfected with the X-tremeGENE HP transfection reagent (Roche) according to manufacturer's protocol. Cells were harvested 24–48 h post-transfection. Renilla luciferase activity was measured using the Dual Luciferase Assay protocol (Promega).

Recombinase-mediated cassette exchange

piB-mir-1/998 and piB-mCherry shmiR, mir-998 donor plasmids were generated by replacing the GFP sequence from piB-GFP (Bateman et al. 2006). GFP sequence was removed with BamHI restriction sites and then filled in with T4 DNA polymerase. mir-1/998 and mCherry shmiR, mir-998 were removed from pLex-7 constructs by restriction digest and filling in with T4 DNA polymerase (see Supplemental Fig. S1 for sequence information). Embryo injection

was done at The Best Gene, Inc. Donor plasmid DNA was injected into F₁ from cross of *w; mir-11 pKO w⁺/TM3* to *M{vas-int. Dm}ZH-2A* (FBti0099694). Progeny were screened for loss of mini-white marker as described in Weng et al. (2009) and then sequenced to confirm insertion.

Northern blot analysis

Ten micrograms of total RNA extracted from third instar larvae using TRIzol (Invitrogen) was run on a 16% polyacrylamide gel with 7 M urea. Resolved RNA was transferred to Zeta-Probe membrane (Bio-Rad), followed by UV crosslinking, and prehybridizing in ULTRAhyb-Oligo (Invitrogen) at 42°C. RNA oligos (IDT) were 5'-end-labeled with [γ -³²P]ATP using T4 polynucleotide kinase (NEB). Hybridization was performed overnight at 42°C, followed by two washes in 2× SSC with 0.1% SDS, and one wash in 0.5× SSC with 0.1% SDS. Membranes were stripped of probes using boiling 0.1× SSC with 0.1% SDS.

Immunohistochemistry

Antibodies used were as follows: rabbit anti-C3 (Cleaved Caspase3), lot 26, 1:100 (Cell Signaling), rat anti-ELAV 1:50, (Developmental Studies Hybridoma Bank), rabbit anti-Pnt P1 (1:200) (Alvarez et al. 2003), rabbit anti-Atonal (1:2000) (from Y. Jan), and Cy3-, and Cy5- conjugated anti-mouse, and anti-rabbit secondary antibodies (Jackson ImmunoResearch Laboratories). Immunofluorescence was done as previously described (Ambrus et al. 2009) using a Zeiss Confocal microscope and images were prepared using Adobe Photoshop CS4. All images are confocal single-plane images unless otherwise stated as projection images.

miRNA target prediction

Comprehensive lists of predicted miR-998 targets and miR-11 targets were compiled from TargetScan (Ruby et al. 2007), MinoTar (Schnall-Levin et al. 2010), PITA (Kertesz et al. 2007), miRanda (Enright et al. 2003), and RNAhybrid (Rehmsmeier et al. 2004), and can be found in Truscott et al. (2014).

Gene expression analysis

For qRT-PCR, total RNA was isolated from 10 third instar larvae, or 30–50 third instar larval eye discs, with TRIzol (Invitrogen). Reverse transcription to measure standard mRNAs was performed using the SensiFAST cDNA Synthesis Kit (Bioline) according to the manufacturer's specifications. Quantitative PCR was performed with SensiFast SYBR No-Rox Mix (Bioline) info on a Roche Light Cycler 480. Mature miRNA expression was measured by TaqMan hydrolysis probe assay (Applied Biosystems) using the SensiFAST Probe No-ROX Kit (Bioline).

For genome-wide expression analysis, RNA was harvested from 30–50 third instar larval eye discs in biological duplicate from Canton S, *mir-11^{KO(w-)}*, *mir-11~998^{RMCE-1}*, and *mir-11~998^{RMCE-2}* and processed for Illumina HiSeq2000 Next Gen RNA Sequencing at the University of Chicago Genomics Facility. Data analysis is described in Supplementary Methods. RNA-seq data files are publicly available through the Gene Expression Omnibus (GEO; Accession number GSE73946).

SUPPLEMENTAL MATERIAL

Supplemental material is available for this article.

ACKNOWLEDGMENTS

We thank Nuria Lopez-Bigas, Barcelona Biomedical Research Park, Spain, for providing the cluster computing facility. We are grateful to Steve Cohen and the Bloomington Stock Center for fly stocks. We thank Zain Paroo for sharing protocols, equipment, and reagents for Northern blot experiments. We thank Jason Bateman for the piB-GFP plasmid, and Jim Skeath for the anti-Pnt P1 antibody. We are thankful to Steve Cohen and Zain Paroo for discussions. This research was supported by grants GM110018 and GM093827 from the National Institutes of Health to M.V.F., by a Scholar Award from the Leukemia & Lymphoma Society to M.V.F., and by a Postdoctoral Fellowship to M.T. from the Fonds de Recherches en Santé Québec.

Received July 12, 2015; accepted October 14, 2015.

REFERENCES

- Alvarez AD, Shi W, Wilson BA, Skeath JB. 2003. *pannier* and *pointedP2* act sequentially to regulate *Drosophila* heart development. *Development* **130**: 3015–3026.
- Ambrus AM, Rasheva VI, Nicolay BN, Frolov MV. 2009. Mosaic genetic screen for suppressors of the *de2f1* mutant phenotype in *Drosophila*. *Genetics* **183**: 79–92.
- Ameres SL, Zamore PD. 2013. Diversifying microRNA sequence and function. *Nat Rev Mol Cell Biol* **14**: 475–488.
- Baek D, Villén J, Shin C, Camargo FD, Gygi SP, Bartel DP. 2008. The impact of microRNAs on protein output. *Nature* **455**: 64–71.
- Bateman JR, Lee AM, Wu CT. 2006. Site-specific transformation of *Drosophila* via ϕ C31 integrase-mediated cassette exchange. *Genetics* **173**: 769–777.
- Ebert MS, Sharp PA. 2012. Roles for microRNAs in conferring robustness to biological processes. *Cell* **149**: 515–524.
- Enright AJ, John B, Gaul U, Tuschl T, Sander C, Marks DS. 2003. MicroRNA targets in *Drosophila*. *Genome Biol* **5**: R1.
- Ge W, Chen Y-W, Weng R, Lim SF, Buescher M, Zhang R, Cohen SM. 2012. Overlapping functions of microRNAs in control of apoptosis during *Drosophila* embryogenesis. *Cell Death Differ* **19**: 839–846.
- Guil S, Cáceres JF. 2007. The multifunctional RNA-binding protein hnRNP A1 is required for processing of miR-18a. *Nat Struct Mol Biol* **14**: 591–596.
- Haley B, Hendrix D, Trang V, Levine M. 2008. A simplified miRNA-based gene silencing method for *Drosophila melanogaster*. *Dev Biol* **321**: 482–490.
- Han J, Lee Y, Yeom K-H, Nam J-W, Heo I, Rhee J-K, Sohn SY, Cho Y, Zhang B-T, Kim VN. 2006. Molecular basis for the recognition of primary microRNAs by the Drosha-DGCR8 complex. *Cell* **125**: 887–901.
- Hornstein E, Shomron N. 2006. Canalization of development by microRNAs. *Nat Genet* **38**: S20–S24.
- Hossain A, Kuo MT, Saunders GF. 2006. Mir-17-5p regulates breast cancer cell proliferation by inhibiting translation of AIB1 mRNA. *Mol Cell Biol* **26**: 8191–8201.
- Kertesz M, Iovino N, Unnerstall U, Gaul U, Segal E. 2007. The role of site accessibility in microRNA target recognition. *Nat Genet* **39**: 1278–1284.
- Kim VN, Han J, Siomi MC. 2009. Biogenesis of small RNAs in animals. *Nat Rev Mol Cell Biol* **10**: 126–139.
- Kozomara A, Griffiths-Jones S. 2011. miRBase: integrating microRNA annotation and deep-sequencing data. *Nucleic Acids Res* **39**: D152–D157.
- Lee Y, Jeon K, Lee J-T, Kim S, Kim VN. 2002. MicroRNA maturation: stepwise processing and subcellular localization. *EMBO J* **21**: 4663–4670.
- Lee Y, Ahn C, Han J, Choi H, Kim J, Yim J, Lee J, Provost P, Rådmark O, Kim S, et al. 2003. The nuclear RNase III Drosha initiates microRNA processing. *Nature* **425**: 415–419.
- Li Y, Vecchiarelli-Federico LM, Li Y-J, Egan SE, Spaner D, Hough MR, Ben-David Y. 2012. The miR-17-92 cluster expands multipotent hematopoietic progenitors whereas imbalanced expression of its individual oncogenic miRNAs promotes leukemia in mice. *Blood* **119**: 4486–4498.
- Moon N-S, Di Stefano L, Dyson N. 2006. A gradient of epidermal growth factor receptor signaling determines the sensitivity of *rbf1* mutant cells to E2F-dependent apoptosis. *Mol Cell Biol* **26**: 7601–7615.
- Morrison CM, Halder G. 2010. Characterization of a dorsal-eye Gal4 Line in *Drosophila*. *Genesis* **48**: 3–7.
- Mukherji S, Ebert MS, Zheng GXY, Tsang JS, Sharp PA, van Oudenaarden A. 2011. MicroRNAs can generate thresholds in target gene expression. *Nat Genet* **43**: 854–859.
- O'Donnell KA, Wentzel EA, Zeller KI, Dang CV, Mendell JT. 2005. c-Myc-regulated microRNAs modulate E2F1 expression. *Nature* **435**: 839–843.
- Rehmsmeier M, Steffen P, Höchsmann M, Giegerich R. 2004. Fast and effective prediction of microRNA/target duplexes. *RNA* **10**: 1507–1517.
- Ruby JG, Stark A, Johnston WK, Kellis M, Bartel DP, Lai EC. 2007. Evolution, biogenesis, expression, and target predictions of a substantially expanded set of *Drosophila* microRNAs. *Genome Res* **17**: 1850–1864.
- Schmiedel JM, Klemm SL, Zheng Y, Sahay A, Blüthgen N, Marks DS, van Oudenaarden A. 2015. Gene expression. MicroRNA control of protein expression noise. *Science* **348**: 128–132.
- Schnall-Levin M, Zhao Y, Perrimon N, Berger B. 2010. Conserved microRNA targeting in *Drosophila* is as widespread in coding regions as in 3' UTRs. *Proc Natl Acad Sci* **107**: 15751–15756.
- Selbach M, Schwanhäusser B, Thierfelder N, Fang Z, Khanin R, Rajewsky N. 2008. Widespread changes in protein synthesis induced by microRNAs. *Nature* **455**: 58–63.
- Shan SW, Lee DY, Deng Z, Shatseva T, Jeyapalan Z, Du WW, Zhang Y, Xuan JW, Yee S-P, Siragam V, et al. 2009. MicroRNA MiR-17 retards tissue growth and represses fibronectin expression. *Nat Cell Biol* **11**: 1031–1038.
- Sokol NS, Xu P, Jan Y-N, Ambros V. 2008. *Drosophila* let-7 microRNA is required for remodeling of the neuromusculature during metamorphosis. *Genes Dev* **22**: 1591–1596.
- Truscott M, Islam ABMMK, López-Bigas N, Frolov MV. 2011. mir-11 limits the proapoptotic function of its host gene, *de2f1*. *Genes Dev* **25**: 1820–1834.
- Truscott M, Islam ABMMK, Lightfoot J, López-Bigas N, Frolov MV. 2014. An intronic microRNA links Rb/E2F and EGFR signaling. *PLoS Genet* **10**: e1004493.
- Weng R, Chen Y-W, Bushati N, Cliffe A, Cohen SM. 2009. Recombinase-mediated cassette exchange provides a versatile platform for gene targeting: knockout of miR-31b. *Genetics* **183**: 399–402.
- Zhang L, Huang J, Yang N, Greshock J, Megraw MS, Giannakakis A, Liang S, Naylor TL, Barchetti A, Ward MR, et al. 2006. microRNAs exhibit high frequency genomic alterations in human cancer. *Proc Natl Acad Sci* **103**: 9136–9141.
- Zuker M. 2003. Mfold web server for nucleic acid folding and hybridization prediction. *Nucleic Acids Res* **31**: 3406–3415.

RNA

A PUBLICATION OF THE RNA SOCIETY

Novel regulation and functional interaction of polycistronic miRNAs

Mary Truscott, Abul B.M.M.K. Islam and Maxim V. Frolov

RNA 2016 22: 129-138 originally published online November 9, 2015

Access the most recent version at doi:[10.1261/rna.053264.115](https://doi.org/10.1261/rna.053264.115)

Supplemental Material

<http://rnajournal.cshlp.org/content/suppl/2015/11/04/rna.053264.115.DC1>

References

This article cites 35 articles, 15 of which can be accessed free at:
<http://rnajournal.cshlp.org/content/22/1/129.full.html#ref-list-1>

Creative Commons License

This article is distributed exclusively by the RNA Society for the first 12 months after the full-issue publication date (see <http://rnajournal.cshlp.org/site/misc/terms.xhtml>). After 12 months, it is available under a Creative Commons License (Attribution-NonCommercial 4.0 International), as described at <http://creativecommons.org/licenses/by-nc/4.0/>.

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).

To subscribe to *RNA* go to:

<http://rnajournal.cshlp.org/subscriptions>