ERRATUM

RNA 22: 839–851 (2016)

Erratum: How many biological replicates are needed in an RNA-seq experiment and which differential expression tool should you use?

NICHOLAS J. SCHURCH, PIETÀ SCHOFIELD, MAREK GIERLIŃSKI, CHRISTIAN COLE, ALEXANDER SHERSTNEV, VIJENDER SINGH, NICOLA WROBEL, KARIM GHARBI, GORDON G. SIMPSON, TOM OWEN-HUGHES, MARK BLAXTER, and GEOFFREY J. BARTON

In the above-noted article, the following information should have been included in the Acknowledgments section:

“We acknowledge funding from the Wellcome Trust (095062; 098439/Z/12; 92530/Z/10/Z; WT097945; WT092340; WT083481), the Medical Research Council (MR/K001744/1), and the Biotechnology and Biological Sciences Research Council (BB/M004155/1; BB/H002286/1).”

This article has been corrected in both the PDF and full-text HTML files online.

doi: 10.1261/rna.058339.116

RNA

A PUBLICATION OF THE RNA SOCIETY

Erratum: How many biological replicates are needed in an RNA-seq experiment and which differential expression tool should you use?

Nicholas J. Schurch, Pietà Schofield, Marek Gierlinski, et al.

RNA 2016 22: 1641

Related Content

How many biological replicates are needed in an RNA-seq experiment and which differential expression tool should you use?

Nicholas J. Schurch, Pietà Schofield, Marek Gierlinski, et al.

[RNA June , 2016 22: 839-851](#)

Open Access

Freely available online through the *RNA* Open Access option.

License

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).

To subscribe to *RNA* go to:

<http://rnajournal.cshlp.org/subscriptions>