Regulation of the vitamin B₁₂ metabolism and transport in bacteria by a conserved RNA structural element

ALEXEY G. VITRESCHAK,^{1,2} DMITRY A. RODIONOV,^{1,3} ANDREY A. MIRONOV,^{1,3} and MIKHAIL S. GELFAND^{1,3}

¹Integrated Genomics–Moscow, 117333, Russia

²Institute for Problems of Information Transmission, Moscow, 101447, Russia

³State Scientific Center GosNIIGenetika, Moscow, 113545, Russia

ABSTRACT

Cobalamin in the form of adenosylcobalamin (Ado-CBL) is known to repress expression of genes for vitamin B₁₂ biosynthesis and be transported by a posttranscriptional regulatory mechanism, which involves direct binding of Ado-CBL to 5′ untranslated gene regions (5′UTR). Using comparative analysis of genes and regulatory regions, we identified a highly conserved RNA structure, the *B12*-element, which is widely distributed in 5′UTRs of vitamin B₁₂-related genes in eubacteria. Multiple alignment of approximately 200 *B12*-elements from 66 bacterial genomes reveals their common secondary structure and several extended regions of sequence conservation, including the previously known *B12*-box motif. In analogy to the model of regulation of the riboflavin and thiamin biosynthesis, we suggest Ado-CBL-mediated regulation based on formation of alternative RNA structures including the *B12*-element. In Gram-negative proteobacteria, as well as in cyanobacteria, actinobacteria, and the CFB group, the cobalamin biosynthesis and vitamin B₁₂ transport genes are predicted to be regulated by inhibition of translation initiation, whereas in the *Bacillus/Clostridium* group of Gram-positive bacteria, these genes seem to be regulated by transcriptional antitermination. Phylogenetic analysis of the *B12*-elements reveals a large number of likely duplications of *B12*-elements in several bacterial genomes. These lineage-specific duplications of RNA regulatory elements seem to be a major evolutionary mechanism for expansion of the vitamin B₁₂ regulon.

Keywords: Bacteria; comparative genomics; regulatory RNA; *B12*-element; cobalamin

INTRODUCTION

Synthesized only by prokaryotic organisms, vitamin B₁₂ or cobalamin (CBL) is an essential cofactor for several important enzymes that catalyze a variety of transmethylation and rearrangement reactions (Martens et al. 2002). Expression of the *Salmonella typhimurium cob* operon, encoding the CBL biosynthetic pathway, and of the *btuB* gene of *Escherichia coli* and *S. typhimurium*, encoding the vitamin B₁₂ transporter, is repressed by addition of vitamin B₁₂ by a post-transcriptional regulatory mechanism (Lundrigan et al. 1991; Richter-Dahlfors and Andersson 1992). As shown by deletion analysis, this regulation requires unusually long 5′-untranslated leader sequences of the corresponding mRNAs, which contain several conserved elements. The leader mRNAs of the *cob* and *btuB* genes contain an evolutionarily

conserved sequence known as the *B12*-box (Franklund and Kadner 1997). Translational regulation of these genes requires also a conserved RNA hairpin that would mask the ribosome-binding site (RBS), thus inhibiting translation initiation and gene expression (Richter-Dahlfors et al. 1994; Nou and Kadner 1998). In addition, a *cis*-acting translational enhancer element in the *cob* leader mRNA is absolutely required to unfold the inhibitory RBS hairpin in the absence of coenzyme B₁₂ (Ravnum and Andersson 2001). In the same work, a secondary structure model for the *cob* leader mRNA was obtained from the chemical probing experiments with single-stranded RNA-modifying agents and combined with the output of a RNA folding computer program. No cobalamin-regulatory genes were identified in bacteria, but it was shown that adenosylcobalamin (Ado-CBL) is an effector molecule involved in the regulation of CBL genes (Nou and Kadner 2000). Recently, the structure-dependent spontaneous cleavage of RNA technique was applied to the *E. coli btuB* leader sequence in the presence and absence of Ado-CBL, and it was shown that this sequence fragment can directly bind Ado-CBL, consequently undergoing conformational changes in the secondary and tertiary

Reprint requests to: Mikhail Gelfand, GosNIIGenetika, 1st Dorozhny pr., 1 Moscow 113545 Russia; e-mail: gelfand@ig-msk.ru; fax: +7-095-3150501.

Article and publication are at <http://www.rnajournal.org/cgi/doi/10.1261/rna.5710303>.

structure of the RNA (Nahvi et al. 2002). The investigators suggested that the mechanism of the *btuB* regulation involves formation of two alternative RNA structures, repressing and antirepressing, in the presence and absence of Ado-CBL, respectively.

The comparative analysis is a powerful approach to identification of regulatory patterns in bacterial genomes (Gelfand et al. 2000). Comparative sequence analysis was used for prediction of conserved RNA secondary structures (Eddy and Durbin 1994; Marck and Grosjean 2002) and detection of novel regulatory RNA elements, for instance, iron-responsive elements in *E. coli* (Dandekar et al. 1998) or S-boxes in Gram-positive bacteria (Grundy and Henkin 1998). In such studies, analysis of complementary substitutions in aligned sequences is used to construct a single conserved structure. The number of known noncoding RNA families is expanding rapidly. This resulted in development of a number of specialized databases, in particular the RNA family database Rfam (Griffiths-Jones et al. 2003), the database of known RNA structures RNABase (Murthy and Rose 2003), and the noncoding RNAs database (Szymanski et al. 2003). Current availability of multiple complete genomes provides an opportunity to identify consensus RNA elements upstream of co-regulated genes. In particular, highly conserved *RFN* and *THI* elements were identified in the regulatory regions of genes involved in the biosynthesis of two different vitamins, riboflavin and thiamin, respectively (Vitreschak et al. 2002; Rodionov et al. 2002). Recently it was confirmed that these RNA elements control expression of the target genes through a post-transcriptional regulatory mechanism (Mironov et al. 2002; Winkler et al. 2002a,b).

Here we applied the comparative approach to analysis of 5'-untranslated regions (5'UTRs) of vitamin *B₁₂*-related genes in ~100 prokaryotic genomes. We report identification of a novel conserved RNA element involved in regulation of *B₁₂*-related genes in eubacteria. This new 5'UTR regulatory RNA, named the *B12*-element, is highly conserved on the sequence and structural levels and includes the previously defined *B12*-box motif. The *B12*-elements are widely distributed in bacteria: ~200 elements were identified in 5'UTRs of *B₁₂*-related genes in 67 bacterial genomes. The common structure of the *B12*-element was inferred and a possible mechanism of the *B12*-element-mediated regulation involving either transcriptional or translational attenuation was proposed for different groups of bacteria.

RESULTS AND DISCUSSION

Conserved structure of the *B12*-element

The *btuB* genes of *E. coli* and *S. typhimurium* have extensive regulatory regions including the conserved *B12*-box sequence (Ravnum and Andersson 1997; Nou and Kadner 2000). We started with identification of orthologs of *btuB* in

related bacteria. The upstream regions of *btuB* orthologs were aligned by the RNAMultAln program and the conserved RNA secondary structure was identified. This novel RNA structure, named the *B12*-element, consist of a number of helices and conserved sequence motifs, including the known *B12*-box. We constructed a pattern, which corresponded to the identified *B12*-element and scanned 107 genomic sequences using the RNA-PATTERN program. As a result, we found ~200 *B12*-elements in 67 bacterial genomes. Multiple alignment of these elements is shown in Figure 1. The *B12*-element is widely distributed in eubacteria, but it has not been observed in Archaea and Eukaryota. Most eubacterial genomes, containing the CBL biosynthesis or transport genes, have 1 to 13 *B12*-elements. The distribution of the elements is different in various taxonomic groups. On the average, Gram-negative α -proteobacteria have five *B12*-elements per genome, γ - and β -proteobacteria have two elements, Gram-positive bacteria have three elements, and cyanobacteria have only one *B12*-element per genome. All of these elements are located upstream of *B₁₂*-related genes. The detailed functional, positional, and phylogenetic analysis of the CBL genes is submitted elsewhere (D.A. Rodionov, A.G. Vitreschak, A.A. Mironov, and M.S. Gelfand, in prep.).

Previously, we described highly conserved 5'UTR structures, namely the *RFN* and *THI* elements, involved in regulation of riboflavin- and thiamin-related genes, respectively (Rodionov et al. 2002; Vitreschak et al. 2002). Similarly to these elements, the *B12*-element has a set of helices closed by a single base stem, and regions of high sequence conservation that are distributed along the entire element (Fig. 2). The conserved core of *B12*-element consists of seven helices (P0 to P6) and single-stranded regions with a high degree of sequence conservation. Existence of the conserved helices is confirmed by compensatory substitutions in base-paired positions. In addition to the conserved core, the *B12*-element has a number of facultative nonconserved stem-loops, designated Add-I and Add-II, and one internal variable structure, named VS. The previously defined *B12*-box is situated within the long internal loop between P0 and P4, slightly overlapping the base stem P0. In a number of cases a part of the *B12* box can fold into a facultative stem-loop structure (Fig. 2). Other identified conserved regions of the *B12*-element are located upstream of the *B12*-box. The first one stretches from P1 to P4 and contains an AG-rich internal loop between P1 and P2 with consensus 5'-AAN AGGGAA-3'. Interestingly, AG-rich conserved regions were also observed in internal loops of the *RFN* and *THI* elements. Two other conserved regions, with consensi 5'-GC CACTG-3' and 5'-YGGGAAGGC-3', are located between P5 and P6, partially overlapping these helices.

Beside the conserved core structure, the *B12*-element includes additional nonconserved stem-loops, Add-I and Add-II. The former seems to be correlated with phylogeny, as it occurs only in genomes of proteobacteria. In contrast,

FIGURE 2. Conserved structure of the B_{12} -element. Capital letters indicate invariant positions. Lowercase letters indicate strongly conserved positions. Degenerate positions: R = A or G; Y = C or U; K = G or U; M = A or C; H = not G; D = not C; N = any nucleotide. Dashes and black dots indicate obligatory and facultative base pairs, respectively. The conserved helices are numbered P0 to P6. Stem-loops of variable and constant lengths are shown by broken and solid lines, respectively. Additional stem-loops are designated Add-I and Add-II. The sequence of the conserved B12-box motif is shaded in gray. The variable structure VS separates two conserved parts of the B_{12} -element, BI and BII. Taxonomic variations of the VS topology are shown in the inset.

the presence of the latter stem-loop does not depend on the phylogenetic position of a genome and occurs in <10% of cases. The maximal observed lengths of the Add-I and Add-II stem-loops are 34 and 122 nucleotides, respectively.

The B_{12} -elements can be classified into two major types based on the existence of a highly conserved stem-loop region, named BII. This part of the B_{12} -element includes

conserved regions, 5'-GCCACTG-3' and 5'-YGGGAAGGC-3', which partially overlap the P5 and P6 helices (Fig. 2). Although most B_{12} -elements are complete, the BII part is absent in a number of genomes: in all B_{12} -elements of cyanobacteria, bacteria *Deinococcus radiodurans*, *Bacillus subtilis*, *Shewanella oneidensis*, *Chloroflexus aurantiacus*, as well as in some B_{12} -elements of actinobacteria and *Pseudomonas aeruginosa* (Table 1). Short B_{12} -elements without the BII part were found upstream of some B_{12} -related genes and, therefore, are functional. In complete B_{12} -elements, the BII part is separated from other conserved parts of the B_{12} -element by a variable linker, which can fold into a nonconserved variable structure, named VS. This structure has different topologies in various taxonomic groups of bacteria. For example, the VS structures are represented by one helix in bacteria from the *Bacillus/Clostridium* group, whereas VS of γ -proteobacteria consist of two adjacent helices. In other cases the VS structure is more complex

(see the lower frame in Fig. 2). Thus, the additional highly conserved BII part can play either an auxiliary role in the function of the B_{12} -element or perform some other function.

The previously proposed secondary structures for the *cbiA* (Ravnum and Andersson 2001) and *btuB* (Nahvi et al. 2002) leader mRNAs from enterobacteria differ in some

TABLE 1. Phylogenetic distribution of B_{12} -elements in bacteria

Taxonomic group	Genomes with B_{12} -elements	Number of B_{12} -elements	Type of B_{12} -elements		Proposed type of regulation
			complete	without BII	
α -proteobacteria	11	58	all	no	Translational
β -proteobacteria	5	11	all	no	Translational
γ -proteobacteria	14	31	yes	3	Translational
δ -proteobacteria (GME)	1	1	all	no	Translational/transcriptional
The <i>Bacillus/Clostridium</i> group	14	37	yes	1	Transcriptional
Actinobacteria	7	22	yes	6	Translational
Cyanobacteria	5	8	no	all	Translational
The <i>Thermus/Deinococcus</i> group	1	3	no	all	Translational
The CFB group (BX, PG, CL)	3	17	all	no	Translational
Spirochetes (LI, TDE)	2	5	all	no	?
<i>Chloroflexaceae</i> (CAU)	1	2	no	all	Translational
<i>Fusobacteriaceae</i> (FN)	1	2	all	no	Transcriptional
<i>Thermotogales</i> (TM)	1	1	all	no	Transcriptional
Total	66	198	174	23	Transcriptional ~20%; translational ~80%

details in their topology. Here we propose a conserved secondary structure for 5'UTRs of B_{12} -regulated genes. Similarly to the riboflavin- and thiamin-specific 5'UTR regulatory RNAs, *RFN* and *THI* elements, this new B_{12} -specific RNA has a compact secondary structure, consisting of a set of conserved helices closed by a single base stem. The previously proposed RNA secondary structure of the *cbiA* regulatory region (Ravnum and Andersson 2001) is not compact, although a number of helices in both structures coincide (namely, P1, P2, P3, and P6). Moreover, despite of compactness of the predicted secondary structure of the *btuB* regulatory region (Nahvi et al. 2002) and similarity in topology of both structures, this structure differs from the B_{12} -element in regions of base pairing of the base stem. However, the suggested conserved structure of the B_{12} -element is mostly consistent with the chemical probing data for the *btuB* and *cbiA* leader regions (Fig. 3).

Regulation of transcription/translation mediated by B_{12} -element

Analysis of the leader sequences of B_{12} -regulated genes allowed us to predict additional RNA regulatory hairpins

downstream of all B_{12} -elements. In some cases these additional hairpins are followed by runs of thymidines and therefore are candidate ρ -independent terminators of transcription. In other cases the hairpins overlap the ribosome-binding site (RBS) of the first gene in the B_{12} -regulated operon and are candidate translational sequesters that prevent ribosome binding to the RBS. Previously, it has been shown that RBS-sequestering hairpins are involved in the regulation of translation of the *btuB* gene of *E. coli* and *S. typhimurium*, as well as the *cbiA* gene of *S. typhimurium*. (Ravnum and Andersson 1997; Nou and Kadner 2000). Most Gram-positive bacteria from the *Bacillus/Clostridium* group, *Chloroflexus aurantiacus*, *Fusobacterium nucleatum*, and *Thermotoga maritima* have a candidate terminator hairpin, whereas most Gram-negative bacteria (proteobacteria and the CFB group), as well as actinomycetes, cyanobacteria, and *D. radiodurans* have candidate RBS-sequestering hairpins downstream of B_{12} -elements. Therefore, the regulation of B_{12} -related genes is likely to operate mainly at the level of transcription in the former group of bacteria and at the level of translation in the latter group (Table 1). Thus, the phylogenetic distribution of the proposed terminators and sequesters in Gram-positive and Gram-negative bacteria, respectively, is similar to previously

observed distribution of the regulatory hairpins in the riboflavin and thiamin regulons (Vitreschak et al. 2002; Rodionov et al. 2002). Analysis of the 5'UTR of the cobalamin biosynthetic operon from *Geobacter metallireducens* reveals two possibilities of regulation. In this case the predicted terminator hairpin overlaps the RBS sequence of the first gene in the operon, and therefore, it can function as both a terminator and a sequester.

The mechanism of the Ado-CBL-dependent regulation involves formation of two alternative RNA conformations. It has been shown in experiments that the RBS sequester of the *S. typhimurium cbi* operon forms in the presence of Ado-CBL, leading to inhibition of the translation initiation, whereas the anti-sequester conformation formed in the absence of Ado-CBL opens the RBS sequence and allows translation to initialize (Ravnum and Andersson 2001).

Previously we proposed the riboswitch mechanism involving the *RFN* and *THI* elements. These elements are stabilized by effector molecules (flavin mononucleotide or thiamin pyrophosphate, respectively), allowing formation of downstream regulatory hairpins

FIGURE 3. Predicted RNA secondary structure of the B_{12} -elements upstream of *btuB* from *E. coli* and *cbiA* from *S. typhimurium* genes. Chemical modifications of the B_{12} -loop region of the *btuB* (A) and *cbiA* (B) 5'UTRs are shown by filled triangles (Ravnum and Andersson 2001; Nahvi et al. 2002).

(transcriptional terminator or a hairpin sequestering the RBS site) that leads to repression of transcription or translation, respectively. In the absence of the effector, a more energetically favorable RNA structure that is alternative to parts of the vitamin-specific element and the regulatory hairpin can fold, thus releasing gene expression (Rodionov et al. 2002; Vitreschak et al. 2002). This riboswitch mechanism was experimentally confirmed for thiamin and riboflavin regulons (Mironov et al. 2002; Winkler et al. 2002a,b). In analogy with the riboflavin- and thiamin-specific riboswitches, we propose here the cobalamin-specific riboswitch.

It seems to have some specific properties. It has been experimentally shown that an additional regulatory element is located between the *B12*-element and the sequestor hairpin of the *cbiA* gene of *S. typhimurium* (Ravnum and Andersson 2001). This element, a translational enhancer, plays the key role in the regulation of translation initiation of *cbiA*. In the absence of cobalamin it interacts with the region, corresponding to the left stem of the RBS sequestor, and thus releases the translation initiation of *cbiA*. Moreover, it has been suggested (Ravnum and Andersson 2001) that in the presence of cobalamin, the enhancer element interacts with the *B12*-element, thus the RBS sequestor forms and repression the translation initiation is repressed.

To elucidate the mechanism of regulation, we analyzed multiple alignments of upstream regions of close orthologs of *btuB* genes from enterobacteria and some α -proteobacteria. It turned out that these alignments contained islands of conservation, corresponding to *B12*-elements, sequestor hairpins, and some additional elements in between (Fig. 4). The first additional conserved element is complementary to the left stem of the RBS sequestor. Thus, it can form an antisequestor structure whose possible role is similar to the translational enhancer. Another conserved region can form a pseudoknot by interacting with the loop region of the P3 helix of the *B12*-element. Similar additional elements in 5'UTRs of *B12*-regulated genes were found in most Gram-positive and Gram-negative bacteria (Fig. 5). In some cases the pseudoknot element overlaps the left base stem of the antisequestor/antiterminator, in other cases (including the cases shown in Fig. 4), it is located within the antisequestor/antiterminator. Thus, the predicted mechanism of the co-

B

FIGURE 4. Multiple alignment of the *btuB* upstream regions from enterobacteria (A) and some α -proteobacteria (B). Gray background denotes the P0 and P3 stems of the *B12*-element and the proposed RBS sequestors. Bold black denotes the main stem of the antisequestor. Arrows show the complementary stems of RNA secondary structures. Candidate pseudoknots are underlined and designated Pkn. RBSs and start codons are shown in bold letters.

balamin riboswitch differs from the thiamin and riboflavin riboswitches, as it involves formation of the pseudoknot (Fig. 6). The *B12*-element stabilized by Ado-CBL interacts with the antisequestor/antiterminator region, forms the pseudoknot, and promotes the sequestor formation. In the absence of Ado-CBL, the nonstabilized *B12*-element cannot form the pseudoknot and the more favorable of antisequestor/antiterminator structure folds, thus allowing for the translation initiation or transcriptional readthrough.

Phylogeny of the *B12*-elements

Global analysis of the *B12*-elements in available bacterial genomes demonstrates that this conserved RNA regulatory

FIGURE 5. Conserved RNA elements upstream of some B12-regulated genes. The P0 and P3 stems of the B12-element are highlighted in green. Proposed regulatory hairpins (sequesters in proteobacteria and actinobacteria; terminators in the *Bacillus/Clostridium* group) are highlighted in blue. Red denotes the base stem of the antiterminator/antisequencer. Arrows in the upper line show the complementary stems of RNA secondary structures. Candidate pseudoknots that overlap the P3 loop and the antiterminator/antisequencer are underlined. RBSs and the start codons are set in blue.

FIGURE 6. Predicted mechanism of the *B12*-mediated regulation of CBL genes: (A) transcriptional attenuation; (B) translational attenuation (inhibition of translation initiation).

element is widely distributed in eubacteria and regulates most genes required for the CBL biosynthesis and some other genes (Table 2; for details see D.A. Rodionov, A.G. Vitreschak, A.A. Mironov, and M.S. Gelfand, in prep.). First, various *cbi* and *cob* biosynthetic genes are regulated by *B12*-elements in most CBL-synthesizing bacteria. These

genes either form single CBL gene clusters or are scattered along the chromosome. Moreover, genes for most known and predicted cobalt transporters, as well as cobalt chelates and reductases, are often preceded by candidate *B12*-elements. Indeed, cobalt ions are required for the de novo CBL biosynthesis. Second, the vitamin *B*₁₂ transport sys-

TABLE 2. Phylogenetic distribution of gene clusters regulated by *B12*-elements

Gene cluster	Function	Taxonomic group
1. CBL biosynthesis <i>cbi</i> and <i>cob</i> <i>cbt</i> , <i>hoxN</i> , <i>cbiMNQO</i> , <i>hupE</i> <i>orfI-cobW-cobN-chlID</i> <i>bluB</i> <i>btuR</i>	cobalamin biosynthesis cobalt transporters cobalt chelation cobalt reduction CBL adenosyltransferase	proteobacteria, the <i>Bacillus/Clostridium</i> group all CBL-synthesizing bacteria α -, β -proteobacteria, <i>Pseudomonadaceae</i> , actinobacteria α -proteobacteria α -, β -proteobacteria, <i>Pseudomonadaceae</i>
2. Vitamin <i>B</i> ₁₂ transport <i>btuB</i> <i>btuFCD</i>	vitamin <i>B</i> ₁₂ receptor vitamin <i>B</i> ₁₂ transporter (ABC components)	proteobacteria α -, β -proteobacteria, <i>Pseudomonadaceae</i> , the <i>Bacillus/Clostridium</i> and CFB groups, <i>Deinococcus</i> <i>radiodurans</i> , actinobacteria, spirochetes, <i>Fusobacteriaceae</i> , <i>Thermotogales</i> , <i>Chloroflexaceae</i>
3. <i>B</i> ₁₂ -dependent or alternative metabolic pathways <i>metE</i> <i>nrd</i> <i>ardX-frdX</i> <i>achX</i>	methionin synthase ribonucleotide reductase predicted enzymes predicted enzymes	various groups various groups α -proteobacteria <i>Deinococcus radiodurans</i> and some other species

Using multiple alignment of identified *B12*-element sequences without additional nonconserved interior loops and the BII part, we constructed the maximum likelihood phylogenetic tree for these RNA elements (Fig. 7). The BII part was not used as it is not obligatory for all *B12*-elements.

The branch of *B12*-elements found upstream of the *cbi* operons of enterobacteria is the most obvious example of possible horizontal transfer, as it clusters with various *B12*-elements from the *Bacillus/Clostridium* group (Fig. 7), and the same holds for phylogenetic trees constructed for each gene from the *cbi* operon (data not shown). These observations allow us to suggest that the complete transcriptional

1094 RNA, Vol. 9, No. 9

unit, including the regulatory B_{12} -element and the *cbi* operon, has been likely transferred from the *Bacillus/Clostridium* group to three enterobacterial genomes.

The phylogenetic tree of B_{12} -elements contains a number of genome-specific branches. In particular, four B_{12} -elements found upstream of the *pccC*, *cbtD*, *nrdA*, and *metE* genes from *Bacteroides fragilis* form a separate branch (Fig. 7). Furthermore, in *D. radiodurans*, *Burkholderia pseudomallei*, *Rhodopseudomonas palustris*, *Rhodobacter capsulatus*, *Heliobacillus mobilis*, *Sphingomonas aromaticivorans*, *Streptomyces coelicolor*, *Clostridium difficile*, *Porphyromonas gingivalis*, *Chlorobium tepidum*, and *Chloroflexus aurantiacus*, we observed other organism-specific branches containing two or three B_{12} -elements. In all of these cases, the regulatory elements are located upstream of various nonhomologous genes. Thus, it seems likely that the evolution of the B_{12} -elements often involved independent lineage-specific duplications with subsequent transfer to a position upstream of a new gene. On the other hand, the phylogenetic tree has branches that correspond to B_{12} -elements occurring upstream of orthologous genes, for example, the *btuB* genes from enterobacteria, the *cobW* genes from pseudomonads, and the *bluB* and *metE* genes from α -proteobacteria. These observations indicate that these B_{12} -elements have co-evolved with the corresponding B_{12} -regulated genes.

Finally, the constructed phylogenetic tree allows us to propose a possible origin of the B_{12} -elements lacking the highly conserved BII part (see bold elements in Fig. 7). Most B_{12} -elements without the BII part, for example, all B_{12} -elements from early-diverged bacteria (*D. radiodurans*, *Chloroflexaceae*, and cyanobacteria), form a separate branch in the phylogenetic tree. The BII part seems to appear after divergence of these taxonomic groups (recall that the BII part was not used in constructing the tree). However, a B_{12} -element from *B. subtilis* lacking the BII part is clustered with several complete B_{12} -elements from other *Bacillus* species. Moreover, another BII-deficient B_{12} element from *Pseudomonas aeruginosa* clusters with other *Pseudomonas* B_{12} -elements with both BI and BII parts. The absence of BII parts in these two B_{12} -elements likely is a consequence of late deletions in these species. Thus, although the additional BII part is highly conserved in a large number of B_{12} -elements, it is possibly not obligatory and can play an auxiliary role in the functioning of the B_{12} -element.

CONCLUSIONS

Regulation of most B_{12} -related genes in bacteria appears to operate through a unique RNA structural element. The B_{12} -element is characterized by its compact secondary structure with a number of conserved helices and extended regions of sequence conservation, which could be necessary for specific metabolite binding. Recently, it has been shown that Ado-CBL specifically binds the leader mRNAs of CBL-

related genes from enterobacteria (Nahvi et al. 2002). However, the B_{12} -element also contains the highly conserved BII structure, which is not obligatory for all bacterial B_{12} -elements, being absent in deeply branching bacterial groups and in some other cases. The role of this additional part of the B_{12} -element is not clear.

On the whole, the mechanism of regulation of vitamin B_{12} -related genes is similar to the previously proposed mechanisms of regulation of riboflavin- and thiamin-related genes (Rodionov et al. 2002; Vitreschak et al. 2002). At that, a highly conserved RNA element (*RFN*, *THI*, or B_{12}) is stabilized by direct binding of the effector (flavin mononucleotide, thiamin pyrophosphate, or adenosylcobalamin, respectively). Thus, an adjacent regulatory hairpin, terminator or sequestor, can fold that leads to transcriptional or translational repression of vitamin-related genes. In the absence of the effector, the unstable vitamin-specific RNA element is replaced by an alternative antiterminator or antisequestor RNA conformation allowing for transcription readthrough or translation initiation. Interestingly, the observed transcriptional and translational types of regulation have similar taxonomic distribution for each of the analyzed vitamin regulons. At that, the transcriptional termination occurs mostly in Gram-positive organisms, whereas the inhibition of translation initiation happens in Gram-negative proteobacteria, *D. radiodurans*, the CFB group and cyanobacteria.

This study once again demonstrates the power of the simultaneous positional and phylogenetic analysis of regulatory elements and genes. This approach provides an opportunity for identification of new RNA regulatory elements in bacterial genomes and prediction of the regulation. Moreover, it uncovers the traces of evolutionary events such as horizontal transfer and lineage-specific duplication of genes and regulatory elements.

MATERIALS AND METHODS

Complete and partial sequences of bacterial genomes were downloaded from GenBank (Benson et al. 2003). Preliminary sequence data were also obtained from the www sites of the Institute for Genomic Research (<http://www.tigr.org>), the University of Oklahoma's Advanced Center for Genome Technology (<http://www.genome.ou.edu/>), the Wellcome Trust Sanger Institute (<http://www.sanger.ac.uk/>), the DOE Joint Genome Institute (<http://jgi.doe.gov>), and the ERGO Database (<http://ergo.integratedgenomics.com/ERGO>) (Overbeek et al. 2003). The genome abbreviations from the ERGO database are used throughout and listed in Table 3.

The conserved secondary structure of the B_{12} -element was derived using the RNAMultAln program (A.A. Mironov, unpubl.). This program simultaneously creates a multiple alignment and a conserved secondary structure for a set of RNA sequences. The RNA-PATTERN program (Vitreschak et al. 2001) was used to search for new B_{12} -elements in bacterial genomes. The input RNA pattern described both the RNA secondary structure and the se-

TABLE 3. Taxonomy and abbreviations for bacterial genomes used in this work

Tax	Genome	Abbreviation
α	<i>Agrobacterium tumefaciens</i>	AU
	<i>Bradyrhizobium japonicum</i>	BJA
	<i>Brucella melitensis</i>	BME
	<i>Caulobacter crescentus</i>	CO
	<i>Mesorhizobium loti</i>	MLO
	<i>Rhodobacter capsulatus</i>	RC
	<i>Rhodobacter sphaeroides</i>	RS
	<i>Rhodospseudomonas palustris</i>	RPA
	<i>Sinorhizobium meliloti</i>	SM
	<i>Sphingomonas aromaticivorans</i>	SAR
β	<i>Burkholderia pseudomallei</i>	BPS
	<i>Methylobacillus flagellatus</i>	MFL
	<i>Nitrosomonas europaea</i>	NE
	<i>Ralstonia eutropha</i>	REU
γ	<i>Ralstonia solanacearum</i>	RSO
	<i>Escherichia coli</i>	EC
	<i>Salmonella typhimurium</i>	SY
	<i>Klebsiella pneumoniae</i>	KP
	<i>Yersinia enterocolitica</i>	YE
	<i>Yersinia pestis</i>	YP
	<i>Erwinia carotovora</i>	EO
	<i>Vibrio cholerae</i>	VC
	<i>Pseudomonas aeruginosa</i>	PA
	<i>Pseudomonas putida</i>	PP
	<i>Pseudomonas fluorescens</i>	PU
	<i>Pseudomonas syringae</i>	PY
	<i>Shewanella oneidensis</i>	SON
	<i>Azotobacter vinelandii</i>	AV
δ	<i>Xanthomonas axonopodis</i>	XAX
	<i>Geobacter metallireducens</i>	GME
	<i>Chloroflexus aurantiacus</i> #	CAU
	<i>Fusobacterium nucleatum</i>	FN
B/C	<i>Thermotoga maritima</i>	TM
	<i>Bacillus subtilis</i>	BS
	<i>Bacillus cereus</i>	ZC
	<i>Bacillus megaterium</i>	BI
	<i>Bacillus halodurans</i>	HD
	<i>Bacillus stearothermophilus</i>	BE
	<i>Listeria monocytogenes</i>	LMO
	<i>Clostridium acetobutylicum</i>	CA
	<i>Clostridium perfringens</i>	CPE
	<i>Clostridium botulinum</i>	CB
	<i>Clostridium difficile</i>	DF
	<i>Thermoanaerobacter tengcongensis</i>	THT
	<i>Enterococcus faecalis</i>	EF
	<i>Heliobacillus mobilis</i>	HMO
Act	<i>Desulfotobacterium halfniense</i>	DHA
	<i>Corynebacterium diphtheriae</i>	DI
	<i>Mycobacterium tuberculosis</i>	MT
	<i>Mycobacterium leprae</i>	ML
Cya	<i>Thermobifida fusca</i>	TFU
	<i>Rhodococcus str.</i>	RK
	<i>Streptomyces coelicolor</i>	SX
	<i>Propionibacterium shermanii</i>	PI
	<i>Anabaena</i> sp.	AN
	<i>Prochlorococcus marinus</i>	PMA
	<i>Synechocystis</i> sp.	CY
	<i>Synechococcus</i> sp.	SN
	<i>Thermosynechococcus elongatus</i>	TEL

(continued)

TABLE 3. Continued

Tax	Genome	Abbreviation
CFB	<i>Porphyromonas gingivalis</i>	PG
	<i>Bacteroides fragilis</i>	BX
	<i>Chlorobium tepidum</i>	CL
T/D	<i>Deinococcus radiodurans</i>	DR
SP	<i>Treponema denticola</i>	TDE
	<i>Leptospira interrogans</i>	LI

The names of taxonomic groups in Tax column (α , β , γ , δ , B/C, Act, Cya, CFB, T/D and SP) stand for α -, β -, γ -, and δ -proteobacteria, the *Bacillus/Clostridium* group, actinobacteria, cyanobacteria, the CFB group, the *Thermus/Deinococcus* group and spirochetes, respectively.

quence consensus motifs. The RNA secondary structure was described as a set of the following parameters: the number of helices, the length of each helix, the loop lengths, and the description of the topology of helix pairs. Additional RNA secondary structures, in particular antiterminators and antisequestors, were predicted using Zuker's algorithm of free energy minimization (Lyngso et al. 1999) implemented in the Mfold program (<http://bioinfo.math.rpi.edu/~mfold/rna>).

Protein similarity search was done using the Smith-Waterman algorithm implemented in the GenomeExplorer program (Mironov et al. 2000). Orthologous proteins were initially defined by the best bidirectional hit criterion (Tatusov et al. 2000) and if necessary, confirmed by construction of phylogenetic trees. The phylogenetic trees of the *B12*-elements and *B₁₂*-related proteins were created by the maximum likelihood method implemented in PHYLIP (Felsenstein 1981). Multiple sequence alignments were constructed using CLUSTALX (Thompson et al. 1997).

Complete FASTA sequences of *B12*-elements, including the additional and variable regions are available from the authors (L.V. at l_veter@mail.ru or M.G. at gelfand@ig-msk.ru).

ACKNOWLEDGMENTS

We are grateful to Andrei Osterman for attention, advice, and encouragement. This study was partially supported by grants from the Ludwig Institute for Cancer Research and the Howard Hughes Medical Institute (55000309).

The publication costs of this article were defrayed in part by payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 USC section 1734 solely to indicate this fact.

Received April 16, 2003; accepted June 3, 2003.

REFERENCES

- Benson, D.A., Karsch-Mizrachi, I., Lipman, D.J., Ostell, J., and Wheeler, D.I. 2003. GenBank. *Nucleic Acids Res.* **31**: 23–27.
- Dandekar, T., Beyer, K., Bork, P., Kenealy, M.R., Pantopoulos, K., Hentze, M., Sonntag-Buck, V., Flouriot, G., Gannon, F., and Schreiber, S. 1998. Systematic genomic screening and analysis of mRNA in untranslated regions and mRNA precursors: Combining experimental and computational approaches. *Bioinformatics* **14**: 271–278.

- Eddy, S.R. and Durbin, R. 1994. RNA sequence analysis using covariance models. *Nucleic Acids Res.* **22**: 2079–2088.
- Felsenstein, J. 1981. Evolutionary trees from DNA sequences: A maximum likelihood approach. *J. Mol. Evol.* **17**: 368–376.
- Franklund, C.V. and Kadner, R.J. 1997. Multiple transcribed elements control expression of the *Escherichia coli* *btuB* gene. *J. Bacteriol.* **179**: 4039–4042.
- Gelfand, M.S., Novichkov, P.S., Novichkova, E.S., and Mironov, A.A. 2000. Comparative analysis of regulatory patterns in bacterial genomes. *Brief Bioinform.* **1**: 357–371.
- Griffiths-Jones, S., Bateman, A., Marshall, M., Khanna, A., and Eddy, S.R. 2003. Rfam: An RNA family database. *Nucleic Acids Res.* **31**: 439–441.
- Grundy, F.J. and Henkin, T.M. 1998. The S box regulon: A new global transcription termination control system for methionine and cysteine biosynthesis genes in gram-positive bacteria. *Mol. Microbiol.* **30**: 737–749.
- Lundrigan, M.D., Koster, W., and Kadner, R.J. 1991. Transcribed sequences of the *Escherichia coli* *btuB* gene control its expression and regulation by vitamin B_{12} . *Proc. Natl. Acad. Sci.* **88**: 1479–1483.
- Lyngso, R.B., Zuker, M., and Pedersen, C.N. 1999. Fast evaluation of internal loops in RNA secondary structure prediction. *Bioinformatics* **15**: 440–445.
- Marck, C. and Grosjean, H. 2002. tRNomics: Analysis of tRNA genes from 50 genomes of Eukarya, Archaea, and Bacteria reveals anticodon-sparing strategies and domain-specific features. *RNA* **8**: 1189–1232.
- Martens, J.H., Barg, H., Warren, M.J., and Jahn, D. 2002. Microbial production of vitamin B_{12} . *Appl. Microbiol. Biotechnol.* **58**: 275–285.
- Mironov, A.A., Vinokurova, N.P., and Gelfand, M.S. 2000. Genome-Explorer: Software for analysis of complete bacterial genomes. *Mol. Biol.* **34**: 222–231.
- Mironov, A.S., Gusarov, I., Rafikov, R., Lopez, L.E., Shatalin, K., Krenova, R.A., Perumov, D.A., and Nudler, E. 2002. Sensing small molecules by nascent RNA: A mechanism to control transcription in bacteria. *Cell* **111**: 747–756.
- Murthy, V.L. and Rose, G.D. 2003. RNABase: An annotated database of RNA structures. *Nucleic Acids Res.* **31**: 502–504.
- Nahvi, A., Sudarsan, N., Ebert, M.S., Zou, X., Brown, K.L., and Breaker, R.R. 2002. Genetic control by a metabolite binding mRNA. *Chem. Biol.* **9**: 1043–1049.
- Nou, X. and Kadner, R.J. 1998. Coupled changes in translation and transcription during cobalamin-dependent regulation of *btuB* expression in *Escherichia coli*. *J. Bacteriol.* **180**: 6719–6728.
- . 2000. Adenosylcobalamin inhibits ribosome binding to *btuB* RNA. *Proc. Natl. Acad. Sci.* **97**: 7190–7195.
- Overbeek, R., Larsen, N., Walunas, T., D'Souza, M., Pusch, G., Selkov Jr., E., Liolios, K., Joukov, V., Kaznadzey, D., Anderson, I., et al. 2003. The ERGO(TM) genome analysis and discovery system. *Nucleic Acids Res.* **31**: 164–171.
- Ravnum, S. and Andersson, D.I. 1997. Vitamin B_{12} repression of the *btuB* gene in *Salmonella typhimurium* is mediated via a translational control which requires leader and coding sequences. *Mol. Microbiol.* **23**: 35–42.
- . 2001. An adenosyl-cobalamin (coenzyme- B_{12})-repressed translational enhancer in the *cob* mRNA of *Salmonella typhimurium*. *Mol. Microbiol.* **39**: 1585–1594.
- Richter-Dahlfors, A.A. and Andersson, D.I. 1992. Cobalamin (vitamin B_{12}) repression of the *cob* operon in *Salmonella typhimurium* requires sequences within the leader and the first translated open reading frame. *Mol. Microbiol.* **6**: 743–749.
- Richter-Dahlfors, A.A., Ravnum, S., and Andersson D.I. 1994. Vitamin B_{12} repression of the *cob* operon in *Salmonella typhimurium*: Translational control of the *cbiA* gene. *Mol. Microbiol.* **13**: 541–553.
- Rodionov, D.A., Vitreschak, A.G., Mironov, A.A., and Gelfand, M.S. 2002. Comparative genomics of thiamin biosynthesis in prokaryotes: New genes and regulatory mechanisms. *J. Biol. Chem.* **277**: 48949–48959.
- Szymanski, M., Erdmann, V.A., and Barciszewski, J. 2003. Noncoding regulatory RNAs database. *Nucleic Acids Res.* **31**: 429–431.
- Tatusov, R.L., Galperin, M.Y., Natale, D.A., and Koonin, E.V. 2000. The COG database: A tool for genome-scale analysis of protein functions and evolution. *Nucleic Acids Res.* **28**: 33–36.
- Thompson, J.D., Gibson, T.J., Plewniak, F., Jeanmougin, F., and Higgins, D.G. 1997. The CLUSTAL_X windows interface: Flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res.* **25**: 4876–4882.
- Vitreschak, A.G., Mironov, A.A., and Gelfand, M.S. 2001. The RNAPattern program: Searching for RNA secondary structure by the pattern rule. In *Proceedings of the 3rd International Conference on "Complex Systems: Control and Modeling Problems."*, September 4–9, 2001, pp. 623–625. The Institute of Control of Complex Systems, Samara, Russia.
- Vitreschak, A.G., Rodionov, D.A., Mironov, A.A., and Gelfand, M.S. 2002. Regulation of riboflavin biosynthesis and transport genes in bacteria by transcriptional and translational attenuation. *Nucleic Acids Res.* **30**: 3141–3151.
- Wolf, Y.I., Rogozin, I.B., Grishin, N.V., Tatusov, R.L., and Koonin, E.V. 2001. Genome trees constructed using five different approaches suggest new major bacterial clades. *BMC Evol. Biol.* **1**: 8.
- Winkler, W., Nahvi, A., and Breaker, R.R. 2002a. Thiamine derivatives bind messenger RNAs directly to regulate bacterial gene expression. *Nature* **419**: 952–956.
- Winkler, W., Cohen-Chalamish, S., and Breaker, R.R. 2002b. An mRNA structure that controls gene expression by binding FMN. *Proc. Natl. Acad. Sci.* **99**: 15908–15913.

RNA

A PUBLICATION OF THE RNA SOCIETY

Regulation of the vitamin B₁₂ metabolism and transport in bacteria by a conserved RNA structural element

ALEXEY G. VITRESCHAK, DMITRY A. RODIONOV, ANDREY A. MIRONOV, et al.

RNA 2003 9: 1084-1097

References

This article cites 31 articles, 7 of which can be accessed free at:
<http://rnajournal.cshlp.org/content/9/9/1084.full.html#ref-list-1>

License

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).

To subscribe to *RNA* go to:
<http://rnajournal.cshlp.org/subscriptions>