Effects of transgene insertion loci and copy number on *Dnmt3L* gene silencing through antisense transgene-derived PIWI-interacting RNAs

SePil Lee¹, Satomi Kuramochi-Miyagawa^{1,2*}, Ippei Nagamori², and Toru Nakano^{1,2*}

¹ Graduate School of Frontier Biosciences and ² Medical School, Department of Pathology, Osaka University, Yamada-oka 2-2 Suita, Osaka 565-0871, Japan

*Address correspondence:

Satomi Kuramochi-Miyagawa (PhD, E-mail: smiya@patho.med.osaka-u.ac.jp)

Toru Nakano (MD, PhD, E-mail: tnakano@patho.med.osaka-u.ac.jp)

(Tel: +81-6-6879-3721, Fax: +81-6-6879-3729)

Running title: Silencing by transgene-derived piRNA

Keywords:

piRNA, PIWI, transcriptional gene silencing (TGS), post-transcriptional gene silencing (PTGS),

germ cell

Abstract

PIWI-interacting RNAs (piRNAs), which are germ cell-specific small RNAs, are essential for spermatogenesis. In fetal mouse germ cells, piRNAs are synthesized from sense and antisense RNAs of transposable element sequences for retrotransposon silencing. In a previous study, we reported that transgenic mice expressing antisense-*Dnmt3L* under the control of the *Miwi2* promoter (Tg-*Miwi2P*-as*Dnmt3L*) exhibited piRNA-mediated DNMT3L downregulation. In this study, two transgene integration loci (B3 and E1) were identified on chromosome 18 of the Tg-*Miwi2P*-as*Dnmt3L* mice; these loci were weak piRNA clusters. Crossbreeding was performed to obtain mice with the transgene cassette inserted into a single locus. DNMT3L was silenced and spermatogenesis was severely impaired in mice with the transgene cassette inserted at the B3 locus (Tg-B mice). In contrast, spermatogenesis in mice bearing the transgene at the E1 locus (Tg-E mice) was normal. The number of piRNAs for *Dnmt3L* in Tg-B mice was eight-fold higher than that in Tg-E mice. Therefore, both gene silencing and impaired spermatogenesis depended on the transgene copy number rather than on the insertion loci. Additionally, the endogenous *Dnmt3L* promoter was not methylated in Tg mice, suggesting that *Dnmt3L* silencing was caused by post-transcriptional gene silencing. Based on these data, we discuss a piRNA-dependent gene silencing mechanism against novel gene insertions.

1 Introduction

2 PIWI-interacting RNAs (piRNAs), a class of germ cell-specific small RNAs, regulate
3 spermatogenesis and maintain genome integrity by repressing transposons (Castaneda et al. 2011;
4 Chuma and Nakano 2013; Toth et al. 2016). Approximately half of the human and mouse
5 genomes comprise transposons and transposon fragments as part of their repetitive sequences
6 (Mouse Genome Sequencing et al. 2002). The transcription of transposons is repressed through
7 piRNA-dependent DNA methylation and repressive histone modifications that play crucial roles
8 in mouse spermatogenesis (Aravin et al. 2007, 2008; Carmell et al. 2007; Kuramochi-Miyagawa
9 et al. 2008; De Fazio et al. 2011; Reuter et al. 2011; Seisenberger et al. 2012; Zheng and Wang
10 2012; Di Giacomo et al. 2013; Pezic et al. 2014). In addition to their role in epigenetic gene
11 silencing, piRNAs mediate the RNA degradation of target transposons. piRNA-PIWI family
12 protein complexes mediate transcriptional gene silencing (TGS) and post-transcriptional gene
13 silencing (PTGS) (Senti et al. 2015; Czech and Hannon 2016).

14 There are three members of the mouse PIWI family of proteins, namely, mouse Piwi (MIWI),
15 MIWI-like (MILI), and MIWI2, all of which are required for spermatogenesis (Kuramochi-
16 Miyagawa et al. 2001, 2004; Deng and Lin 2002; Carmell et al. 2007). In fetal male germ cells,
17 the vast majority of piRNAs are derived from retrotransposon genes, and two PIWI family
18 proteins, MILI and MIWI2, are involved in this process. Fetal piRNAs are mainly generated from
19 transcripts of piRNA-producing regions called piRNA clusters, although the molecular
20 mechanisms underlying piRNA clusters are currently unknown. The piRNA clusters expressed in
21 the fetal testis are genomic regions that accumulate loss-of-function transposon fragments or full-
22 length transposons (Girard et al. 2006; Aravin et al. 2007; Brennecke et al. 2007; Zanni et al.
23 2013). Primary piRNAs, which exhibit a sequence bias for uridine at the first position of the 5'
24 end (1U bias), were produced and incorporated into MILI. Next, piRNA-bound MILI, which

functions as an endonuclease (slicer activity), cleaves the complementary target RNAs annealed to the primary piRNAs between nucleotides 10 and 11 (Aravin et al. 2007, 2008). The secondary piRNA produced from MILI cleavage exhibits a sequence bias for adenine at the tenth position (10A bias) and is further incorporated into MILI or MIWI2 (Aravin et al. 2008; De Fazio et al. 2011). piRNA-bound MILI represses transposon expression in the cytoplasm through the cleavage of transposon RNAs. In contrast, piRNA-bound MIWI2 migrates to the nucleus, where piRNAs guide binding to nascent RNAs and induce DNA methylation of retrotransposon genes, such as *LINE* and *IAP* (Carmell et al. 2007; Molaro et al. 2014; Manakov et al. 2015).

DNA methylation is one of the gene expression control mechanisms that regulates cell differentiation and development. Germ cells undergo global epigenetic reprogramming, including DNA methylation. In mice, DNA methylation is globally erased by embryonic day 12.5 (E12.5) during the development of germ cells; consequently, parental epigenetic imprinting is erased (Hajkova et al. 2002; Sasaki and Matsui 2008). Subsequently, global *de novo* DNA methylation occurs in approximately 76% of the genome of mouse gonocytes, the male germ cells, around E16.5 (Molaro et al. 2014). DNMT3L and DNMT3A, which are involved in *de novo* DNA methylation, target transposable elements and paternally imprinted genes (*H19*, *Dlk1*, and *Rasgrf1*) (Li et al. 2004; Kato et al. 2007; La Salle et al. 2007; Watanabe et al. 2011). Notably, as DNA methylation of retrotransposon genes is dependent on piRNAs, a similar phenotype is observed in MILI-deficient mice, in which almost no piRNAs are produced.

Previously, we demonstrated that the expression of sense and antisense RNAs in fetal male germ cells is sufficient for piRNA production and subsequent *de novo* DNA methylation (Itou et al. 2015). We previously generated transgenic (Tg) mice harboring the antisense-*EGFP* gene under the control of the *Miwi2* promoter (Tg-*Miwi2P*-as*EGFP*). *Miwi2* is expressed only in the male fetal germ cells. Crossing Tg-*Miwi2P*-as*EGFP* mice with *EGFP*-expressing Tg mice

1 resulted in EGFP silencing in the progeny. Next, Tg mice expressing antisense-*Dnmt3L* under the
2 control of the *Miwi2* promoter (Tg-*Miwi2P*-as*Dnmt3L*) were generated. Tg-*Miwi2P*-as*Dnmt3L*
3 mice exhibited downregulated endogenous *Dnmt3L* RNA expression and a phenotype similar to
4 that of DNMT3L-deficient mice. In these Tg mice, piRNAs against *EGFP* and *Dnmt3L* were
5 produced, and DNA methylation occurred in their corresponding genomic loci. However, the
6 piRNAs of these Tg mice were mapped not only to the target genes *EGFP* and *Dnmt3L*, but also
7 to *Miwi2* promoter regions. These data indicated that the transcription of the piRNA precursors
8 against as*EGFP* and as*Dnmt3L* was not only regulated by the *Miwi2* promoter but also by some
9 unknown promoters outside of the inserted Tg cassette. It is reasonable to assume that these
10 transgenes have been incorporated into piRNA clusters (Shoji and Katsuma 2015; Kuramochi-
11 Miyagawa and Nakano 2015).

12 In this study, whole-genome sequencing of Tg-*Miwi2P*-as*Dnmt3L* mice led to the
13 identification of two insertion sites: the B3 and E1 loci on chromosome 18 (ch18). The analysis
14 of the two loci revealed that each of them had different roles in piRNA production and,
15 consequently, in mouse phenotypes. The E1 locus, which is presumably a weak piRNA cluster,
16 produced some piRNAs. However, the mice harboring the transgene at the E1 locus were fertile.
17 In contrast, the mice harboring the transgene at the B3 locus (a very weak piRNA cluster)
18 exhibited infertility due to the post-transcriptional silencing of *Dnmt3L*. Further analyses revealed
19 that piRNA-mediated *Dnmt3L* silencing was dependent on the transgene copy number rather than
20 on the transgene insertion locus.

21

Results

Miwi2P-asDnmt3L transgene insertion locus in the mouse genome

Whole-genome DNA sequencing of Tg-*Miwi2P-asDnmt3L* mice was performed to examine whether the Tg cassette insertion loci corresponded to piRNA clusters. Four junction sites between the transgene and chromosome (two at the B3 locus and two at the E1 locus on ch18) were identified (Fig. 1A). *Dnmt3L*-polyA ($\Delta 604$) - 41273991 (NCBI/mm10) and *Miwi2P* - 41311357 (NCBI/mm10) were detected at the B3 locus, while the polyA - 61646220 (NCBI/mm10) and *Miwi2P* - 61648546 (NCBI/mm10) junctions were detected at the E1 locus (Fig. 1A).

Transgene insertions did not reduce the mapping reads by half in any of the two genomic areas. The DNA sequencing read counts around the junction sites of the transgene increased by 1.5-fold in both loci compared to other regions of the same chromosome (Fig. 1B–C). These findings indicate that the 37 kb and 2.3 kb regions of the 18qB3 and 18qE1 loci, respectively, underwent duplication. Polymerase chain reaction (PCR) was performed using primers designed to flank the 2.3 kb region (Supplemental Fig. S1A). The amplified sequences confirmed duplication (Supplemental Fig. S1B). The duplication of the B3 locus could not be confirmed due to the length of the duplicated region. Therefore, similar to E1, B3 was assumed to have undergone duplication.

piRNAs of the flanking regions of transgene insertion sites

To determine whether these two loci were part of any piRNA cluster, previously reported data on MILI-associated and MIWI2-associated piRNAs of E16.5 wild-type (WT) and MILI^{DAH} (MILI endonuclease-impaired mutant) mice in the piRNA database (piRBase: <http://regulatoryrna.org/database/piRNA/index.html>) were analyzed (De Fazio et al. 2011; Zhang

et al. 2014). The expression level of primary piRNAs in MILI^{DAH} mice was higher than that in WT mice because mutant mice cannot produce secondary piRNAs. Some MILI- and MIWI2-bound piRNAs were detected around the E1 locus in both WT and MILI^{DAH} mice (Supplemental Fig. S1C). In contrast, only a small number of piRNAs were annotated around the B3 locus. The number of piRNAs or coverage, and the average values of each category in the 50 kb regions surrounding the E1 and B3 loci are shown in Supplemental Figure S1C.

Next, these two regions were analyzed using piRNAQuest (<http://bicresources.jcbose.ac.in/zhumur/pirnaquest/index.html>), which annotates piRNA clusters (Sarkar et al. 2014). Each of the two regions was included in a piRNA cluster, although both cluster scores were low (Supplemental Table S1). These data suggest that the E1 and B3 loci are weak piRNA clusters.

Some sense and antisense piRNAs were detected in both 50 kb regions surrounding the insertion sites in WT mice (Supplemental Fig. S1C–D). Therefore, the piRNAs (23–30 nt) in the 2.3 kb duplicated region of the E1 locus and the 2.3 kb regions adjacent to the insertion sites of the B3 locus (B3-5' and B3-3') were analyzed (Supplemental Fig. S1D). The number of each piRNA was normalized based on the length and number of total small RNAs per million after excluding miRNAs and tRNAs. The 2.5 kb of the *Miwi2* promoter used for the transgene was not a piRNA cluster region, considering that very few piRNAs were mapped to this region; therefore, it was used as the negative control. In contrast, the number of antisense piRNAs was high in the E1 and B3-5' regions, whereas that of sense-strand piRNAs was low. These findings indicate that the insertion sites have the potential to produce antisense piRNAs, and that antisense piRNAs are transcribed in this region. Thus, both regions were considered weak piRNA clusters.

Phenotypes of Tg mice with single transgene insertions

1 To determine which locus was critical for silencing *Dnmt3L*, crossbreeding was performed to
2 segregate each insertion into different mice of the progeny. Segregation was confirmed by PCR
3 (Fig. 2A). Mice with the transgene cassette inserted at the B3 locus (Tg-B) exhibited lower testis
4 size and weight relative to body weight than the control mice. In contrast, mice with the cassette
5 at the E1 locus (Tg-E) showed a normal phenotype (Fig. 2B–C).

6 Western blot analysis revealed that the expression of endogenous DNMT3L was
7 downregulated in the testes of Tg-B mice at E16.5, and in Tg mice with transgenes inserted at
8 both E1 and B3 loci (Tg-BE; Tg-*Miwi2P-asDnmt3L*). However, the expression level of DNMT3L
9 in Tg-E mice was similar to that in the control mice (Fig. 2D). Hematoxylin and eosin (HE)
10 staining revealed severe spermatogenesis defects in the testes of Tg-B and Tg-BE mice. In
11 contrast, the testes of Tg-E mice did not exhibit any histological defects (Fig. 2E). These findings
12 suggest that the phenotype of Tg-*Miwi2P-asDnmt3L* mice is only due to the insertion in the B3
13 locus. We then examined the meiotic chromosome spreads and stained them with antibodies
14 against SCP1 and SCP3 to evaluate meiotic defects in DNMT3L mutants. We found that most
15 meiotic cells in Tg-B mice were at the zygotene or zygotene-like stages, similar to the phenotype
16 of DNMT3L-deficient mice.

17 Next, bisulfite sequencing of the target genes of DNMT3L, such as retrotransposons and
18 imprinted genes, was performed. Germ cells were sorted based on the epithelial adhesion
19 molecule Ep-CAM, a spermatogonial stem cell marker, at postnatal day 10 (P10) (Kanatsu-
20 Shinohara et al. 2011). We found that retrotransposon genes (*LINE1* type A, *LINE1* Tf family,
21 and LTR element *L1PIΔI*) and paternally imprinted genes (*H19*, *Dkl1*, and *Rasgrfl*) were
22 methylated in the germ cells of non-Tg mice (Fig. 3A–B). However, consistent with our previous
23 study, retrotransposon genes and *Rasgrfl* in the germ cells of Tg-BE mice exhibited decreased
24 methylation levels. Similarly, germ cells of Tg-B mice exhibited decreased levels of DNA

1 methylation. Although Tg-E mice did not exhibit any histological defects at the different stages
2 of spermatogenesis, their germ cells exhibited partial methylation (Fig. 3A–B).

3 We performed northern blotting to evaluate the transcription levels of the transposable
4 elements *LINE1* and *IAP* and their correlation with the bisulfite sequencing data. As shown in
5 Figure 3D, both *LINE1* and *IAP* were highly expressed in Tg-B and Tg-BE cells. In contrast, Tg-
6 E showed a lower expression. This indicates that decreased DNA methylation and increased
7 transcription levels of DNMT3L target genes in Tg-*Miwi2P-asDnmt3L* (Tg-BE) mice were
8 mainly caused by the insertion in B3. However, the phenotypic differences between Tg-B and
9 Tg-E mice could not be explained by the insertion site of the transgene, given that both insertion
10 loci were weak piRNA clusters. Therefore, transgene copy numbers were examined.

12 ***Copy number of the antisense-Dnmt3L transgene and its correlation with phenotype***

13 Genomic quantitative PCR (qPCR) analysis revealed that Tg-E mice harbored approximately
14 10 copies of the transgene cassette. In contrast, 11 of the 12 Tg-B mice harbored 28–51 copies,
15 and 1 mouse harbored only 9 copies. The copy number of the transgene cassette in Tg-BE mice
16 varied from 16 to 61 (Fig. 4A). The reasons for these differences are unclear. One possible
17 explanation is the occurrence of recombination events among tandemly arrayed transgenes,
18 similar to the recombination observed in SINE and LINE transposons.

19 To determine the correlation between copy number and phenotype, HE staining was
20 performed, and the proportion of cells arrested at meiosis and those that had completed it in the
21 seminiferous tubules was examined (Figs. 4B–C; Supplemental Fig. S2A–B). Tg-E mice
22 exhibited normal spermatogenic progression. In contrast, most Tg-B mice exhibited severe
23 spermatogenic defects; the exception was the mouse with a low copy number (that with nine
24 copies), which exhibited normal spermatogenesis. In Tg-BE mice, the severity of spermatogenic

1 defects was almost directly proportional to the transgene copy number. These findings
2 demonstrate that high transgene copy numbers are correlated with the degree of spermatogenic
3 defects.

4 Next, the phenotypes of Tg-B mice with low transgene copy numbers and Tg-E mice with
5 high transgene copy numbers were examined. Crossing female Tg-B mice with WT male mice
6 resulted in mice harboring various copies of the transgene. However, Tg-B mice with transgene
7 copy numbers below 40 were not obtained. Crossing male and female Tg-E mice resulted in mice
8 with 2-fold the number of transgenes on both alleles of the E locus (Tg-E/E mice). In Tg-E/E
9 mice, 30%–50% of the cells in the seminiferous tubules were arrested during meiosis (Fig. 4D;
10 Supplemental Fig. S2C). Additionally, testis size and weight relative to body weight were
11 significantly decreased in Tg-E/E mice (Fig. 4E; Supplemental Fig. S2C). The correlation
12 between transgene copy number and the progression of spermatogenesis or testis weight is shown
13 in Figure 4F–G and Supplemental Figure S2D. It was observed that 10 copies of the transgene
14 did not affect spermatogenesis, whereas more than 25 completely inhibited it. Interestingly, Tg
15 mice harboring approximately 20 copies of the transgene exhibited partial inhibition of
16 spermatogenesis. These data suggest that the phenotype is dependent on the copy number of the
17 transgene rather than on the transgene insertion locus.

18 19 *piRNA expression in the fetal testes of Tg-BE, Tg-B, and Tg-E mice.*

20 Next, the expression of piRNAs against *Dnmt3L* was examined. Small RNAs isolated from
21 the testes at E16.5, were subjected to deep sequencing analysis. Sequences with sizes ranging
22 from 15 to 45 nt were selected. After removing both miRNAs and tRNAs, transgene-annotated
23 small RNAs with up to two mismatches were selected (Supplemental Fig. S3A). Tg-BE and Tg-
24 B mice exhibited similar numbers of transgene-annotated piRNAs. However, the number of

1 piRNAs in Tg-E mice was approximately eight-fold lower than that in Tg-B mice (Fig. 5A;
2 Supplemental Table S2). These data suggest that piRNA copy number determines the silencing
3 of DNMT3L and, consequently, the phenotype.

4 Analysis of the first and tenth nucleotides of the transgene-annotated small RNAs from Tg
5 mice revealed that the 1U bias in the transgene sense strand was more pronounced than that in the
6 antisense strand. The antisense strand exhibited a 10A bias (Supplemental Fig. S3B). The length
7 distribution and proportion of 1U in the sense strand and of 10A in the antisense strand were
8 similar in both Tg-B and Tg-E mice (Fig. 5B). The analysis of small RNAs with 1U bias revealed
9 a prominent peak of 27 nt in the sense strand but not in the antisense strand.

10 Figure 5C–D shows the proportion of piRNAs derived from the sense transcripts of the *Miwi2*
11 promoter (light blue) and as*Dnmt3L* (blue) and from the antisense transcripts of the *Miwi2*
12 promoter (yellow) and sense *Dnmt3L* (orange). piRNAs corresponding to the first three types of
13 transcripts derived from the transgene, given that the expression of these piRNAs was extremely
14 low in control mice testes. However, it was impossible to distinguish between piRNAs from the
15 sense transcript of the endogenous *Dnmt3L* gene and those from the antisense transcript of the
16 as*Dnmt3L* transgene. Histograms showing the piRNA levels of the transgene sequence are shown
17 in Figure 5E (Tg-B) and Supplemental Figure S4F (piRNA data from Tg-E, Tg-BE, and previous
18 publications). The number of piRNAs for antisense *Dnmt3L* was higher than that for the *Miwi2*
19 promoter. Hence, most piRNAs of antisense *Dnmt3L* should derive from transcripts driven by the
20 *Miwi2* promoter.

21 Previously, we reported that endogenous *Dnmt3L* silencing can be attributed to piRNA-
22 mediated DNA methylation. Hence, DNA methylation of the promoter region of *Dnmt3L* was
23 examined using bisulfite sequencing. Methylation was not observed in any of the Tg-E, Tg-B, or
24 Tg-BE lines (Fig. 6). However, as shown in Supplemental Figure S3C, the nuclear localization of

- 1 MIWI2 was the same in the control and Tg-B mice. These data showed that MIWI2-mediated
- 2 silencing was not affected in Tg-B mice.
- 3

Discussion

Previously, we reported that the expression of antisense transcripts from a *Miwi2P-asDnmt3L* transgene promoted piRNA-mediated endogenous gene silencing in fetal male germ cells (Itou et al. 2015). However, it was pointed out that the Tg cassette was incorporated into piRNA clusters, and its expression was affected by the insertion loci (Shoji and Katsuma 2015). In this study, whole-genome sequencing and piRNA next-generation sequencing analyses revealed that, although the transgenes were incorporated into weak piRNA clusters, the copy numbers of the inserted transgenes were more important for silencing than the insertion loci.

Transgene insertion sites, copy numbers, and silencing

We defined that the insertion sites of the transgene were weak piRNA clusters because piRNAs derived from the neighboring region of the insertion sites were relatively abundant in the wild-type mouse (Supplemental Fig. S1C-D). Moreover, the whole genome DNA sequencing data indicated that the majority of the transgenes were inserted in a tandem orientation, and only 1% showed head-to-head or tail-to-tail, namely, inverted orientation. PCR analysis failed to detect inverted orientation insertion in our hands, showing that there should be few inverted junctions. Inverted orientation insertions can induce the expression of antisense transcripts, but tandem orientation insertions do not originate from molecular mechanisms. Therefore, we hypothesized that the antisense strand of the transgene was transcribed as a piRNA precursor, albeit in a small amount, by an unknown promoter outside of the inserted Tg. This supports the idea that the insertion regions are weak clusters. However, phenotypic differences between Tg-B and Tg-E mice could not be explained by the insertion site of the transgene, but rather depended on the copy number. Specifically, the copy number of the transgene determines the amount of piRNAs produced and the subsequent piRNA-mediated silencing.

1 The transgene was inserted into two loci on ch18 of Tg-*Miwi2P*-as*Dnmt3L* mice. Therefore,
2 crossbreeding was performed to obtain mice with a transgene inserted in only one of the two loci.
3 We found that different mice exhibited different transgene copy numbers. However, no transgenic
4 insertion sites containing more than 60 copies were observed. Therefore, the number of transgenes
5 in these sites either had some limitations or tended to decrease. Moreover, although the copy
6 number of Tg-E mice was stable, those of Tg-B and Tg-BE mice showed some tendency to change.
7 These results suggest that a high copy number of the transgene was not appropriately maintained.
8 After several passages, the copy number was set to approximately 40, maybe due to structural
9 stability issues.

10 Further analyses revealed that the silencing of *Dnmt3L* was dependent on the copy number of
11 the transgene and not on the insertion site (Figs. 4F–G; Supplemental Fig. S2D). The high copy
12 number of the transgenes may increase the transcription of as*Dnmt3L* under the control of the
13 *Miwi2* promoter, leading to enhanced expression of piRNAs with an as*Dnmt3L* sequence and,
14 consequently, to the silencing of the endogenous *Dnmt3L*. The findings of this study indicate that
15 a transgene copy number below 10 is not sufficient to silence DNMT3L expression and arrest
16 spermatogenesis. The silencing of *Dnmt3L* is determined by the balance between the expression
17 levels of endogenous genes and the number of piRNAs.

18 In addition to the possibility that the transgene was inserted into a piRNA cluster, pre-existing
19 genome defense mechanisms (either from the piRNA cluster or from other genomic locations)
20 may have been activated in response to the insertion. High-copy number tandem repeats of the
21 transgenes may have been recognized as novel piRNA clusters. Nevertheless, the high copy
22 number of the inserted transgene of the antisense strand is important for gene silencing in the fetal
23 testis.

1 ***Timing of DNMT3L suppression***

2 *Dnmt3L* expression was silenced, and the expression of DNMT3L protein was downregulated
3 at E16.5, in Tg-B and Tg-BE mice (Fig. 2D). Consequently, the methylation levels of DNMT3L
4 targets, such as retrotransposons and *Rasgrfl*, were diminished (Fig. 3). However, the DNA
5 methylation levels of *H19* and *Dlk1*, which are also DNMT3L targets, were unaffected. piRNA-
6 dependent DNA methylation is slightly delayed compared to piRNA-independent DNA
7 methylation, such as that of *H19* and *Dlk1* (Molaro et al. 2014). Therefore, piRNA-dependent
8 silencing of *Dnmt3L* in Tg-*Miwi2P-asDnmt3L* mice may occur after the DNA methylation of *H19*
9 and *Dlk1*. Meanwhile, DNA methylation of *LINE1* is delayed when compared with that of *IAP*
10 (Molaro et al. 2014; Kuramochi-Miyagawa et al. 2008). In addition, we observed that the
11 reduction in DNA methylation of *IAP* was not as severe as that of the *LINE1* elements *L1A* and
12 *L1Tf* (Fig. 3A), which could be explained by the DNA methylation timing differences between
13 *LINE1* and *IAP*.

14 DNA methylation of DNMT3L target genes in Tg-E mice, which did not exhibit any
15 histological defects, was partially impaired at P10. The endogenous *Dnmt3L* in the testis at E16.5,
16 was partially silenced because the number of piRNAs for antisense *Dnmt3L* was low. However,
17 threshold DNA methylation levels of DNMT3L target genes may be required to impair
18 spermatogenesis.

20 ***Interpretation of the discrepancies between studies***

21 The methylation level of the endogenous *Dnmt3L* promoter and the piRNA next-generation
22 sequencing data reported in this study were not consistent with those of a previous study (Itou et
23 al. 2015) (Supplemental Table S3). In that study, the endogenous *Dnmt3L* promoter was
24 methylated, whereas methylation was not observed here. However, gene silencing was observed

1 in both studies. This could be attributed to DNA methylation-mediated silencing in the previous
2 study and only to PTGS in this study.

3 In this study, the percentage of antisense *Dnmt3L* piRNAs with 1U bias was high (80%), and
4 the peak length of the piRNAs was 27 nt, which suggested that the antisense *Dnmt3L* piRNAs
5 were primary piRNAs bound to MILI (Fig. 5B). The endogenous *Dnmt3L* RNAs may have been
6 targeted by MILI with complementary piRNA sequences and undergone MILI-mediated
7 degradation in the cytoplasm, which resulted in PTGS. In the previous study, the percentage of
8 antisense *Dnmt3L* piRNAs with 10A bias was high (49%), and the peak length of the piRNA was
9 28 nt, which suggested that antisense *Dnmt3L* piRNAs were secondary piRNAs bound to MIWI2
10 (Supplemental Fig. S3B–E). MIWI2-bound piRNAs were translocated to the nucleus where they
11 bound *Dnmt3L* nascent RNA, resulting in DNA methylation of *Dnmt3L*. Considering the lack of
12 slicer activity of MIWI2, it is likely that silencing at the post-transcriptional level is determined
13 by whether the piRNA against antisense *Dnmt3L* binds to MILI. Regarding silencing at the
14 transcriptional level, the binding of piRNA against antisense *Dnmt3L* to MIWI2 as a secondary
15 piRNA is thought to be important. Taken together, although the data are circumstantial, we
16 consider that DNA methylation was not present in the current Tg mice due to the low levels of
17 MIWI2-binding antisense *Dnmt3L* piRNAs for transcriptional silencing.

18 The mice examined in this study exhibited only PTGS. In our previous study, we speculated
19 that both PTGS and TGS had occurred in Tg mice because the methylation of the endogenous
20 *Dnmt3L* promoter was partial (Itou et al. 2015). Additionally, almost two-thirds (61%) of all
21 antisense *Dnmt3L* piRNAs exhibited a 1U bias with a peak length of 27 nt (Supplemental Fig.
22 S3B–C), suggesting that these piRNAs were primary piRNAs bound to MILI. Thus, both TGS by
23 DNA methylation and PTGS by target RNA degradation play an important role in piRNA-
24 mediated silencing in the fetal testis, where transcripts of both sense and antisense strands of

- 1 retrotransposons are abundant. However, PTGS may be sufficient for silencing endogenous genes
- 2 by piRNAs on the antisense strand, as suggested in the current study.
- 3

1 **Materials and Methods**

2 ***Mice breeding***

3 All animal experiments were performed in accordance with the general guidelines of The
4 Institute of Experimental Animal Science, Osaka University Medical School. Tg-*Miwi2P-*
5 *asDnmt3L* mice were generated by injecting the *Miwi2 promoter-antisense Dnmt3L-polyA*
6 transgene cassette (*Miwi2P-asDnmt3L*) into fertilized mouse eggs (Itou et al., 2015). Transgene-
7 positive female mice were crossed with WT B6 male mice. Mice generated by crossing with WT
8 B6 mice more than five times were used in this study. Primer sequences used for genotyping are
9 listed in Table S4.

11 ***Whole-genome sequencing***

12 The genomic DNA of Tg-*Miwi2P-asDnmt3L* mice was subjected to whole-genome
13 sequencing. Paired-end sequencing of approximately 150 bases was performed using an Illumina
14 HiSeq X system (Hokkaido System Science Co., Ltd, Japan). Sequencing data were analyzed
15 using the Bowtie 2. Paired sequences consisting of one element that mapped against the *Miwi2P-*
16 *asDnmt3L* transgene and one that mapped against the genome were extracted, whereas those pairs
17 whose elements mapped against the transgene only were excluded. The extracted sequences were
18 mapped against the mouse genome. Most of these sequences were mapped to the genomic regions
19 of *Miwi2* and *Dnmt3L*. The four regions considered to be insertion sites were detected on ch18.
20 The precise insertion sites were confirmed by PCR sequencing with primers targeting the vicinity
21 of the insertion site and inside the transgene. The BioSample Accession number of DDBJ is
22 SAMD00324390.

24 ***Western blotting***

1 Fetal testes were lysed with 30 μ L of radioimmunoprecipitation buffer (50 mM Tris-HCl pH
2 8.0, 0.1% sodium dodecyl sulfate, 150 mM NaCl, 0.5% DOC, and 1% NP-40) and 10 μ L of 4 $\times$
3 sample buffer (250 mM Tris-HCl pH 6.8, 20% β -mercaptoethanol, 8% sucrose, and 0.01%
4 bromophenol blue). The lysates were boiled at 98 °C for 3 min and incubated with Benzonase
5 nuclease (Novagen) at 22 °C for 5 min. Testes samples (a half of a testis each) were boiled at
6 98 °C for 3 min and loaded onto a 10% polyacrylamide gel. The resolved proteins were transferred
7 to PVDF Immobilon-P membranes (0.45 μ m) (Millipore). The membrane was then probed with
8 either anti-DNMT3L (E1Y7Q CST#13451; Cell Signaling Technology) (1:1000) or anti- β -actin
9 (Clone AC-15, A5441; Sigma-Aldrich) (1:2500) antibodies at 4 °C overnight. Next, the
10 membranes were washed with Tris-buffered saline containing 0.1% Tween-20 (TBS-T) and
11 incubated with the secondary antibody (ECL-anti-rabbit-IgG-HRP, NA9340V; or ECL-anti-
12 mouse-IgG-HRP, NA9310V; GE Healthcare) (1:2000) for 1 h. The membranes were washed, and
13 the immunoreactive signals were detected using the BioRad Clarity Western ECL Substrate
14 #1705060 (BioRad) and a ChemiDoc MP imaging system (BioRad).

15

16 ***HE staining of paraffin-embedded sections and spermatid counting***

17 The testes were fixed with Bouin's solution overnight at 4 °C. Next, the samples were
18 dehydrated with EtOH, equalized with xylene, and embedded in paraffin. The samples were
19 sectioned and subjected to hematoxylin and eosin staining. The images of the sections were
20 captured using a BZ-X710 microscope (KEYENCE). The number of cells in the seminiferous
21 tubules was counted using the ImageJ Plugins Analyze-Cell counter. The presence of even a few
22 spermatids in the seminiferous tubule was scored as "beyond meiosis" even if there were many
23 cells arrested at meiosis during spermatogenesis. Partial spermatogenesis was scored in more than
24 200 seminiferous tubules.

1

2 *Meiotic chromosomal spread analysis*

3 The testes were freshly dissected from adult mice (7 weeks) and decapsulated. The testes were
4 ground in PBS, and the cells were collected and centrifuged. The cell pellets were washed several
5 times in PBS and suspended in hypotonic buffer (30 mM Tris at pH 7.5, 17 mM Tris sodium
6 citrate, 5 mM EDTA, and 50 mM sucrose). After keeping for 5 min, the cells were centrifuged,
7 and the supernatant was discarded, followed by making suspension in 100 mM sucrose. The cell
8 suspension was spread onto slides dipped in formaldehyde solution (1% formaldehyde, 0.15%
9 Triton X-100 in water adjusted with sodium borate to pH 9.2). Slides were then placed in a
10 humidified chamber overnight and air-dried at room temperature. For immunostaining, slides
11 were rinsed for 5 min in PBS, and then incubated for 30 min in blocking buffer (3% BSA, 10%
12 normal goat serum in PBS). The slides were incubated with primary antibodies for Scp1 (NB300-
13 229, Novus) and Scp3 (sc-74569, Santa Cruz Biotechnology) in blocking buffer overnight at 4°C,
14 followed by incubation with goat anti-mouse IgG (H+L) Alexa Fluor 568 (A11004) and goat anti-
15 rabbit IgG (H+L) Alexa Fluor 488 (A11008) (Invitrogen) secondary antibodies and DAPI for 1 h
16 at room temperature. The slides were then washed with PBS and mounted.

17

18 *Immunohistochemical staining of embryonic testes*

19 E16.5 testes were fixed by incubation with 4% paraformaldehyde (PFA) in PBS for 30 min at
20 4°C. Cryosections were permeabilized with 0.2% Triton X-100 in phosphate-buffered saline
21 (PBS) for 30 min. Sections were traversed with HistoVT One (Nacalai) at 70°C for 20 min. Slides
22 were treated with blocking buffer (3% BSA and 10% normal goat serum in PBS) for 30 min at
23 room temperature. Following treatment with primary antibodies for MILI (clone17.8, Active

motif) or MIWI2 (25D11) (Kojima-Kita et al. 2016) overnight at 4°C, the sections were incubated with secondary antibodies for 1 h at room temperature.

Germ cell sorting and bisulfite sequencing

The resected testes were treated with collagenase B (Roche) and DNase I, followed by treatment with trypsin and DNase I. The samples were washed with phosphate-buffered saline, passed through a cell strainer (FALCON #352235), and sorted using a FACS Aria II flow cytometer (BD Bioscience). The germ cells from 10-day-old mouse testes were purified by sorting with phycoerythrin (PE)-conjugated anti-mouse CD326 (Ep-CAM) antibody (BioLegend). Alternatively, germ cells from the testes of 0-day-old mice crossed with Tg-OctEGFP mice were purified by sorting with an anti-EGFP antibody. Genomic DNA was purified and treated with bisulfite using the EpiTect Fast DNA bisulfite kit (QIAGEN). To test the sorting quality, the methylation level of *Snrpn*, a maternally imprinted gene that is hypomethylated in male germ cells, was examined. The methylation level of *Snrpn* was close to 0% in all samples, which indicated high germ cell purity. The primer sequences used for amplification of the bisulfite target sites are listed in Table S7.

Northern blotting

Total RNA was prepared from 3-wk and adult testes using Isogen (Nippon Gene). Northern blotting analysis was performed as previously described in a previous study (Kuramochi-Miyagawa et al. 2008). Five micrograms of total RNA per lane were run through the gel. The probes of IAP and Line-1 were as follows: the 3' non-coding region and part of the 3'LTR region of IAP (nucleotides 6486-6793 in GenBank Accession No. M17551) and the common region of type A and Tf of Line-1 (nucleotides 2211-2625 in D84391). To compare the amount of RNA

1 loaded, the filters were stained with 0.04% methylene blue/ 0.5 M sodium acetate for 5 – 10' at
2 room temperature gently shaking after transfer and rinsed with H₂O to show the amount of 28S
3 and 18S.

5 ***Measurement of transgene copy number using qPCR***

6 To quantify the transgene copy number, qPCR was performed using a primer set targeting
7 *Dnmt3L* that can detect both the endogenous *Dnmt3L* and the antisense *Dnmt3L* Tg cassette.
8 qPCR was performed using a THUNDERBIRD™ SYBR qPCR Mix (Toyobo) and a CFX384
9 Touch™ Real-Time PCR analysis system (Bio-Rad). The expression level of *Dnmt3L* was
10 normalized to that of *mTert* (murine telomerase), which does not exhibit any sequence similarity
11 with WT genomic DNA. The copy number of *Dnmt3L* in non-Tg mice is two. Thus, the copy
12 number in the Tg mice was subtracted by two to obtain the copy number of the transgene cassette.
13 Primer sequences are listed in Table S6.

15 ***Small RNAs in fetal testes of Tg-Miwi2P-asDnmt3L***

16 Total RNA was isolated from the testes of each Tg mouse at E16.5 using ISOGEN (Nippon
17 Gene CO., LTD., Tokyo, Japan). Small RNA libraries were generated using a TruSeq small RNA
18 Library Prep kit and analyzed using an Illumina HiSeq system (Macrogen Inc.). The raw sequence
19 data in all samples were processed using CLC Genomics Workbench software (Filgen Inc.) to
20 trim miRNAs (miRbase) and tRNAs (GtRNAdb), allowing up to two mismatches after removing
21 adapter sequences, low-quality reads, and reads with lengths < 15 nt and > 45 nt. The small RNA
22 reads of length 15–45 nt were mapped to the *Miwi2P-asDnmt3L* transgene sequence, allowing up
23 to two mismatches (Table S2). This strategy is summarized in Fig. S3A.

Small RNAs derived from the genomic region around the transgene insertion sites were extracted from the fetal testis library of WT and non-Tg mice (Fig. S1D). The small RNA reads of length 15–45 nt were mapped against a 2.3-kb genomic region before and after the 18qB3 transgene insertion site (B3-3' and B3-5') and an overlapping 2.3 kb genomic region (E1) on both sides of the 18qE1 transgene.

Acknowledgments

The authors thank Ms. N. Asada for technical assistance and Ms. S. Tuda for secretarial work. This work was supported in part by a Grant-in-Aid for Scientific Research (B) (#19H03421) and Innovative Areas (#26112511) from the Japan Society for the Promotion of Science (JSPS) and AMED-CREST (#120701467) from the Japan Agency for Medical Research and Development.

Author Contributions

L.S.P. and S.K.M. performed the experiments and analyzed the data. L. S. P., S.K.M., and T.N. designed the experiments and wrote the manuscript. I.N. supported the whole-genome sequencing analysis.

Declaration of interests

The authors declare no competing interests.

Reference

- Aravin AA, Sachidanandam R, Bourc'his D, Schaefer C, Pezic D, Toth KF, Bestor T, Hannon GJ. 2008. A piRNA pathway primed by individual transposons is linked to de novo DNA methylation in mice. *Mol Cell* **31**: 785-799.
- Aravin AA, Sachidanandam R, Girard A, Fejes-Toth K, Hannon GJ. 2007. Developmentally regulated piRNA clusters implicate MILI in transposon control. *Science* **316**: 744-747.
- Brennecke J, Aravin AA, Stark A, Dus M, Kellis M, Sachidanandam R, Hannon GJ. 2007. Discrete small RNA-generating loci as master regulators of transposon activity in *Drosophila*. *Cell* **128**: 1089-1103.
- Carmell MA, Girard A, van de Kant HJ, Bourc'his D, Bestor TH, de Rooij DG, Hannon GJ. 2007. MIWI2 is essential for spermatogenesis and repression of transposons in the mouse male germline. *Dev Cell* **12**: 503-514.
- Castaneda J, Genzor P, Bortvin A. 2011. piRNAs, transposon silencing, and germline genome integrity. *Mutat Res* **714**: 95-104.
- Chuma S, Nakano T. 2013. piRNA and spermatogenesis in mice. *Philos Trans R Soc Lond B Biol Sci* **368**: 20110338.
- Czech B, Hannon GJ. 2016. One Loop to Rule Them All: The Ping-Pong Cycle and piRNA-Guided Silencing. *Trends Biochem Sci* **41**: 324-337.
- De Fazio S, Bartonicek N, Di Giacomo M, Abreu-Goodger C, Sankar A, Funaya C, Antony C, Moreira PN, Enright AJ, O'Carroll D. 2011. The endonuclease activity of Mili fuels piRNA amplification that silences LINE1 elements. *Nature* **480**: 259-263.
- Deng W, Lin H. 2002. miwi, a murine homolog of piwi, encodes a cytoplasmic protein essential for spermatogenesis. *Dev Cell* **2**: 819-830.
- Di Giacomo M, Comazzetto S, Saini H, De Fazio S, Carrieri C, Morgan M, Vasiliauskaite L, Benes V, Enright AJ, O'Carroll D. 2013. Multiple epigenetic mechanisms and the piRNA pathway enforce LINE1 silencing during adult spermatogenesis. *Mol Cell* **50**: 601-608.
- Girard A, Sachidanandam R, Hannon GJ, Carmell MA. 2006. A germline-specific class of small RNAs binds mammalian Piwi proteins. *Nature* **442**: 199-202.
- Hajkova P, Erhardt S, Lane N, Haaf T, El-Maarri O, Reik W, Walter J, Surani MA. 2002. Epigenetic reprogramming in mouse primordial germ cells. *Mech Dev* **117**: 15-23.
- Itou D, Shiromoto Y, Yukiho SY, Ishii C, Nishimura T, Ogonuki N, Ogura A, Hasuwa H, Fujihara Y, Kuramochi-Miyagawa S et al. 2015. Induction of DNA methylation by artificial piRNA production in male germ cells. *Curr Biol* **25**: 901-906.
- Kanatsu-Shinohara M, Takashima S, Ishii K, Shinohara T. 2011. Dynamic changes in EPCAM expression during spermatogonial stem cell differentiation in the mouse testis. *PLoS One* **6**: e23663.

- 1 Kato Y, Kaneda M, Hata K, Kumaki K, Hisano M, Kohara Y, Okano M, Li E, Nozaki M, Sasaki
2 H. 2007. Role of the Dnmt3 family in de novo methylation of imprinted and repetitive
3 sequences during male germ cell development in the mouse. *Hum Mol Genet* **16**: 2272-
4 2280.
- 5 Kojima-Kita, K., Kuramochi-Miyagawa, S., Nagamori, I., Ogonuki, N., Ogura, A., Hasuwa, H.,
6 Akazawa, T., Inoue, N., and Nakano, T. 2016. MIWI2 as an Effector of DNA Methylation
7 and Gene Silencing in Embryonic Male Germ Cells. *Cell Rep* **16**: 2819-2828.
- 8 Kuramochi-Miyagawa S, Kimura T, Ijiri TW, Isobe T, Asada N, Fujita Y, Ikawa M, Iwai N, Okabe
9 M, Deng W et al. 2004. Mili, a mammalian member of piwi family gene, is essential for
10 spermatogenesis. *Development* **131**: 839-849.
- 11 Kuramochi-Miyagawa S, Kimura T, Yomogida K, Kuroiwa A, Tadokoro Y, Fujita Y, Sato M,
12 Matsuda Y, Nakano T. 2001. Two mouse piwi-related genes: miwi and mili. *Mech Dev*
13 **108**: 121-133.
- 14 Kuramochi-Miyagawa S, Nakano T. 2015. Reply to Shoji and Katsuma. *Curr Biol* **25**: R710.
- 15 Kuramochi-Miyagawa S, Watanabe T, Gotoh K, Totoki Y, Toyoda A, Ikawa M, Asada N, Kojima
16 K, Yamaguchi Y, Ijiri TW et al. 2008. DNA methylation of retrotransposon genes is
17 regulated by Piwi family members MILI and MIWI2 in murine fetal testes. *Genes Dev*
18 **22**: 908-917.
- 19 La Salle S, Oakes CC, Neaga OR, Bourc'his D, Bestor TH, Trasler JM. 2007. Loss of
20 spermatogonia and wide-spread DNA methylation defects in newborn male mice
21 deficient in DNMT3L. *BMC Dev Biol* **7**: 104.
- 22 Li JY, Lees-Murdock DJ, Xu GL, Walsh CP. 2004. Timing of establishment of paternal
23 methylation imprints in the mouse. *Genomics* **84**: 952-960.
- 24 Manakov SA, Pezic D, Marinov GK, Pastor WA, Sachidanandam R, Aravin AA. 2015. MIWI2
25 and MILI Have Differential Effects on piRNA Biogenesis and DNA Methylation. *Cell*
26 *Rep* **12**: 1234-1243.
- 27 Molaro A, Falciatori I, Hodges E, Aravin AA, Marran K, Rafii S, McCombie WR, Smith AD,
28 Hannon GJ. 2014. Two waves of de novo methylation during mouse germ cell
29 development. *Genes Dev* **28**: 1544-1549.
- 30 Mouse Genome Sequencing C Waterston RH Lindblad-Toh K Birney E Rogers J Abril JF Agarwal
31 P Agarwala R Ainscough R Alexandersson M et al. 2002. Initial sequencing and
32 comparative analysis of the mouse genome. *Nature* **420**: 520-562.
- 33 Pezic D, Manakov SA, Sachidanandam R, Aravin AA. 2014. piRNA pathway targets active
34 LINE1 elements to establish the repressive H3K9me3 mark in germ cells. *Genes Dev* **28**:
35 1410-1428.
- 36 Reuter M, Berninger P, Chuma S, Shah H, Hosokawa M, Funaya C, Antony C, Sachidanandam

- 1 R, Pillai RS. 2011. Miwi catalysis is required for piRNA amplification-independent
2 LINE1 transposon silencing. *Nature* **480**: 264-267.
- 3 Sarkar A, Maji RK, Saha S, Ghosh Z. 2014. piRNAQuest: searching the piRNAome for silencers.
4 *BMC Genomics* **15**: 555.
- 5 Sasaki H, Matsui Y. 2008. Epigenetic events in mammalian germ-cell development:
6 reprogramming and beyond. *Nat Rev Genet* **9**: 129-140.
- 7 Seisenberger S, Andrews S, Krueger F, Arand J, Walter J, Santos F, Popp C, Thienpont B, Dean
8 W, Reik W. 2012. The dynamics of genome-wide DNA methylation reprogramming in
9 mouse primordial germ cells. *Mol Cell* **48**: 849-862.
- 10 Senti KA, Jurczak D, Sachidanandam R, Brennecke J. 2015. piRNA-guided slicing of transposon
11 transcripts enforces their transcriptional silencing via specifying the nuclear piRNA
12 repertoire. *Genes Dev* **29**: 1747-1762.
- 13 Shoji K, Katsuma S. 2015. Is the expression of sense and antisense transgenes really sufficient
14 for artificial piRNA production? *Curr Biol* **25**: R708-710.
- 15 Toth KF, Pezic D, Stuwe E, Webster A. 2016. The piRNA Pathway Guards the Germline Genome
16 Against Transposable Elements. *Adv Exp Med Biol* **886**: 51-77.
- 17 Watanabe T, Tomizawa S, Mitsuya K, Totoki Y, Yamamoto Y, Kuramochi-Miyagawa S, Iida N,
18 Hoki Y, Murphy PJ, Toyoda A et al. 2011. Role for piRNAs and noncoding RNA in de
19 novo DNA methylation of the imprinted mouse Rasgrfl locus. *Science* **332**: 848-852.
- 20 Zanni V, Eymery A, Coiffet M, Zytnicki M, Luyten I, Quesneville H, Vaury C, Jensen S. 2013.
21 Distribution, evolution, and diversity of retrotransposons at the flamenco locus reflect the
22 regulatory properties of piRNA clusters. *Proc Natl Acad Sci USA* **110**: 19842-19847.
- 23 Zhang P, Si X, Skogerbo G, Wang J, Cui D, Li Y, Sun X, Liu L, Sun B, Chen R et al. 2014.
24 piRBase: a web resource assisting piRNA functional study. *Database (Oxford)* **2014**:
25 bau110.
- 26 Zheng K, Wang PJ. 2012. Blockade of pachytene piRNA biogenesis reveals a novel requirement
27 for maintaining post-meiotic germline genome integrity. *PLoS Genet* **8**: e1003038.

28

29

1 **Figure legends**

2 **Fig 1. Transgene insertion sites on chromosome 18 (ch18) of the mouse genome**

3 (A) The four transgene insertion sites found on ch18 and the sequences of the junctions are shown.
 4 The numbers represent the location on the genome (NCBI/mm10 assembly). Genomic sequences
 5 and unknown sequences are shown in black and green letters, respectively. Transgene sequences
 6 are highlighted in blue. (B and C) Read counts around the insertion loci were determined using
 7 whole-genome sequencing. The overall lengths of (B) and (C) are 300 kb for Ch18qB3 and 10 kb
 8 for Ch18qE1, respectively. The read counts show a 100-bp window. The regions flanked by the
 9 insertion sites of the transgenic cassette in both B3 and E1 regions have a high genomic read
 10 count (red arrow). This indicates a model in which the transgene is sandwiched between
 11 duplicated genomic regions.

13 **Fig 2. Transgenic mice (Tg) phenotypes**

14 (A) Schematic illustration of polymerase chain reaction (PCR) primer annealing sites to confirm
 15 Tg cassette insertion sites. “Tg internal” legends indicate the junctions between *Miwi2* promoter
 16 and *asDnmt3L*. “Tg head” shows the junction between genomic DNA and the *Miwi2* promoter at
 17 Ch18qB3 and E1 sites, and “Tg tail” shows the junction between the tail of the Tg cassette and
 18 genomic DNA at each insertion site. The primer sequences are shown in Table S4. The PCR
 19 products were subjected to agarose gel electrophoresis with Tris-acetate EDTA buffer (right
 20 panel). (B) Morphology of testes of adult mice with transgenes inserted at the E1 locus (Tg-E),
 21 B3 locus (Tg-B), and B3 and E1 loci (Tg-BE). Black bar indicates a 2 mm scale. (C) Weights of
 22 the testes of Tg-E, Tg-B, and Tg-BE mice. Each dot represents the weight of each testis. The
 23 range of the whisker lines represents the minimum and maximum values; outliers were excluded.
 24 The box ranges indicate the interquartile range (IQR), the horizontal bars in the box indicate the

median, and the “X” marks indicate the mean weights. (98.4, 90.8, 32.0, and 29.8 mg in non-Tg, Tg-E, Tg-B, and Tg-BE mice, respectively; T-test; * $p = 0.15$; ** $p < 0.01$; *** $p = 0.72$). (D) DNMT3L expression in mice testes at E16.5 was analyzed using western blotting. Upper panel shows DNMT3L and lower panel shows β -actin. Each lane was loaded with protein extracts from one half of a testis. The band intensity of DNMT3L was normalized to that of β -actin, as well as to that of DNMT3L in non-Tg mice. Analysis was performed using a Gel analyzer 2010. (E) Hematoxylin and eosin-stained histological sections showing developmental differences between non-Tg, Tg-E, Tg-B, and Tg-BE mouse testes.

Fig 3. Bisulfite sequencing and Northern blot of DNMT3L *de novo* DNA methylation targets

(A) Methylation levels of retrotransposon genes (*L1A*, *LINE1* type A; *L1Tf*, *LINE1* Tf; and *IAP*, *IAP1Δ1*) in germ cells of non-Tg and mice with transgenes inserted at the E1 locus (Tg-E), B3 locus (Tg-B), and B3 and E1 loci (Tg-BE). (B) Methylation levels of paternally imprinted genes (*Rasgrfl*, *H19*, and *Dlk1*). (C) Methylation levels of *Snrpn*, a maternally imprinted gene, representing sorted cell purity. Black and white circles indicate methylated and unmethylated CpGs, respectively. (D) Northern blotting of *LINE1* and *IAP* in testes of 3 weeks in the non-Tg and Tg mice. To compare the amount of RNA loaded, the filters were stained with methylene blue after transfer to show the amount of 28S and 18S.

Fig 4. Copy numbers of as*Dnmt3L* transgenes and their correlation with phenotypes

(A) as*Dnmt3L* transgene copy numbers in mice with transgenes inserted at the E1 locus (Tg-E), B3 locus (Tg-B), and B3 and E1 loci (Tg-BE) were confirmed using quantitative polymerase chain reaction. Each dot indicates the copy number. The range of the whisker lines indicates the minimum and maximum values excluding outliers. The box range indicates the interquartile range

1 (IQR), the horizontal bars in the box indicate the median, and the “X” marks indicate the mean
 2 value. (10, 38, and 42 copies in Tg-E, Tg-B, and Tg-BE mice, respectively). (B) Transgene copy
 3 numbers represented through a bar graph. (C) Percentage of cells in seminiferous tubules that
 4 have undergone meiosis. The blue and red bars represent the cells that have completed meiosis
 5 and those that are arrested at meiosis, respectively. (D) Progression of spermatogenesis in mice
 6 with single or double alleles of the transgene cassette inserted at the E1 locus (Tg-E and Tg-E/E
 7 mice, respectively). Each dot represents the percentage of cells in the tubules of a mouse that have
 8 completed meiosis. The range of the whisker lines indicates the minimum and maximum values
 9 excluding outliers. The box range indicates the interquartile range (IQR), the horizontal bars in
 10 the box indicate the median, and the “X” marks indicate the mean value. (97% and 72% in cells
 11 with single and double transgenic alleles, respectively). Student’s *t*-test; * $p < 0.01$. (E) Testicular
 12 weight relative to the body weight of mice with single or double transgenic alleles. Each dot
 13 represents testis weight relative to body weight (mg/g) and horizontal bars represent averages (3.2
 14 and 1.8 mg/g for mice with single and double transgenic alleles, respectively). Student’s *t*-test; *
 15 $p < 0.01$. (F–G) Correlation between *Dnmt3L* copy number and the progression of
 16 spermatogenesis (F) or testicular weight relative to body weight (G). Black, yellow, orange, blue,
 17 and green dots indicate cells derived from non-Tg mice, Tg-E mice, mice with double transgenic
 18 alleles (Tg-E/E), Tg-B mice, and Tg-BE mice, respectively. The fitting curves were adjusted to a
 19 sigmoidal model (Min = 6.03, Max = 95.81, half-maximal effective concentration [EC50] = 23.72,
 20 $R^2 = 0.98$ for (F); and Min = 0.82, Max = 3.25, EC50 = 20.5, $R^2 = 0.94$ for (G)).

21

22 **Fig 5. Small RNAs from *Miwi2P-asDnmt3L* in fetal testes**

23 (A) Numbers of small RNAs annotated as from the *Miwi2P-asDnmt3L* transgene among total
 24 reads in the testes of non-transgenic (Tg) mice and mice with transgene cassettes inserted at the

1 E1 locus (Tg-E), B3 locus (Tg-B), and B3 and E1 loci (Tg-BE) at E16.5. Blue bars show the small
 2 RNAs from the sense *Miwi2* promoter and antisense *Dnmt3L* (transgene sense strand), while the
 3 orange bars show the small RNAs from the antisense *Miwi2* promoter and sense *Dnmt3L*
 4 (transgene antisense strand). (B) Size distribution of transgene-annotated small RNAs. The upper
 5 side represents the transgene sense strand where blue and purple indicate small RNAs with and
 6 without uracil as their first nucleotide, respectively. The lower side represents the transgene
 7 antisense strand where orange and purple indicate small RNAs with and without adenine at the
 8 tenth position, respectively. (C) Schematic diagram showing the directions of small RNAs. Red
 9 arrow indicates the expected transcript of as*Dnmt3L* driven by the *Miwi2* promoter. Light blue,
 10 blue, and gray arrows indicate transgene sense strands, while yellow, orange, and brown arrows
 11 indicate transgene antisense strands. (D) Ratio of transgene-annotated small RNAs in the fetal
 12 testis of Tg mice. The colors of the pie charts correspond to the colors of the arrows in Fig. 5C.
 13 The shaded orange color contains piRNAs from the transgene and the endogenous *Dnmt3L*. (E)
 14 Mapping of piRNAs of the Tg-B mice onto the transgene cassette sequence. The upper and the
 15 lower sides represent the sense strand in blue and the antisense strand in orange, respectively.

16

17 **Fig 6. Bisulfite sequencing of *Dnmt3L* promoter**

18 The top panel shows a diagram of the *Dnmt3L* promoter and its methylation positions (white
 19 circles). Methylation levels of the *Dnmt3L* promoter in germ cells of 0-day-old (P0) non-
 20 transgenic (Non-Tg) mice and of mice with a transgene cassette inserted at the B3 locus (Tg-B)
 21 as well as in those of 10-day-old (P10) Non-Tg mice and mice with transgene cassettes inserted
 22 at the E1 locus (Tg-E) and at both the E1 and B3 loci (Tg-BE). Black and white circles indicate
 23 methylated and unmethylated CpGs, respectively.

(A)

(B)

(C)

(D)

(A)

(B)

(C)

(E)

(D)

Fig.6 Li et.al.

RNA

A PUBLICATION OF THE RNA SOCIETY

Effects of transgene insertion loci and copy number on Dnmt3L gene silencing through antisense transgene-derived PIWI-interacting RNAs

SePil Lee, Satomi Kuramochi-Miyagawa, Ippei Nagamori, et al.

RNA published online February 10, 2022

Supplemental Material

<http://rnajournal.cshlp.org/content/suppl/2022/02/10/rna.078905.121.DC1>

P<P

Published online February 10, 2022 in advance of the print journal.

Accepted Manuscript

Peer-reviewed and accepted for publication but not copyedited or typeset; accepted manuscript is likely to differ from the final, published version.

Open Access

Freely available online through the *RNA* Open Access option.

Creative Commons License

This article, published in *RNA*, is available under a Creative Commons License (Attribution-NonCommercial 4.0 International), as described at <http://creativecommons.org/licenses/by-nc/4.0/>.

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).

To subscribe to *RNA* go to:
<http://rnajournal.cshlp.org/subscriptions>