

RNA

A PUBLICATION OF THE RNA SOCIETY

VOL. 15, NO. 11



November 2009

CONTENTS

Mini-Review

- Domain-elongation NMR spectroscopy yields new insights into RNA dynamics and adaptive recognition 1941
Qi Zhang and Hashim M. Al-Hashimi

Review

- Three-way RNA junctions with remote tertiary contacts: A recurrent and highly versatile fold 1949
Marcos de la Peña, David Dufour, and José Gallego

Letters to the Editor

- Distinct extremely abundant siRNAs associated with cosuppression in petunia 1965^{OA}
Emanuele De Paoli, Ana Dorantes-Acosta, Jixian Zhai, Monica Accerbi, Dong-Hoon Jeong, Sunhee Park, Blake C. Meyers, Richard A. Jorgensen, and Pamela J. Green
- Combined proteomic–RNAi screen for host factors involved in human hepatitis delta virus replication 1971
Dan Cao, Dirk Haussecker, Yong Huang, and Mark A. Kay
- The XIAP IRES activates 3' cistron expression by inducing production of monocistronic mRNA in the β gal/CAT bicistronic reporter system 1980
Holly A. Saffran and James R. Smiley

Report

- Slow formation of a pseudoknot structure is rate limiting in the productive co-transcriptional folding of the self-splicing *Candida* intron 1986
Libin Zhang, Penghui Bao, Michael J. Leibowitz, and Yi Zhang

Articles

- The *Thermus thermophilus* DEAD box helicase Hera contains a modified RNA recognition motif domain loosely connected to the helicase core 1993
Markus G. Rudolph and Dagmar Klostermeier

(continued)

Cover Illustration: Crystal structure of the *Pyrococcus abyssi* exosome RNase PH complexed with a single-stranded 10-mer poly(A) RNA (Protein Data Bank code: 2po1; Navarro MVAS, Oliveira CC, Zanchin NIT, Guimaraes BG. 2008. Insights into the mechanism of progressive RNA degradation by the archaeal exosome. *J Biol Chem* **283**: 14120–14131.) Image details show RNA: surface representation, coloring according to the molecular electrostatic potential; proteins: cartoon representation, Rrp41 subunit—green, Rrp42 subunit—orange. The image was generated with the Accelrys Discovery Studio Visualizer. Cover image provided by the Jena Library of Biological Macromolecules—JenaLib (www.fli-leibniz.de/IMAGE.html).

Formation of She2p tetramers is required for mRNA binding, mRNP assembly, and localization <i>Marisa Müller, Klaus Richter, Alexander Heuck, Elisabeth Kremmer, Johannes Buchner, Ralf-Peter Jansen, and Dierk Niessing</i>	2002 ^{OA}
Complex architecture and regulated expression of the <i>Sox2ot</i> locus during vertebrate development <i>Paulo P. Amaral, Christine Neyt, Simon J. Wilkins, Marjan E. Askarian-Amiri, Susan M. Sunkin, Andrew C. Perkins, and John S. Mattick</i>	2013
Quantitative miRNA expression analysis: Comparing microarrays with next-generation sequencing <i>Hanni Willenbrock, Jesper Salomon, Rolf Søkilde, Kim Bundvig Barken, Thomas Nøhr Hansen, Finn Cilius Nielsen, Søren Møller, and Thomas Litman</i>	2028 ^{OA}
Direct selection for ribozyme cleavage activity in cells <i>Xi Chen, Lisa Denison, Matthew Levy, and Andrew D. Ellington</i>	2035 ^{OA}
A variant riboswitch aptamer class for S-adenosylmethionine common in marine bacteria <i>Elena Poiata, Michelle M. Meyer, Tyler D. Ames, and Ronald R. Breaker</i>	2046 ^{OA}
Methods	
Real-time fluorescence detection of exoribonucleases <i>Flore Sinturel, Olivier Pellegrini, Song Xiang, Liang Tong, Ciarán Condon, and Lionel Bénard</i>	2057
In vivo detection of RNA-binding protein interactions with cognate RNA sequences by fluorescence resonance energy transfer <i>Martina Huranová, Joseph A. Jablonski, Aleš Benda, Martin Hof, David Staněk, and Massimo Caputi</i>	2063
Erratum	2072
RNA: Instructions for contributors	2073

^{OA}Open Access paper