

RNA

A PUBLICATION OF THE RNA SOCIETY

VOL. 15, NO. 6



June 2009

CONTENTS

Letter to the Editor

- Post-transcriptional control of DGCR8 expression by the Microprocessor 1005
Robinson Triboulet, Hao-Ming Chang, Robert J. LaPierre, and Richard I. Gregory

Bioinformatics

- Small RNA analysis in *Petunia hybrida* identifies unusual tissue-specific expression patterns of conserved miRNAs and of a 24mer RNA 1012^{OA}
Philip Tedder, Elena Zubko, David R. Westhead, and Peter Meyer

Reports

- Equal transcription rates of productively and nonproductively rearranged immunoglobulin μ heavy chain alleles in a pro-B cell line 1021
Andrea B. Eberle, Kai Herrmann, Hans-Martin Jäck, and Oliver Mühlemann
- Co-occupancy of two Pumilio molecules on a single hunchback NRE 1029
Yogesh K. Gupta, Tammy H. Lee, Thomas A. Edwards, Carlos R. Escalante, Lyudmila Y. Kadyrova, Robin P. Wharton, and Aneel K. Aggarwal
- Structure–activity relationships in *Kluyveromyces lactis* γ -toxin, a eukaryal tRNA anticodon nuclease 1036
Niroshika Keppetipola, Ruchi Jain, Birthe Meineke, Melinda Diver, and Stewart Shuman
- Regulation of NAB2 mRNA 3'-end formation requires the core exosome and the Trf4p component of the TRAMP complex 1045
Kelly M. Roth, Joel Byam, Feng Fang, and J. Scott Butler
- The C-terminal domains of human TNRC6A, TNRC6B, and TNRC6C silence bound transcripts independently of Argonaute proteins 1059^{OA}
Daniela Lazzaretti, Isabelle Tournier, and Elisa Izaurralde

(continued)

Cover Illustration: Crystal structure of the N-terminal domain of human La protein complexed with the RNA oligomer AUUU (Protein Data Bank code: 2vop; Kotik-Kogan, O., Valentine, E.R., Sanfelice, D., Conte, M.R., and Curry, S. 2008. Structural analysis reveals conformational plasticity in the recognition of RNA 3' ends by the human La protein. *Structure* **16**: 852–862.) Image details show protein: cartoon representation, coloring according to secondary structure; RNA: ribbon–plate representation, transparent surface, A—red, U—magenta, G—green. C—blue. The image was generated with the Accelrys Discovery Studio Visualizer. Cover image provided by the Jena Library of Biological Macromolecules—JenaLib (www.fli-leibniz.de/IMAGE.html).

Articles

- A C-terminal silencing domain in GW182 is essential for miRNA function 1067^{OA}
Ana Eulalio, Sigrun Helms, Christoph Fritsch, Maria Fauser, and Elisa Izaurralde
- Mammalian GW182 contains multiple Argonaute-binding sites and functions in microRNA-mediated translational repression 1078
Koji Takimoto, Motoaki Wakiyama, and Shigeyuki Yokoyama
- A single *C. elegans* PUF protein binds RNA in multiple modes 1090
Yvonne Yiling Koh, Laura Opperman, Craig Stumpf, Arpita Mandan, Sunduz Keles, and Marvin Wickens
- Accuracy modulating mutations of the ribosomal protein S4-S5 interface do not necessarily destabilize the rps4-rps5 protein–protein interaction 1100
Haritha Vallabhaneni and Philip J. Farabaugh
- Specific and global regulation of mRNA stability during osmotic stress in *Saccharomyces cerevisiae* 1110
Lorena Romero-Santacreu, Joaquín Moreno, José E. Pérez-Ortín, and Paula Alepuz
- Vg1RBP phosphorylation by Erk2 MAP kinase correlates with the cortical release of Vg1 mRNA during meiotic maturation of *Xenopus* oocytes 1121
Anna Git, Rachel Allison, Eusebio Perdiguer, Angel R. Nebreda, Evelyn Houliston, and Nancy Standart
- The *yfiC* gene of *E. coli* encodes an adenine-N6 methyltransferase that specifically modifies A37 of tRNA^{Val} (cmo⁵UAC) 1134
Anna Y. Golovina, Petr V. Sergiev, Andrey V. Golovin, Marina V. Serebryakova, Irina Demina, Vadim M. Govorun, and Olga A. Dontsova
- A comparative genomics approach identifies a PPR-DYW protein that is essential for C-to-U editing of the *Arabidopsis* chloroplast *accD* transcript 1142
John C. Robbins, Wade P. Heller, and Maureen R. Hanson
- Gene regulation by sense–antisense overlap of polyadenylation signals 1154
Rui Gu, Zuo Zhang, Joshua N. DeCervo, and Gordon G. Carmichael
- Cold-inducible RNA binding protein (CIRP) expression is modulated by alternative mRNAs 1164
Mohamed B. Al-Fageeh and C. Mark Smales
- The fraction of RNA that folds into the correct branched secondary structure determines hepatitis delta virus type 3 RNA editing levels 1177
Sarah D. Linnstaedt, Wojciech K. Kasprzak, Bruce A. Shapiro, and John L. Casey
- SHQ1 is required prior to NAF1 for assembly of H/ACA small nucleolar and telomerase RNPs 1188
Petar N. Grozdanov, Sujayita Roy, Nupur Kittur, and U. Thomas Meier
- The conserved 3' end domain of U6atac snRNA can direct U6 snRNA to the minor spliceosome 1198
Rosemary C. Dietrich, Richard A. Padgett, and Girish C. Shukla
- An extra double-stranded RNA binding domain confers high activity to a squid RNA editing enzyme 1208
Juan Pablo Palavicini, Mary A. O'Connell, and Joshua J.C. Rosenthal

Method

- DOCK 6: Combining techniques to model RNA–small molecule complexes 1219
P. Therese Lang, Scott R. Brozell, Sudipto Mukherjee, Eric F. Pettersen, Elaine C. Meng, Veena Thomas, Robert C. Rizzo, David A. Case, Thomas L. James, and Irwin D. Kuntz

RNA: Instructions for contributors